Full wwPDB NMR Structure Validation Report (i)

Feb 12, 2022 – 05:17 PM EST

PDB ID	:	1FYO
Title	:	EUKARYOTIC DECODING REGION A-SITE RNA
Authors	:	Lynch, S.R.; Puglisi, J.D.
Deposited on	:	2000-10-02

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/NMRValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity	:	4.02b-467
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
RCI	:	v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV	:	Wang et al. (2010)
ShiftChecker	:	2.26
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.26

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
SOLUTION NMR

The overall completeness of chemical shifts assignment was not calculated.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	158937	12864
RNA backbone	4643	676

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Mol	Chain	Length	Quality of chain
1	A	27	30% 59% 11%

2 Ensemble composition and analysis (i)

This entry contains 25 models. This entry does not contain polypeptide chains, therefore identification of well-defined residues and clustering analysis are not possible. All residues are included in the validation scores.

3 Entry composition (i)

There is only 1 type of molecule in this entry. The entry contains 867 atoms, of which 294 are hydrogens and 0 are deuteriums.

- Molecule 1 is a RNA chain called FRAGMENT OF 18S RIBOSOMAL RNA.

Mol	Chain	Residues	Atoms						Trace
			Total	C	H	N	O	P	
1	A	27	867	256	294	102	189	26	0

4 Residue-property plots (i)

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

• Molecule 1: FRAGMENT OF 18S RIBOSOMAL RNA

Chain A: 

4.2 Scores per residue for each member of the ensemble

Colouring as in section 4.1 above.

4.2.1 Score per residue for model 1

• Molecule 1: FRAGMENT OF 18S RIBOSOMAL RNA

Chain A: 

4.2.2 Score per residue for model 2

• Molecule 1: FRAGMENT OF 18S RIBOSOMAL RNA

Chain A: 



4.2.3 Score per residue for model 3

• Molecule 1: FRAGMENT OF 18S RIBOSOMAL RNA

Chain A: 19% 74% 7%

G1 G2 C3 G4 U5 C6 G7 C8 A9 C10 C11 U12 U13 C14 G15 G16 G17 U18 G19 A20 A21 G22 U23 C24 C27

4.2.4 Score per residue for model 4

• Molecule 1: FRAGMENT OF 18S RIBOSOMAL RNA

Chain A: 52% 37% 11%

G1 G2 U5 C6 G7 C11 U12 U13 C14 G15 G19 A20 A21 G22 C27

4.2.5 Score per residue for model 5

• Molecule 1: FRAGMENT OF 18S RIBOSOMAL RNA

Chain A: 37% 56% 7%

G1 G2 U5 C6 C11 U12 U13 C14 G15 G16 G17 U18 G19 A20 A21 G22 U23 C24 C27

4.2.6 Score per residue for model 6

• Molecule 1: FRAGMENT OF 18S RIBOSOMAL RNA

Chain A: 37% 52% 11%

G1 G2 C3 G4 U5 C6 C11 U12 U13 C14 G15 G16 G17 U18 G19 A20 A21 G22 U23 C27

4.2.7 Score per residue for model 7

• Molecule 1: FRAGMENT OF 18S RIBOSOMAL RNA

Chain A: 26% 67% 7%

G1 G2 U5 C6 G7 C8 A9 C10 C11 U12 U13 C14 G15 G16 G17 U18 G19 A20 A21 G22 U23 C24 C27

4.2.8 Score per residue for model 8

- Molecule 1: FRAGMENT OF 18S RIBOSOMAL RNA

Chain A: 22% 63% 15%

4.2.9 Score per residue for model 9

- Molecule 1: FRAGMENT OF 18S RIBOSOMAL RNA

Chain A: 44% 44% 11%

4.2.10 Score per residue for model 10

- Molecule 1: FRAGMENT OF 18S RIBOSOMAL RNA

Chain A: 44% 44% 11%



4.2.11 Score per residue for model 11

- Molecule 1: FRAGMENT OF 18S RIBOSOMAL RNA

Chain A: 33% 56% 11%

4.2.12 Score per residue for model 12

- Molecule 1: FRAGMENT OF 18S RIBOSOMAL RNA

Chain A: 26% 63% 11%



4.2.13 Score per residue for model 13

• Molecule 1: FRAGMENT OF 18S RIBOSOMAL RNA

Chain A: 26% 63% 11%



4.2.14 Score per residue for model 14

• Molecule 1: FRAGMENT OF 18S RIBOSOMAL RNA

Chain A: 37% 56% 7%

4.2.15 Score per residue for model 15

• Molecule 1: FRAGMENT OF 18S RIBOSOMAL RNA

Chain A: 30% 63% 7%



4.2.16 Score per residue for model 16

• Molecule 1: FRAGMENT OF 18S RIBOSOMAL RNA

Chain A: 33% 59% 7%



4.2.17 Score per residue for model 17

• Molecule 1: FRAGMENT OF 18S RIBOSOMAL RNA

Chain A: 37% 52% 11%

4.2.18 Score per residue for model 18

- Molecule 1: FRAGMENT OF 18S RIBOSOMAL RNA

Chain A: 26% 67% 7%



4.2.19 Score per residue for model 19

- Molecule 1: FRAGMENT OF 18S RIBOSOMAL RNA

Chain A: 33% 56% 11%

4.2.20 Score per residue for model 20

- Molecule 1: FRAGMENT OF 18S RIBOSOMAL RNA

Chain A: 44% 48% 7%



4.2.21 Score per residue for model 21

- Molecule 1: FRAGMENT OF 18S RIBOSOMAL RNA

Chain A: 22% 67% 11%



4.2.22 Score per residue for model 22

- Molecule 1: FRAGMENT OF 18S RIBOSOMAL RNA

Chain A: 41% 48% 11%



4.2.23 Score per residue for model 23

- Molecule 1: FRAGMENT OF 18S RIBOSOMAL RNA

Chain A: 33% 56% 11%



4.2.24 Score per residue for model 24

- Molecule 1: FRAGMENT OF 18S RIBOSOMAL RNA

Chain A: 48% 41% 11%



4.2.25 Score per residue for model 25

- Molecule 1: FRAGMENT OF 18S RIBOSOMAL RNA

Chain A: 37% 44% 19%



5 Refinement protocol and experimental data overview (i)

The models were refined using the following method: *simulated annealing molecular dynamics.*

Of the 100 calculated structures, 25 were deposited, based on the following criterion: *structures with the least restraint violations,structures with the lowest energy.*

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
X-PLOR	refinement	3.83
X-PLOR	structure solution	3.83

No chemical shift data was provided.

6 Model quality (i)

6.1 Standard geometry (i)

There are no covalent bond-length or bond-angle outliers.

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

There are no planarity outliers.

6.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	573	294	294	17±2
All	All	14325	7350	7350	430

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 20.

All unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:20:A:O2'	1:A:21:A:H5"	1.05	1.51	7	25
1:A:20:A:HO2'	1:A:21:A:H8	0.93	1.02	7	20
1:A:21:A:HO2'	1:A:22:G:H8	0.92	1.04	1	19
1:A:21:A:O2'	1:A:22:G:H8	0.76	1.63	24	23
1:A:15:G:H3'	1:A:15:G:N3	0.75	1.96	25	8
1:A:20:A:O2'	1:A:21:A:H8	0.67	1.71	25	25
1:A:20:A:H3'	1:A:20:A:OP2	0.64	1.92	13	8
1:A:19:G:N2	1:A:20:A:N7	0.64	2.46	21	7
1:A:14:C:H4'	1:A:15:G:O5'	0.61	1.96	3	24
1:A:17:G:O2'	1:A:18:U:H5'	0.60	1.97	1	16
1:A:20:A:O2'	1:A:21:A:C8	0.60	2.54	6	9
1:A:11:C:H2'	1:A:12:U:O4'	0.60	1.96	19	12
1:A:6:C:H2'	1:A:7:G:O4'	0.59	1.98	3	11
1:A:20:A:O2'	1:A:21:A:C5'	0.58	2.45	6	7

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:11:C:O2'	1:A:12:U:H5'	0.58	1.99	14	15
1:A:5:U:H2'	1:A:6:C:C6	0.57	2.34	25	22
1:A:13:U:O2'	1:A:14:C:H5'	0.57	2.00	9	24
1:A:21:A:O2'	1:A:22:G:C8	0.56	2.59	3	4
1:A:1:G:O2'	1:A:2:G:H5'	0.56	2.01	16	23
1:A:19:G:C2	1:A:20:A:N7	0.56	2.74	5	21
1:A:20:A:HO2'	1:A:21:A:H5"	0.55	1.56	17	6
1:A:18:U:O2'	1:A:19:G:H5'	0.54	2.02	7	3
1:A:9:A:O2'	1:A:10:C:H5'	0.54	2.02	7	6
1:A:23:U:H2'	1:A:24:C:O4'	0.53	2.03	8	12
1:A:21:A:C2	1:A:22:G:C5	0.52	2.98	3	2
1:A:15:G:N3	1:A:15:G:C3'	0.52	2.70	25	1
1:A:22:G:H4'	1:A:22:G:OP1	0.50	2.05	20	3
1:A:22:G:O2'	1:A:23:U:H5'	0.50	2.06	15	1
1:A:8:C:C2	1:A:20:A:N6	0.50	2.79	18	1
1:A:16:G:O2'	1:A:17:G:H5'	0.50	2.07	16	3
1:A:8:C:C2	1:A:21:A:N6	0.49	2.80	8	2
1:A:20:A:C2	1:A:21:A:C4	0.49	2.99	3	1
1:A:23:U:H2'	1:A:24:C:C6	0.48	2.42	19	2
1:A:10:C:O2'	1:A:11:C:H5'	0.48	2.08	19	6
1:A:2:G:H2'	1:A:3:C:C6	0.48	2.44	25	1
1:A:15:G:H2'	1:A:16:G:O4'	0.48	2.08	13	1
1:A:3:C:H2'	1:A:4:G:O4'	0.47	2.08	16	9
1:A:7:G:N1	1:A:21:A:N1	0.47	2.63	8	1
1:A:22:G:C5	1:A:23:U:C5	0.46	3.03	16	1
1:A:15:G:N2	1:A:16:G:C8	0.46	2.83	25	1
1:A:7:G:C2	1:A:22:G:C6	0.46	3.04	4	9
1:A:7:G:N2	1:A:20:A:H61	0.45	2.09	3	1
1:A:1:G:H2'	1:A:2:G:O4'	0.45	2.11	6	1
1:A:8:C:N3	1:A:20:A:N6	0.45	2.65	20	1
1:A:7:G:N2	1:A:21:A:N1	0.44	2.64	18	2
1:A:14:C:H4'	1:A:15:G:N3	0.43	2.28	4	3
1:A:9:A:C2	1:A:10:C:C2	0.43	3.06	2	1
1:A:22:G:C6	1:A:23:U:C4	0.43	3.05	6	2
1:A:7:G:O2'	1:A:8:C:H5'	0.43	2.12	21	1
1:A:4:G:C6	1:A:5:U:C2	0.43	3.06	6	1
1:A:20:A:H1'	1:A:21:A:O4'	0.43	2.14	19	1
1:A:21:A:O2'	1:A:22:G:O4'	0.43	2.36	20	2
1:A:20:A:C6	1:A:21:A:C6	0.42	3.06	20	1
1:A:9:A:C2	1:A:19:G:C2	0.42	3.08	3	3
1:A:14:C:H4'	1:A:15:G:H5'	0.42	1.90	25	1

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:4:G:N2	1:A:25:G:C4	0.42	2.87	1	3
1:A:4:G:C2	1:A:25:G:C5	0.42	3.08	25	1
1:A:2:G:C2	1:A:3:C:C2	0.41	3.07	19	1
1:A:7:G:N2	1:A:21:A:C6	0.41	2.88	10	1
1:A:17:G:O2'	1:A:18:U:C5'	0.41	2.67	5	1
1:A:11:C:C5	1:A:12:U:C4	0.41	3.09	4	1
1:A:3:C:H6	1:A:3:C:O5'	0.41	1.98	25	1
1:A:17:G:C2	1:A:18:U:C2	0.41	3.08	20	1
1:A:6:C:N4	1:A:7:G:C6	0.40	2.89	10	1
1:A:1:G:C2	1:A:2:G:C4	0.40	3.09	12	1
1:A:14:C:C4'	1:A:15:G:H5'	0.40	2.47	25	1
1:A:18:U:O2'	1:A:19:G:C5'	0.40	2.70	7	1

6.3 Torsion angles (i)

6.3.1 Protein backbone (i)

There are no protein molecules in this entry.

6.3.2 Protein sidechains (i)

There are no protein molecules in this entry.

6.3.3 RNA (i)

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers	Suiteness
1	A	26/27 (96%)	3±1 (11±3%)	0±0 (0±0%)	0.49±0.03
All	All	650/675 (96%)	70 (11%)	0 (0%)	0.49

The overall RNA backbone suiteness is 0.49.

All unique RNA backbone outliers are listed below:

Mol	Chain	Res	Type	Models (Total)
1	A	21	A	25
1	A	20	A	22
1	A	22	G	19
1	A	2	G	2
1	A	7	G	1
1	A	15	G	1

There are no RNA pucker outliers to report.

6.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates (i)

There are no monosaccharides in this entry.

6.6 Ligand geometry (i)

There are no ligands in this entry.

6.7 Other polymers (i)

There are no such molecules in this entry.

6.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

7 Chemical shift validation (i)

No chemical shift data were provided