Full wwPDB X-ray Structure Validation Report (i)

Sep 6, 2023 – 07:12 PM EDT

|		
PDB ID	:	4FYY
Title	:	E. coli Aspartate Transcarbamoylase Complexed with CTP, UTP, and Mg2+
Authors	:	Cockrell, G.M.; Kantrowitz, E.R.
Deposited on	:	2012-07-05
Resolution	:	1.94 Å(reported)

1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 1.94 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.

Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4310 (1.96-1.92)
Clashscore	141614	1023 (1.94-1.94)
Ramachandran outliers	138981	1007 (1.94-1.94)
Sidechain outliers	138945	1007 (1.94-1.94)
RSRZ outliers	127900	4250 (1.96-1.92)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	310	4% ... 87% ... 12% ·
1	C	310	5% ... 87% ... 11% ·
2	B	153	2% ... 85% ... 10% · ·
2	D	153	2% ... 80% ... 14% · ·

2 Entry composition (i)

There are 7 unique types of molecules in this entry. The entry contains 7834 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Aspartate carbamoyltransferase catalytic chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	310	Total	C	N	O	S	0	0	0
			2415	1527	423	456	9			
1	C	310	Total	C	N	O	S	0	0	0
			2415	1527	423	456	9			

- Molecule 2 is a protein called Aspartate carbamoyltransferase regulatory chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	147	Total	C	N	O	S	0	0	0
			1151	723	203	220	5			
2	D	148	Total	C	N	O	S	0	0	0
			1159	729	205	220	5			

- Molecule 3 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	1	Total	Zn	0	0
			1	1		
3	D	1	Total	Zn	0	0
			1	1		

- Molecule 4 is URIDINE 5'-TRIPHOSPHATE (three-letter code: UTP) (formula: $C_9H_{15}N_2O_{15}P_3$).

UTP

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	B	1	Total	C	N	O	P	0	0
			29	9	2	15	3		
4	D	1	Total	C	N	O	P	0	0
			29	9	2	15	3		

- Molecule 5 is CYTIDINE-5'-TRIPHOSPHATE (three-letter code: CTP) (formula: $C_9H_{16}N_3O_{14}P_3$).

CTP

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	B	1	Total	C	N	O	P	0	0
			29	9	3	14	3		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
5	D	1	29	9	3	14	3	0	0

- Molecule 6 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Mg		
6	B	1	1	1	0	0
6	D	1	1	1	0	0

- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
7	A	193	193	193	0	0
7	B	84	84	84	0	0
7	C	206	206	206	0	0
7	D	91	91	91	0	0

3 Residue-property plots ⓘ

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Aspartate carbamoyltransferase catalytic chain


- Molecule 1: Aspartate carbamoyltransferase catalytic chain

- Molecule 2: Aspartate carbamoyltransferase regulatory chain

- Molecule 2: Aspartate carbamoyltransferase regulatory chain

4　Data and refinement statistics ⓘ

Property	Value	Source
Space group	P 3 2 1	Depositor
Cell constants a, b, c, α, β, γ	121.13Å　　121.13Å　　141.44Å 90.00°　　90.00°　　120.00°	Depositor
Resolution (Å)	49.18　–　1.94 49.18　–　1.94	Depositor EDS
% Data completeness (in resolution range)	100.0 (49.18-1.94) 99.8 (49.18-1.94)	Depositor EDS
R_{merge}	0.14	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	3.22 (at 1.94Å)	Xtriage
Refinement program	PHENIX	Depositor
R, R_{free}	0.193　,　0.236 0.204　,　0.245	Depositor DCC
R_{free} test set	4462 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å^2)	33.5	Xtriage
Anisotropy	0.102	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.29 , 26.0	EDS
L-test for twinning[2]	$< \lvert L \rvert > = 0.42, < L^2 > = 0.25$	Xtriage
Estimated twinning fraction	0.199 for -h,-k,l	Xtriage
F_o,F_c correlation	0.97	EDS
Total number of atoms	7834	wwPDB-VP
Average B, all atoms (Å^2)	41.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.31% of the height of the origin peak. No significant pseudotranslation is detected.*

---

[1]Intensities estimated from amplitudes.

[2]Theoretical values of $< \lvert L \rvert >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5   Model quality (i)

5.1   Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: UTP, ZN, CTP, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#|Z| >5	RMSZ	#|Z| >5
1	A	0.36	0/2461	0.56	0/3339
1	C	0.38	0/2461	0.59	0/3339
2	B	0.32	0/1168	0.52	0/1579
2	D	0.33	0/1176	0.52	0/1590
All	All	0.36	0/7266	0.56	0/9847

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2   Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2415	0	2422	22	0
1	C	2415	0	2422	24	0
2	B	1151	0	1170	15	0
2	D	1159	0	1183	18	0
3	B	1	0	0	0	0
3	D	1	0	0	0	0
4	B	29	0	11	1	0
4	D	29	0	11	2	0
5	B	29	0	12	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	D	29	0	12	2	0
6	B	1	0	0	0	0
6	D	1	0	0	0	0
7	A	193	0	0	6	0
7	B	84	0	0	2	0
7	C	206	0	0	6	0
7	D	91	0	0	3	0
All	All	7834	0	7243	79	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (79) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:7:LYS:NZ	7:C:595:HOH:O	2.18	0.76
2:B:150:VAL:O	2:B:153:ASN:ND2	2.18	0.76
1:C:189:PRO:HB3	1:C:243:VAL:HG21	1.71	0.73
1:A:17:ARG:NH2	7:A:536:HOH:O	2.21	0.72
1:C:29:LYS:HB3	1:C:310:LEU:HD13	1.74	0.70
1:C:305:ASN:ND2	7:C:606:HOH:O	2.25	0.70
1:C:76:SER:OG	1:C:77:ALA:N	2.26	0.69
2:D:82:THR:OG1	2:D:96:ARG:NH2	2.28	0.67
1:A:17:ARG:NH1	7:A:535:HOH:O	2.28	0.67
5:D:203:CTP:O2A	7:D:330:HOH:O	2.12	0.67
1:A:35:GLN:NE2	7:A:478:HOH:O	2.27	0.66
1:C:9:ILE:HG21	1:C:299:LEU:HD21	1.78	0.66
2:D:134:ILE:HG23	2:D:147:HIS:CG	2.32	0.64
1:A:9:ILE:HG21	1:A:299:LEU:HD21	1.80	0.62
2:B:10:GLU:OE1	7:B:330:HOH:O	2.16	0.61
1:A:231:GLN:H	1:A:231:GLN:CD	2.04	0.61
2:D:151:LEU:O	7:D:390:HOH:O	2.16	0.60
1:A:33:ASN:ND2	7:A:451:HOH:O	2.26	0.60
2:D:6:LYS:HG2	2:D:7:LEU:HD13	1.82	0.59
1:C:33:ASN:O	7:C:461:HOH:O	2.17	0.58
1:A:113:ARG:NH2	7:A:489:HOH:O	2.37	0.58
2:B:128:ARG:NH1	2:B:130:ARG:HG2	2.18	0.58
1:C:80:SER:OG	1:C:81:LEU:N	2.37	0.57
2:B:55:ARG:NH1	7:B:346:HOH:O	2.35	0.57
1:C:269:ARG:HB2	7:C:563:HOH:O	2.04	0.57
2:D:45:GLY:HA3	2:D:48:LEU:HD11	1.87	0.56

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:29:LYS:HD3	1:C:310:LEU:HD13	1.88	0.56
1:C:232:LYS:H	1:C:232:LYS:HE2	1.71	0.55
2:D:130:ARG:HD3	2:D:133:ASP:OD1	2.06	0.55
2:D:50:SER:OG	4:D:202:UTP:O2G	2.19	0.55
2:B:99:LEU:HD12	2:B:100:PRO:HD2	1.89	0.55
1:A:49:PHE:O	1:A:75:ASP:HB3	2.06	0.55
2:D:127:VAL:HG13	2:D:134:ILE:HD11	1.89	0.54
2:B:128:ARG:NH2	2:B:144:GLU:OE1	2.27	0.54
2:D:19:ASP:HA	2:D:58:LEU:HD12	1.90	0.53
2:B:128:ARG:HE	2:B:144:GLU:CD	2.11	0.53
1:C:243:VAL:HG22	1:C:247:PHE:HE2	1.73	0.52
2:D:69:ASP:HB2	7:D:389:HOH:O	2.10	0.51
2:B:69:ASP:O	2:B:73:GLN:HG2	2.11	0.51
1:A:84:LYS:HB2	1:A:86:GLU:O	2.11	0.50
2:B:128:ARG:HD2	2:B:130:ARG:NH1	2.26	0.50
2:D:94:LYS:NZ	5:D:203:CTP:O2G	2.43	0.50
2:B:19:ASP:OD2	2:B:20:HIS:N	2.45	0.50
2:B:128:ARG:NE	2:B:144:GLU:OE2	2.27	0.49
1:A:105:ARG:NH2	1:A:167:ARG:HH22	2.12	0.48
1:C:232:LYS:H	1:C:232:LYS:CE	2.25	0.48
1:C:80:SER:C	1:C:82:GLY:HA3	2.34	0.48
2:B:49:PRO:HG2	4:B:202:UTP:H4'	1.94	0.48
1:A:213:SER:O	1:A:246:GLN:HG3	2.13	0.48
1:C:218:VAL:O	1:C:222:VAL:HG13	2.14	0.47
1:A:244:LYS:HA	1:A:247:PHE:CZ	2.50	0.47
1:C:242:ASN:OD1	7:C:548:HOH:O	2.21	0.46
2:D:65:PHE:CD2	2:D:85:ARG:HG2	2.51	0.45
1:A:12:ILE:HD13	1:A:135:PRO:HA	1.99	0.45
1:A:78:ASN:OD1	1:A:78:ASN:N	2.42	0.45
1:A:254:LEU:HD22	1:A:261:MET:HE1	1.99	0.44
1:C:84:LYS:NZ	1:C:86:GLU:H	2.16	0.44
2:B:23:ALA:HB2	2:B:55:ARG:HD3	1.99	0.43
1:C:84:LYS:HZ1	1:C:86:GLU:H	1.66	0.43
1:C:236:ASP:HB3	1:C:239:GLU:HB3	2.01	0.43
2:B:44:ILE:HB	2:D:44:ILE:HB	2.00	0.43
2:D:6:LYS:HE3	2:D:7:LEU:HD13	2.00	0.43
2:D:130:ARG:NH1	2:D:133:ASP:OD2	2.50	0.43
1:A:81:LEU:HD23	1:A:81:LEU:HA	1.83	0.42
1:C:189:PRO:HB3	1:C:243:VAL:CG2	2.47	0.42
2:B:128:ARG:HH12	2:B:130:ARG:HG2	1.83	0.42
1:A:125:LEU:HD23	1:A:299:LEU:HD23	2.01	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:233:GLU:O	1:A:235:LEU:N	2.53	0.42
2:D:99:LEU:HD12	2:D:99:LEU:HA	1.91	0.42
1:C:13:ASN:ND2	7:C:583:HOH:O	2.34	0.41
1:C:35:GLN:OE1	1:C:38:LEU:HD13	2.21	0.41
1:A:279:LYS:HG3	7:A:488:HOH:O	2.21	0.41
1:A:232:LYS:HA	1:A:232:LYS:HD2	1.90	0.41
2:D:52:GLU:HG2	4:D:202:UTP:O3G	2.20	0.41
1:C:12:ILE:HG13	1:C:171:SER:OG	2.20	0.41
1:A:272:GLU:H	1:A:272:GLU:CD	2.24	0.41
1:C:24:LEU:HD13	1:C:143:PHE:HA	2.03	0.41
2:D:130:ARG:NE	2:D:135:ALA:HB2	2.36	0.40
1:A:36:PRO:HA	1:A:65:ARG:O	2.21	0.40

There are no symmetry-related clashes.

5.3　Torsion angles ⓘ

5.3.1　Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	308/310 (99%)	298 (97%)	8 (3%)	2 (1%)	25	13
1	C	308/310 (99%)	293 (95%)	11 (4%)	4 (1%)	12	4
2	B	145/153 (95%)	140 (97%)	5 (3%)	0	100	100
2	D	146/153 (95%)	137 (94%)	9 (6%)	0	100	100
All	All	907/926 (98%)	868 (96%)	33 (4%)	6 (1%)	22	11

All (6) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	270	VAL
1	A	234	ARG
1	C	76	SER

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	C	84	LYS
1	C	78	ASN
1	C	79	THR

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	261/261 (100%)	248 (95%)	13 (5%)	24	9
1	C	261/261 (100%)	250 (96%)	11 (4%)	30	14
2	B	131/137 (96%)	128 (98%)	3 (2%)	50	38
2	D	132/137 (96%)	125 (95%)	7 (5%)	22	9
All	All	785/796 (99%)	751 (96%)	34 (4%)	29	14

All (34) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	12	ILE
1	A	40	LYS
1	A	59	PHE
1	A	134	HIS
1	A	199	LEU
1	A	232	LYS
1	A	233	GLU
1	A	234	ARG
1	A	246	GLN
1	A	269	ARG
1	A	271	ASP
1	A	285	TYR
1	A	309	VAL
2	B	55	ARG
2	B	102	ARG
2	B	153	ASN
1	C	59	PHE
1	C	84	LYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	C	134	HIS
1	C	153	ASP
1	C	232	LYS
1	C	233	GLU
1	C	269	ARG
1	C	279	LYS
1	C	285	TYR
1	C	305	ASN
1	C	306	ARG
2	D	7	LEU
2	D	8	GLN
2	D	14	ARG
2	D	80	GLN
2	D	85	ARG
2	D	134	ILE
2	D	151	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
2	D	8	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no monosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 8 ligands modelled in this entry, 4 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#|Z| > 2$	Counts	RMSZ	$\#|Z| > 2$
5	CTP	D	203	6	26,30,30	0.87	2 (7%)	39,47,47	1.04	2 (5%)
5	CTP	B	203	6	26,30,30	0.87	1 (3%)	39,47,47	1.01	1 (2%)
4	UTP	B	202	6	22,30,30	1.61	4 (18%)	27,47,47	1.16	4 (14%)
4	UTP	D	202	6	22,30,30	1.58	4 (18%)	27,47,47	1.34	4 (14%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	CTP	D	203	6	-	0/22/38/38	0/2/2/2
5	CTP	B	203	6	-	2/22/38/38	0/2/2/2
4	UTP	B	202	6	-	0/20/38/38	0/2/2/2
4	UTP	D	202	6	-	0/20/38/38	0/2/2/2

All (11) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	B	202	UTP	C3'-C2'	-4.97	1.39	1.53
4	D	202	UTP	C3'-C2'	-4.85	1.40	1.53
4	B	202	UTP	C2'-C1'	-2.80	1.49	1.53
4	D	202	UTP	C2'-C1'	-2.58	1.49	1.53
4	B	202	UTP	C2-N3	-2.46	1.33	1.38
4	D	202	UTP	C2-N3	-2.35	1.33	1.38
5	D	203	CTP	C4-N3	2.20	1.39	1.34
4	D	202	UTP	O4'-C4'	-2.19	1.40	1.45
4	B	202	UTP	O4'-C4'	-2.18	1.40	1.45
5	B	203	CTP	C6-C5	2.10	1.39	1.35
5	D	203	CTP	C6-C5	2.08	1.39	1.35

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
4	D	202	UTP	PB-O3B-PG	-3.36	121.30	132.83
5	D	203	CTP	PB-O3B-PG	-3.03	122.43	132.83
4	D	202	UTP	PB-O3A-PA	-2.90	122.87	132.83
4	D	202	UTP	C3'-C2'-C1'	2.84	105.25	100.98
4	B	202	UTP	PB-O3B-PG	-2.74	123.41	132.83
4	B	202	UTP	C3'-C2'-C1'	2.69	105.03	100.98
5	D	203	CTP	PB-O3A-PA	-2.66	123.71	132.83
5	B	203	CTP	PB-O3B-PG	-2.64	123.76	132.83
4	D	202	UTP	O5'-C5'-C4'	2.22	116.63	108.99
4	B	202	UTP	PB-O3A-PA	-2.19	125.32	132.83
4	B	202	UTP	O5'-C5'-C4'	2.15	116.39	108.99

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	B	203	CTP	C5'-O5'-PA-O3A
5	B	203	CTP	PB-O3A-PA-O2A

There are no ring outliers.

3 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	D	203	CTP	2	0
4	B	202	UTP	1	0
4	D	202	UTP	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

Ligand CTP D 203

Bond lengths

Bond angles

Torsions

Rings

Ligand CTP B 203

Bond lengths

Bond angles

Torsions

Rings

Ligand UTP B 202

Bond lengths

Bond angles

Torsions

Rings

Ligand UTP D 202

Bond lengths

Bond angles

Torsions

Rings

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å^2)	Q<0.9
1	A	310/310 (100%)	-0.06	13 (4%)	36	43	20, 32, 73, 213	0
1	C	310/310 (100%)	-0.25	15 (4%)	30	38	18, 28, 73, 163	0
2	B	147/153 (96%)	-0.16	3 (2%)	65	71	26, 42, 77, 181	0
2	D	148/153 (96%)	-0.22	3 (2%)	65	71	24, 42, 77, 135	0
All	All	915/926 (98%)	-0.17	34 (3%)	41	49	18, 35, 77, 213	0

All (34) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	81	LEU	26.7
1	A	82	GLY	20.8
1	A	80	SER	13.8
1	A	83	LYS	10.9
2	D	153	ASN	8.4
1	A	77	ALA	7.3
1	C	81	LEU	6.3
1	C	79	THR	6.2
1	C	77	ALA	6.0
1	A	79	THR	5.8
1	C	83	LYS	5.7
1	A	310	LEU	5.7
2	B	130	ARG	5.7
1	C	80	SER	4.7
1	A	78	ASN	4.5
1	A	76	SER	4.2
1	A	309	VAL	3.9
1	C	309	VAL	3.8
2	D	6	LYS	3.8
1	C	76	SER	3.8
2	B	153	ASN	3.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	131	ALA	3.6
1	A	84	LYS	3.6
1	C	82	GLY	3.6
1	C	310	LEU	3.5
1	C	78	ASN	3.3
1	A	247	PHE	3.2
1	C	84	LYS	3.0
1	C	246	GLN	2.9
2	D	132	ASN	2.5
1	C	85	GLY	2.4
1	C	240	TYR	2.3
1	A	1	ALA	2.2
1	C	247	PHE	2.2

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no monosaccharides in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

| Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
6	MG	B	204	1/1	0.84	0.03	45,45,45,45	0
6	MG	D	204	1/1	0.88	0.05	44,44,44,44	0
4	UTP	B	202	29/29	0.95	0.07	29,41,53,94	0
4	UTP	D	202	29/29	0.95	0.08	32,46,57,72	0
5	CTP	B	203	29/29	0.96	0.09	30,45,60,77	0
5	CTP	D	203	29/29	0.96	0.08	33,41,60,64	0
3	ZN	B	201	1/1	0.99	0.08	25,25,25,25	0
3	ZN	D	201	1/1	0.99	0.08	27,27,27,27	0

The following is a graphical depiction of the model fit to experimental electron density of all

instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.


Electron density around UTP B 202:

$2mF_o\text{-}DF_c$ (at 0.7 rmsd) in gray
$mF_o\text{-}DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)

Electron density around UTP D 202:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

Electron density around CTP B 203:

$2mF_o\text{-}DF_c$ (at 0.7 rmsd) in gray
$mF_o\text{-}DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)

Electron density around CTP D 203:

$2mF_o\text{-}DF_c$ (at 0.7 rmsd) in gray
$mF_o\text{-}DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)

6.5 Other polymers ⓘ

There are no such residues in this entry.