Full wwPDB X-ray Structure Validation Report (i)

Oct 23, 2021 – 11:02 AM EDT

PDB ID	:	1FZR
Title	:	CRYSTAL STRUCTURE OF BACTERIOPHAGE T7 ENDONUCLEASE I
Authors	:	Hadden, J.M.; Convery, M.A.; Declais, A.C.; Lilley, D.M.J.; Phillips, S.E.V.
Deposited on	:	2000-10-04
Resolution	:	2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.23.2
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.23.2

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5197 (2.10-2.10)
Clashscore	141614	5710 (2.10-2.10)
Ramachandran outliers	138981	5647 (2.10-2.10)
Sidechain outliers	138945	5648 (2.10-2.10)
RSRZ outliers	127900	5083 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	138	5% 68% 25% • 7%
1	B	138	4% 78% 15% 7%
1	C	138	5% 63% 30% • 7%
1	D	138	3% 71% 20% • 7%

2 Entry composition (i)

There are 2 unique types of molecules in this entry. The entry contains 4702 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ENDONUCLEASE I.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	129	1049	681	178	189	1	0	0	0
1	B	129	1060	688	178	193	1	0	0	0
1	C	129	1048	680	176	191	1	0	0	0
1	D	129	1060	689	179	191	1	0	0	0

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	65	LYS	GLU	engineered mutation	UNP P00641
B	65	LYS	GLU	engineered mutation	UNP P00641
C	65	LYS	GLU	engineered mutation	UNP P00641
D	65	LYS	GLU	engineered mutation	UNP P00641

- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
2	A	111	111	111	0	0
2	B	151	151	151	0	0
2	C	77	77	77	0	0
2	D	146	146	146	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: ENDONUCLEASE I

Chain A:





• Molecule 1: ENDONUCLEASE I

Chain B:



• Molecule 1: ENDONUCLEASE I

Chain C:

• Molecule 1: ENDONUCLEASE I

Chain D:



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α, β, γ	123.66Å 135.19Å 61.58Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	28.81 – 2.10 29.88 – 2.10	Depositor EDS
% Data completeness (in resolution range)	99.1 (28.81-2.10) 99.1 (29.88-2.10)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	2.45 (at 2.10Å)	Xtriage
Refinement program	CNS 0.9	Depositor
R, R_{free}	0.198 , 0.235 0.200 , 0.235	Depositor DCC
R_{free} test set	6061 reflections (10.02%)	wwPDB-VP
Wilson B-factor (Å^2)	46.8	Xtriage
Anisotropy	0.022	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.34 , 58.3	EDS
L-test for twinning[2]	$< \|L\| > = 0.49$, $< L^2 > = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o,F_c correlation	0.96	EDS
Total number of atoms	4702	wwPDB-VP
Average B, all atoms (Å^2)	54.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.44% of the height of the origin peak. No significant pseudotranslation is detected.*

[1] Intensities estimated from amplitudes.

[2] Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\|>5$	RMSZ	$\#\|Z\|>5$
1	A	0.50	0/1079	0.71	1/1456 (0.1%)
1	B	0.58	0/1090	0.76	2/1470 (0.1%)
1	C	0.45	0/1078	0.66	0/1457
1	D	0.55	0/1090	0.75	2/1469 (0.1%)
All	All	0.52	0/4337	0.72	5/5852 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand.A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	B	0	1
1	C	0	1
All	All	0	3

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
1	D	75	ARG	NE-CZ-NH2	-7.12	116.74	120.30
1	D	68	GLY	N-CA-C	-6.30	97.36	113.10
1	B	68	GLY	N-CA-C	-6.05	97.97	113.10
1	A	68	GLY	N-CA-C	-5.30	99.84	113.10
1	B	144	ARG	NE-CZ-NH2	-5.21	117.69	120.30

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	52	TYR	Sidechain

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Group
1	B	52	TYR	Sidechain
1	C	52	TYR	Sidechain

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1049	0	1037	21	0
1	B	1060	0	1054	14	0
1	C	1048	0	1028	28	0
1	D	1060	0	1061	25	0
2	A	111	0	0	0	0
2	B	151	0	0	4	0
2	C	77	0	0	3	0
2	D	146	0	0	5	0
All	All	4702	0	4180	79	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (79) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:141:ARG:HB3	1:D:141:ARG:HH11	1.30	0.92
1:D:141:ARG:HH11	1:D:141:ARG:CB	1.94	0.80
1:A:108:SER:OG	1:A:111:GLU:HG3	1.83	0.78
1:B:48:SER:OG	1:B:50:HIS:HE1	1.68	0.76
1:A:82:ARG:HH22	1:A:118:ILE:HA	1.50	0.76
1:A:73:ASP:O	1:A:77:LYS:HG2	1.92	0.69
1:B:89:ASP:HB2	1:B:135:LYS:HE3	1.73	0.69
1:C:48:SER:OG	1:C:50:HIS:HE1	1.76	0.69
1:C:72:SER:O	1:C:76:LYS:HG3	1.94	0.68
1:A:82:ARG:NH2	1:A:118:ILE:HA	2.11	0.65
1:D:141:ARG:HH11	1:D:141:ARG:CG	2.10	0.64
1:C:137:VAL:HB	1:C:139:PHE:CE1	2.33	0.63
1:C:33:LYS:O	1:D:59:PRO:HG3	1.99	0.62

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:72:SER:HA	1:C:75:ARG:NH1	2.14	0.62
1:D:131:LYS:HE3	2:D:193:HOH:O	1.99	0.62
1:D:141:ARG:HB3	1:D:141:ARG:NH1	2.11	0.60
1:C:91:ARG:HG2	1:C:119:LYS:HB2	1.84	0.60
1:C:79:LEU:O	1:C:82:ARG:HB3	2.03	0.59
1:D:18:GLY:O	1:D:22:LYS:HG3	2.02	0.59
1:A:55:ASP:OD2	1:A:65:LYS:HD2	2.03	0.58
1:A:17:SER:OG	1:A:20:GLU:HB2	2.06	0.55
1:C:60:ASN:HA	1:C:134:LYS:HG3	1.87	0.55
1:C:87:GLU:CD	1:C:87:GLU:H	2.08	0.55
1:C:135:LYS:HE3	2:C:185:HOH:O	2.06	0.55
1:C:77:LYS:O	1:C:81:ILE:HG13	2.07	0.54
1:C:60:ASN:HB2	1:C:134:LYS:HA	1.90	0.53
1:A:48:SER:OG	1:A:50:HIS:HE1	1.90	0.53
1:D:108:SER:OG	1:D:111:GLU:HG3	2.08	0.53
1:D:116:HIS:HE1	2:D:192:HOH:O	1.92	0.53
1:A:107:THR:HA	1:A:111:GLU:OE2	2.10	0.52
1:A:45:ILE:HD12	1:B:50:HIS:CE1	2.45	0.52
1:A:130:ILE:CD1	2:B:189:HOH:O	2.57	0.51
1:D:19:LEU:O	1:D:23:VAL:HG23	2.10	0.51
1:C:62:ILE:HD13	1:C:91:ARG:CZ	2.41	0.51
1:D:24:SER:O	1:D:28:GLU:HG3	2.11	0.51
1:C:85:HIS:N	1:C:86:PRO:HD3	2.26	0.51
1:D:17:SER:OG	1:D:20:GLU:HB2	2.11	0.50
1:A:82:ARG:NH2	1:A:117:GLY:O	2.45	0.50
1:C:50:HIS:CE1	1:D:45:ILE:HD12	2.47	0.50
1:D:75:ARG:NH2	1:D:101:LEU:O	2.45	0.49
1:D:35:GLU:HG2	1:D:38:GLU:CB	2.42	0.49
1:C:135:LYS:HD3	2:C:187:HOH:O	2.12	0.48
1:D:141:ARG:CG	1:D:141:ARG:NH1	2.73	0.48
1:C:123:LYS:C	1:C:124:LEU:HD22	2.34	0.47
1:A:137:VAL:HB	1:A:139:PHE:CE1	2.50	0.47
1:C:123:LYS:HB3	1:C:124:LEU:HD22	1.97	0.47
1:C:85:HIS:HA	1:C:87:GLU:OE2	2.15	0.46
1:B:100:LYS:HE3	1:B:106:PRO:O	2.16	0.46
1:C:40:LYS:NZ	2:C:163:HOH:O	2.49	0.46
1:B:48:SER:OG	1:B:50:HIS:CE1	2.58	0.46
1:A:110:GLY:O	1:A:114:GLU:HG3	2.17	0.45
1:A:130:ILE:HD11	2:B:189:HOH:O	2.16	0.45
1:A:123:LYS:O	1:A:124:LEU:HD13	2.17	0.45
1:D:48:SER:OG	1:D:50:HIS:HE1	2.00	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:44:VAL:HG22	1:D:49:ASN:ND2	2.32	0.44
1:D:33:LYS:HA	1:D:33:LYS:HD2	1.68	0.44
1:C:102:TYR:CE1	1:C:105:SER:HB2	2.52	0.44
1:A:128:GLU:H	1:A:128:GLU:CD	2.21	0.44
1:A:124:LEU:HG	1:B:26:GLN:HG3	2.00	0.44
1:B:103:LYS:HG3	1:C:121:ALA:HA	2.00	0.44
1:A:57:LEU:HB3	1:B:35:GLU:HB2	2.00	0.43
1:D:98:ARG:NH1	2:D:245:HOH:O	2.51	0.43
1:A:91:ARG:HG2	1:A:119:LYS:HB2	2.01	0.43
1:B:73:ASP:HB3	2:B:191:HOH:O	2.18	0.42
1:B:89:ASP:CB	1:B:135:LYS:HE3	2.47	0.42
1:C:128:GLU:H	1:C:128:GLU:CD	2.22	0.42
1:B:105:SER:HA	1:B:106:PRO:HD3	1.77	0.42
1:C:44:VAL:HG22	1:D:49:ASN:HD22	1.83	0.42
1:A:102:TYR:CE1	1:A:105:SER:HB2	2.55	0.42
1:D:128:GLU:HG3	2:D:246:HOH:O	2.20	0.42
1:D:98:ARG:HD3	1:D:98:ARG:HA	1.67	0.42
1:C:18:GLY:O	1:C:22:LYS:HG3	2.20	0.41
1:C:84:GLN:C	1:C:86:PRO:HD3	2.42	0.41
1:D:83:GLU:HG3	2:D:239:HOH:O	2.20	0.41
1:A:59:PRO:HG3	1:B:33:LYS:O	2.20	0.40
1:B:87:GLU:HG3	1:B:138:PRO:HG3	2.03	0.40
1:B:71:GLU:HB2	2:B:191:HOH:O	2.21	0.40
1:C:55:ASP:OD2	1:C:65:LYS:HD2	2.22	0.40
1:D:45:ILE:HA	1:D:46:PRO:HD3	1.87	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	127/138 (92%)	123 (97%)	3 (2%)	1 (1%)	19 15

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	127/138 (92%)	123 (97%)	4 (3%)	0	100	100
1	C	127/138 (92%)	123 (97%)	3 (2%)	1 (1%)	19	15
1	D	127/138 (92%)	124 (98%)	3 (2%)	0	100	100
All	All	508/552 (92%)	493 (97%)	13 (3%)	2 (0%)	34	32

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	103	LYS
1	C	103	LYS

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	113/124 (91%)	109 (96%)	4 (4%)	36	38
1	B	116/124 (94%)	113 (97%)	3 (3%)	46	50
1	C	113/124 (91%)	106 (94%)	7 (6%)	18	15
1	D	116/124 (94%)	111 (96%)	5 (4%)	29	29
All	All	458/496 (92%)	439 (96%)	19 (4%)	30	31

All (19) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	27	LEU
1	A	58	LEU
1	A	124	LEU
1	A	142	LEU
1	B	98	ARG
1	B	124	LEU
1	B	142	LEU
1	C	58	LEU
1	C	73	ASP

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	C	74	ASP
1	C	87	GLU
1	C	98	ARG
1	C	107	THR
1	C	142	LEU
1	D	35	GLU
1	D	58	LEU
1	D	124	LEU
1	D	141	ARG
1	D	142	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	50	HIS
1	B	49	ASN
1	B	50	HIS
1	C	50	HIS
1	D	49	ASN
1	D	50	HIS

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

There are no ligands in this entry.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å^2)	Q<0.9
1	A	129/138 (93%)	0.02	7 (5%)	25	31	33, 49, 75, 95	0
1	B	129/138 (93%)	-0.00	5 (3%)	39	45	33, 42, 63, 73	0
1	C	129/138 (93%)	0.16	7 (5%)	25	31	37, 59, 88, 108	0
1	D	129/138 (93%)	0.16	4 (3%)	49	55	32, 45, 92, 102	0
All	All	516/552 (93%)	0.09	23 (4%)	33	38	32, 50, 86, 108	0

All (23) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	102	TYR	6.5
1	C	102	TYR	4.2
1	A	103	LYS	3.8
1	D	39	TRP	3.7
1	C	105	SER	3.4
1	C	140	ASP	3.4
1	C	106	PRO	3.3
1	D	32	ILE	3.1
1	A	140	ASP	3.0
1	B	88	LEU	3.0
1	D	142	LEU	3.0
1	B	81	ILE	2.9
1	A	104	GLY	2.8
1	C	137	VAL	2.7
1	B	90	ILE	2.7
1	C	107	THR	2.6
1	C	104	GLY	2.6
1	B	17	SER	2.3
1	A	41	VAL	2.2
1	D	29	SER	2.2
1	A	43	TYR	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	105	SER	2.2
1	B	85	HIS	2.1

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

There are no ligands in this entry.

6.5 Other polymers (i)

There are no such residues in this entry.