



Full wwPDB X-ray Structure Validation Report i

May 16, 2020 – 05:46 pm BST

PDB ID : 4FZF
Title : Crystal structure of MST4-MO25 complex with DKI
Authors : Shi, Z.B.; Zhou, Z.C.
Deposited on : 2012-07-06
Resolution : 3.64 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>
with specific help available everywhere you see the i symbol.

The following versions of software and data (see [references](#) ①) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.11
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.64 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1341 (3.78-3.50)
Clashscore	141614	1439 (3.78-3.50)
Ramachandran outliers	138981	1391 (3.78-3.50)
Sidechain outliers	138945	1391 (3.78-3.50)
RSRZ outliers	127900	1242 (3.78-3.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain				
1	A	328	4%	67%	27%	...	
2	B	283		66%	26%	6%	•

2 Entry composition (i)

There are 3 unique types of molecules in this entry. The entry contains 4887 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Calcium-binding protein 39.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	325	Total	C 2650	N 1695	O 442	S 502	11	0	0

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	7	GLY	-	EXPRESSION TAG	UNP Q9Y376
A	8	ALA	-	EXPRESSION TAG	UNP Q9Y376
A	9	MET	-	EXPRESSION TAG	UNP Q9Y376
A	10	ALA	-	EXPRESSION TAG	UNP Q9Y376


- Molecule 2 is a protein called Serine/threonine-protein kinase MST4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	279	Total	C 2209	N 1423	O 359	S 419	8	0	0

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	15	GLY	-	EXPRESSION TAG	UNP Q9P289
B	16	ALA	-	EXPRESSION TAG	UNP Q9P289
B	17	MET	-	EXPRESSION TAG	UNP Q9P289
B	162	ALA	ASP	ENGINEERED MUTATION	UNP Q9P289

- Molecule 3 is 5-AMINO-3-{|4-(AMINOSULFONYL)PHENYL]AMINO}-N-(2,6-DIFLUOROPHENYL)-1H-1,2,4-TRIAZOLE-1-CARBOTHIOAMIDE (three-letter code: DKI) (formula: C₁₅H₁₃F₂N₇O₂S₂).




Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
3	B	1	Total	C	F	N	O	S	0	0
			28	15	2	7	2	2		


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Calcium-binding protein 39



- Molecule 2: Serine/threonine-protein kinase MST4



4 Data and refinement statistics (i)

Property	Value	Source
Space group	I 4 3 2	Depositor
Cell constants a, b, c, α , β , γ	239.46 Å 239.46 Å 239.46 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	38.85 – 3.64 38.85 – 3.64	Depositor EDS
% Data completeness (in resolution range)	99.1 (38.85-3.64) 99.2 (38.85-3.64)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ ¹	4.50 (at 3.66 Å)	Xtriage
Refinement program	REFMAC 5.6.0117	Depositor
R , R_{free}	0.252 , 0.309 0.229 , 0.291	Depositor DCC
R_{free} test set	657 reflections (4.94%)	wwPDB-VP
Wilson B-factor (Å ²)	105.4	Xtriage
Anisotropy	0.000	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.28 , 63.7	EDS
L-test for twinning ²	$< L > = 0.51$, $< L^2 > = 0.35$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	4887	wwPDB-VP
Average B, all atoms (Å ²)	116.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 1.74% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [\(i\)](#)

5.1 Standard geometry [\(i\)](#)

Bond lengths and bond angles in the following residue types are not validated in this section: DKI

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z > 5	RMSZ	# Z > 5
1	A	0.39	0 / 2696	0.59	2 / 3634 (0.1%)
2	B	0.50	1 / 2253 (0.0%)	0.64	0 / 3039
All	All	0.44	1 / 4949 (0.0%)	0.61	2 / 6673 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	203	TRP	CD2-CE2	5.41	1.47	1.41

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	333	LEU	CA-CB-CG	6.46	130.16	115.30
1	A	238	LEU	CA-CB-CG	5.72	128.46	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2650	0	2644	54	0
2	B	2209	0	2227	57	0
3	B	28	0	13	6	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	4887	0	4884	113	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

All (113) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:60:ALA:HB3	2:B:61:GLU:HA	1.28	1.09
2:B:272:LYS:H	2:B:272:LYS:HE2	1.22	1.01
2:B:60:ALA:CB	2:B:61:GLU:HA	1.91	0.99
2:B:64:ILE:H	2:B:64:ILE:HD12	1.33	0.92
1:A:11:SER:HB2	1:A:12:PRO:HD2	1.53	0.91
2:B:102:LEU:HD21	2:B:159:LYS:HG3	1.55	0.89
1:A:88:GLN:HB2	1:A:138:GLU:HG3	1.61	0.82
2:B:237:PRO:HG2	2:B:255:LEU:HD13	1.63	0.80
2:B:173:GLN:HG2	2:B:179:PHE:HA	1.63	0.79
1:A:314:THR:N	1:A:315:GLU:HB2	1.97	0.79
3:B:301:DKI:HAK	3:B:301:DKI:NAP	1.96	0.77
2:B:223:HIS:CD2	2:B:224:PRO:HD2	2.21	0.76
2:B:295:LYS:HE2	2:B:296:ALA:O	1.89	0.72
2:B:272:LYS:H	2:B:272:LYS:CE	1.98	0.72
1:A:87:LEU:HB2	1:A:139:ILE:HG21	1.73	0.70
2:B:272:LYS:HE2	2:B:272:LYS:N	2.01	0.69
1:A:301:LYS:HA	1:A:304:GLU:HG3	1.75	0.69
1:A:90:ILE:HG23	1:A:91:ASP:H	1.57	0.67
1:A:87:LEU:HD12	1:A:90:ILE:HG21	1.76	0.67
2:B:60:ALA:CB	2:B:61:GLU:CA	2.72	0.66
1:A:134:TYR:HE2	1:A:184:ALA:HA	1.60	0.65
1:A:190:ASP:O	1:A:194:ARG:HB2	1.96	0.65
2:B:64:ILE:N	2:B:64:ILE:HD12	2.12	0.61
1:A:11:SER:HB2	1:A:12:PRO:CD	2.28	0.61
2:B:173:GLN:CG	2:B:179:PHE:HA	2.31	0.61
3:B:301:DKI:NAP	3:B:301:DKI:CAK	2.62	0.60
1:A:314:THR:HG22	1:A:320:ASN:HD21	1.66	0.59
1:A:160:ILE:HD12	1:A:160:ILE:H	1.68	0.58
2:B:230:LEU:HB3	2:B:234:ASN:HD22	1.67	0.58
2:B:102:LEU:HD21	2:B:159:LYS:CG	2.31	0.57
2:B:230:LEU:HD22	2:B:234:ASN:HD21	1.68	0.57
2:B:109:ASP:O	2:B:112:ARG:HG3	2.05	0.57
2:B:102:LEU:HD22	2:B:151:LEU:HB2	1.87	0.56

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:230:LEU:HB3	2:B:234:ASN:ND2	2.19	0.56
1:A:152:ARG:HA	1:A:194:ARG:HD3	1.88	0.56
2:B:82:VAL:HG12	2:B:162:ALA:HB2	1.87	0.56
2:B:102:LEU:HD11	2:B:159:LYS:HD2	1.87	0.55
1:A:240:ARG:HH21	1:A:243:PHE:HD1	1.52	0.55
2:B:30:ILE:HD11	3:B:301:DKI:NAP	2.22	0.54
2:B:117:ASP:HB2	2:B:120:GLN:H	1.72	0.54
1:A:87:LEU:CB	1:A:139:ILE:HG21	2.37	0.54
2:B:86:TYR:HB2	2:B:98:ILE:HG22	1.88	0.54
1:A:316:ASP:HB3	1:A:319:PHE:HB3	1.90	0.54
2:B:157:ASP:N	2:B:157:ASP:OD1	2.40	0.54
2:B:31:GLY:O	3:B:301:DKI:HAI	2.07	0.54
2:B:198:SER:O	2:B:201:ASP:HB2	2.08	0.53
2:B:183:PRO:HB3	2:B:228:LEU:HD12	1.90	0.53
2:B:57:LEU:HB2	2:B:93:SER:O	2.09	0.52
2:B:30:ILE:HG12	2:B:31:GLY:N	2.24	0.52
1:A:221:GLU:HA	1:A:222:ASN:O	2.08	0.52
2:B:184:PHE:HE1	2:B:227:VAL:HG21	1.74	0.52
2:B:214:LYS:HE2	2:B:240:VAL:O	2.10	0.52
1:A:68:ALA:HB3	1:A:69:GLN:HE21	1.74	0.52
1:A:262:LEU:O	1:A:265:ASP:HB2	2.11	0.51
1:A:106:LEU:HA	1:A:116:THR:HG21	1.93	0.50
1:A:188:PHE:O	1:A:192:LEU:HG	2.12	0.49
2:B:45:ARG:HG2	2:B:46:THR:HG23	1.95	0.49
1:A:19:LEU:HA	1:A:23:MET:HB2	1.95	0.49
1:A:246:MET:HG3	1:A:247:THR:N	2.28	0.49
1:A:255:ASN:HB2	1:A:281:PHE:HZ	1.78	0.48
1:A:276:HIS:HA	1:A:279:LYS:HE2	1.94	0.48
1:A:153:HIS:HB2	1:A:156:LEU:HB2	1.96	0.47
1:A:91:ASP:HB3	1:A:94:GLY:HA3	1.96	0.47
1:A:108:ARG:O	1:A:115:PRO:HD2	2.14	0.47
1:A:29:GLN:HE21	1:A:30:ASP:H	1.64	0.46
2:B:155:GLN:H	2:B:155:GLN:CD	2.17	0.46
2:B:155:GLN:N	2:B:155:GLN:CD	2.69	0.46
1:A:203:LEU:HB2	1:A:245:ILE:HG12	1.97	0.46
1:A:172:TYR:O	1:A:175:MET:HB2	2.16	0.46
1:A:320:ASN:HA	1:A:323:LYS:HB2	1.98	0.46
1:A:294:ILE:O	1:A:298:ASN:ND2	2.49	0.45
1:A:135:GLU:HG3	1:A:172:TYR:HE1	1.82	0.45
2:B:146:LYS:HB3	2:B:185:TRP:CE2	2.52	0.45
1:A:260:MET:HA	1:A:263:LEU:HD12	1.97	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:267:SER:O	1:A:271:GLN:HG3	2.16	0.45
1:A:314:THR:HG22	1:A:320:ASN:ND2	2.30	0.45
1:A:124:GLN:C	1:A:126:ILE:H	2.20	0.45
2:B:223:HIS:HA	2:B:224:PRO:HD3	1.90	0.45
2:B:184:PHE:HB2	2:B:185:TRP:CE3	2.52	0.44
1:A:239:ASP:O	1:A:240:ARG:HB2	2.17	0.44
1:A:92:PHE:CE2	2:B:78:ASP:HB2	2.52	0.44
3:B:301:DKI:SAE	3:B:301:DKI:FAG	2.65	0.43
2:B:236:PRO:HG3	2:B:257:LYS:HG3	2.01	0.43
2:B:280:LYS:HD2	2:B:282:SER:H	1.83	0.43
1:A:90:ILE:HG23	1:A:91:ASP:N	2.30	0.43
2:B:231:ILE:HB	2:B:232:PRO:HD3	1.99	0.43
1:A:333:LEU:HA	1:A:334:LYS:HA	1.58	0.43
1:A:26:LEU:HD11	1:A:43:VAL:HG11	2.01	0.42
1:A:207:TYR:HE1	1:A:252:LYS:HZ2	1.68	0.42
2:B:172:THR:O	2:B:178:THR:HB	2.19	0.42
2:B:211:GLU:HG3	2:B:216:GLU:HA	2.00	0.42
1:A:154:GLU:HA	1:A:195:HIS:CE1	2.54	0.42
2:B:132:LEU:HD11	2:B:145:ILE:HD13	2.01	0.42
1:A:126:ILE:H	1:A:126:ILE:HG13	1.64	0.42
2:B:211:GLU:HG3	2:B:217:PRO:HD3	2.02	0.42
2:B:64:ILE:CD1	2:B:64:ILE:N	2.76	0.42
1:A:151:ILE:HG22	1:A:160:ILE:HD13	2.02	0.42
1:A:152:ARG:HG2	1:A:194:ARG:HH11	1.85	0.42
2:B:124:MET:HG2	2:B:158:VAL:HG21	2.02	0.42
2:B:276:LYS:HG3	2:B:276:LYS:O	2.19	0.41
2:B:118:GLU:OE2	2:B:243:PHE:HA	2.20	0.41
2:B:211:GLU:O	2:B:215:GLY:N	2.43	0.41
1:A:77:SER:HB3	1:A:79:LEU:HD13	2.02	0.41
2:B:33:GLY:HA3	3:B:301:DKI:HAH	2.02	0.41
1:A:82:THR:HA	1:A:85:ALA:HB3	2.02	0.41
2:B:182:THR:HA	2:B:183:PRO:HD3	1.76	0.41
2:B:295:LYS:HG3	2:B:295:LYS:O	2.21	0.41
1:A:114:THR:OG1	1:A:117:VAL:HG23	2.20	0.41
1:A:104:ASN:HA	1:A:107:ARG:HD3	2.03	0.41
1:A:57:THR:O	1:A:57:THR:HG23	2.21	0.41
1:A:323:LYS:O	1:A:327:VAL:HG23	2.21	0.40
2:B:237:PRO:HG2	2:B:255:LEU:CD1	2.42	0.40
2:B:79:SER:HB2	2:B:134:TYR:CD2	2.56	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	323/328 (98%)	268 (83%)	45 (14%)	10 (3%)	4 32
2	B	275/283 (97%)	240 (87%)	27 (10%)	8 (3%)	4 33
All	All	598/611 (98%)	508 (85%)	72 (12%)	18 (3%)	4 32

All (18) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	63	GLU
2	B	171	ASP
2	B	174	ILE
1	A	11	SER
1	A	28	LYS
1	A	299	GLN
2	B	30	ILE
1	A	222	ASN
2	B	60	ALA
1	A	30	ASP
1	A	60	LYS
1	A	81	SER
1	A	177	THR
1	A	194	ARG
2	B	32	LYS
2	B	195	ALA
2	B	196	TYR
1	A	90	ILE

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	291/299 (97%)	263 (90%)	28 (10%)	8 36
2	B	240/244 (98%)	208 (87%)	32 (13%)	4 23
All	All	531/543 (98%)	471 (89%)	60 (11%)	6 29

All (60) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	14	ASP
1	A	23	MET
1	A	25	VAL
1	A	37	GLU
1	A	42	GLU
1	A	50	MET
1	A	82	THR
1	A	89	LEU
1	A	90	ILE
1	A	107	ARG
1	A	123	GLN
1	A	124	GLN
1	A	149	GLU
1	A	161	LEU
1	A	165	GLN
1	A	175	MET
1	A	187	THR
1	A	194	ARG
1	A	238	LEU
1	A	240	ARG
1	A	247	THR
1	A	258	LEU
1	A	261	ASN
1	A	311	ASN
1	A	312	ASP
1	A	329	GLN
1	A	333	LEU
1	A	334	LYS
2	B	17	MET
2	B	21	GLU
2	B	29	ARG
2	B	37	GLU
2	B	45	ARG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	B	50	VAL
2	B	59	GLU
2	B	61	GLU
2	B	62	ASP
2	B	63	GLU
2	B	64	ILE
2	B	72	THR
2	B	94	LYS
2	B	99	MET
2	B	109	ASP
2	B	119	PHE
2	B	128	ILE
2	B	146	LYS
2	B	155	GLN
2	B	157	ASP
2	B	159	LYS
2	B	169	LEU
2	B	177	ASN
2	B	196	TYR
2	B	228	LEU
2	B	269	LEU
2	B	272	LYS
2	B	276	LYS
2	B	280	LYS
2	B	285	THR
2	B	286	GLU
2	B	295	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (15) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	29	GLN
1	A	63	GLN
1	A	69	GLN
1	A	142	ASN
1	A	165	GLN
1	A	219	HIS
1	A	255	ASN
1	A	271	GLN
1	A	298	ASN
1	A	310	GLN
1	A	320	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	B	68	GLN
2	B	223	HIS
2	B	234	ASN
2	B	277	ASN

5.3.3 RNA [\(i\)](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [\(i\)](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [\(i\)](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	DKI	B	301	-	28,30,30	3.73	8 (28%)	35,44,44	2.11	8 (22%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	DKI	B	301	-	-	5/12/18/18	0/3/3/3

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	301	DKI	CAZ-CAU	10.94	1.50	1.38
3	B	301	DKI	CAY-SBB	-9.92	1.61	1.77
3	B	301	DKI	CAZ-CAV	9.80	1.49	1.38
3	B	301	DKI	CAS-SAE	-6.06	1.58	1.67
3	B	301	DKI	CAS-NAQ	-3.50	1.36	1.41
3	B	301	DKI	CAZ-NAQ	-3.15	1.37	1.43
3	B	301	DKI	CAW-NAR	-2.32	1.35	1.40
3	B	301	DKI	CAX-NAP	2.01	1.36	1.33

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	301	DKI	FAF-CAU-CAZ	6.47	123.28	117.70
3	B	301	DKI	OAD-SBB-OAC	-5.25	110.13	118.76
3	B	301	DKI	FAG-CAV-CAZ	3.86	121.03	117.70
3	B	301	DKI	OAD-SBB-CAY	3.80	111.59	107.35
3	B	301	DKI	CAI-CAU-CAZ	-3.47	118.97	122.88
3	B	301	DKI	CAZ-NAQ-CAS	3.15	127.62	122.99
3	B	301	DKI	CAW-NAR-CAX	-2.21	122.82	129.23
3	B	301	DKI	CAJ-CAV-CAZ	-2.13	120.48	122.88

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	B	301	DKI	SAE-CAS-NAQ-CAZ
3	B	301	DKI	NBA-CAS-NAQ-CAZ
3	B	301	DKI	CAM-CAY-SBB-NAB
3	B	301	DKI	CAN-CAY-SBB-NAB
3	B	301	DKI	CAM-CAY-SBB-OAC


There are no ring outliers.

1 monomer is involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	301	DKI	6	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will

also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	325/328 (99%)	-0.09	13 (4%) 38 25	93, 145, 211, 233	0
2	B	279/283 (98%)	-0.34	0 100 100	39, 73, 132, 172	0
All	All	604/611 (98%)	-0.21	13 (2%) 62 46	39, 110, 205, 233	0

All (13) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	62	PRO	5.0
1	A	110	ILE	3.3
1	A	109	GLN	3.0
1	A	64	THR	2.9
1	A	333	LEU	2.8
1	A	330	ILE	2.6
1	A	334	LYS	2.3
1	A	154	GLU	2.2
1	A	77	SER	2.2
1	A	82	THR	2.1
1	A	305	PHE	2.1
1	A	112	THR	2.0
1	A	155	PRO	2.0

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)


There are no carbohydrates in this entry.

6.4 Ligands [\(i\)](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	DKI	B	301	28/28	0.94	0.22	67,77,83,86	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.