



# Full wwPDB X-ray Structure Validation Report ⓘ

May 14, 2020 – 10:18 pm BST

PDB ID : 5G0O  
Title : Structure of rat neuronal nitric oxide synthase D597N mutant heme domain in complex with 4-METHYL-6-(2-(5-(4-METHYLPIPERAZIN-1-YL) PYRIDIN-3-YL)ETHYL)PYRIDIN-2-AMINE  
Authors : Li, H.; Poulos, T.L.  
Deposited on : 2016-03-21  
Resolution : 1.85 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.85 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	2469 (1.86-1.86)
Clashscore	141614	2625 (1.86-1.86)
Ramachandran outliers	138981	2592 (1.86-1.86)
Sidechain outliers	138945	2592 (1.86-1.86)
RSRZ outliers	127900	2436 (1.86-1.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	422	
1	B	422	

## 2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 7322 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called NITRIC OXIDE SYNTHASE, BRAIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	408	Total 3327	C 2130	N 568	O 607	S 22	0	4	1
1	B	411	Total 3360	C 2150	N 575	O 613	S 22	0	4	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	597	ASN	ASP	engineered mutation	UNP P29476
B	597	ASN	ASP	engineered mutation	UNP P29476

- Molecule 2 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula:  $C_{34}H_{32}FeN_4O_4$ ).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
			Total	C	Fe	N			O
2	A	1	Total 43	C 34	Fe 1	N 4	O 4	0	0

*Continued on next page...*

Continued from previous page...


Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
			Total	C	Fe	N			O
2	B	1	43	34	1	4	4	0	0

- Molecule 3 is 5,6,7,8-TETRAHYDROBIOPTERIN (three-letter code: H4B) (formula:  $C_9H_{15}N_5O_3$ ).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
3	A	1	17	9	5	3	0	0
3	B	1	17	9	5	3	0	0

- Molecule 4 is 4-methyl-6-(2-(5-(4-methylpiperazin-1-yl)pyridin-3-yl)ethyl)pyridin-2-amine (three-letter code: W64) (formula:  $C_{18}H_{25}N_5$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	N	0	0
			23	18	5		
4	B	1	Total	C	N	0	0
			23	18	5		

- Molecule 5 is ACETATE ION (three-letter code: ACT) (formula:  $C_2H_3O_2$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			4	2	2		
5	B	1	Total	C	O	0	0
			4	2	2		

- Molecule 6 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	1	Total 1	Zn 1	0	0


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	214	Total 214	O 214	0	0
7	B	246	Total 246	O 246	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: NITRIC OXIDE SYNTHASE, BRAIN



- Molecule 1: NITRIC OXIDE SYNTHASE, BRAIN



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	51.70Å 110.71Å 164.17Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	49.31 – 1.85 49.31 – 1.85	Depositor EDS
% Data completeness (in resolution range)	99.5 (49.31-1.85) 99.7 (49.31-1.85)	Depositor EDS
$R_{merge}$	0.12	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.36 (at 1.86Å)	Xtrriage
Refinement program	PHENIX (PHENIX.REFINE)	Depositor
R, $R_{free}$	0.193 , 0.233 0.190 , 0.231	Depositor DCC
$R_{free}$ test set	4008 reflections (4.94%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	30.5	Xtrriage
Anisotropy	1.103	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.37 , 49.8	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	7322	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	40.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.77% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: HEM, ZN, H4B, W64, ACT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.38	0/3429	0.52	0/4654
1	B	0.39	0/3465	0.51	0/4697
All	All	0.38	0/6894	0.51	0/9351

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3327	0	3241	24	0
1	B	3360	0	3282	25	0
2	A	43	0	30	3	0
2	B	43	0	30	4	0
3	A	17	0	15	0	0
3	B	17	0	15	1	0
4	A	23	0	0	0	0
4	B	23	0	0	0	0
5	A	4	0	3	0	0
5	B	4	0	3	0	0
6	A	1	0	0	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
7	A	214	0	0	1	0
7	B	246	0	0	3	0
All	All	7322	0	6619	49	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (49) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:523:LEU:HD22	1:A:531:PRO:HB2	1.63	0.79
2:A:750:HEM:HMC2	2:A:750:HEM:HBC2	1.68	0.73
1:B:478:GLN:HB2	1:B:481:ARG:HG3	1.73	0.71
2:B:750:HEM:HBC2	2:B:750:HEM:HMC2	1.72	0.71
1:A:475:TRP:HB2	1:A:523:LEU:HB3	1.73	0.69
1:B:349:ARG:HG2	1:B:353:GLN:HG3	1.76	0.67
1:A:330:ILE:HD11	1:B:696:LEU:HD22	1.79	0.64
1:A:480:ILE:HD13	1:A:541:VAL:HG13	1.82	0.62
2:A:750:HEM:HBB2	2:A:750:HEM:HHC	1.83	0.61
1:A:706:TYR:OH	2:A:750:HEM:O2D	2.17	0.59
1:A:632:GLU:OE2	1:B:628:GLN:NE2	2.36	0.58
1:A:478:GLN:HB2	1:A:481:ARG:HG3	1.86	0.57
2:B:750:HEM:O2A	7:B:2245:HOH:O	2.19	0.53
1:A:475:TRP:CZ2	1:A:531:PRO:HG3	2.46	0.51
1:A:330:ILE:HD11	1:B:696:LEU:HB3	1.92	0.51
1:A:455:LEU:HD12	1:A:587:TRP:HB3	1.94	0.49
1:B:386:LYS:HA	1:B:389:GLU:HG2	1.95	0.49
1:A:628:GLN:NE2	1:B:632:GLU:OE2	2.43	0.48
1:B:403:TYR:CE2	1:B:407:HIS:CE1	3.01	0.48
2:B:750:HEM:CGA	3:B:760:H4B:HN22	2.27	0.47
1:A:628:GLN:HG3	1:B:631:VAL:HG11	1.97	0.46
1:A:525:GLN:HG3	1:A:529:ASN:O	2.16	0.46
1:A:322:LEU:HB2	1:A:699:ARG:HB2	1.98	0.46
1:B:595:VAL:O	1:B:599:CYS:HB2	2.16	0.45
1:B:325:GLY:O	1:B:332:MET:HG3	2.15	0.45
1:B:510:TRP:CE2	1:B:521:PRO:HD3	2.51	0.45
1:A:609:GLU:HG3	7:A:2167:HOH:O	2.16	0.45
1:A:596:ARG:O	1:A:600:ASP:HB2	2.17	0.45
1:B:659:ILE:O	1:B:663:GLU:HG3	2.17	0.44
1:A:355:PHE:CE1	1:A:385:ASN:HB2	2.54	0.43
1:B:351:LYS:HB3	1:B:351:LYS:HE2	1.76	0.43

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:403:TYR:CE1	1:A:407:HIS:CE1	3.07	0.43
1:B:409:TRP:CE3	1:B:421:TRP:HA	2.54	0.42
1:B:336:MET:HE2	1:B:336:MET:HB2	1.87	0.42
1:A:709:ASP:HA	1:A:710:PRO:HD3	1.87	0.42
1:B:302:LYS:HE3	1:B:302:LYS:HB2	1.81	0.42
2:B:750:HEM:O1A	7:B:2244:HOH:O	2.21	0.42
1:A:436:HIS:CE1	1:A:534:PHE:HE2	2.38	0.42
1:A:569:ASN:H	1:A:569:ASN:HD22	1.67	0.42
1:B:527:ASN:ND2	7:B:2131:HOH:O	2.53	0.42
1:B:485:TYR:CE2	1:B:514:ARG:HA	2.55	0.41
1:B:680:VAL:HA	1:B:681:PRO:HD3	1.91	0.41
1:A:694:GLU:HB3	1:B:335:ILE:HD13	2.02	0.41
1:B:462:PHE:HB2	1:B:581:ALA:HB3	2.03	0.41
1:A:680:VAL:HA	1:A:681:PRO:HD3	1.92	0.41
1:B:569:ASN:H	1:B:569:ASN:HD22	1.66	0.41
1:A:510:TRP:CE2	1:A:521:PRO:HD3	2.56	0.41
1:B:304:LYS:HD3	1:B:306:TRP:CE2	2.56	0.40
1:B:475:TRP:HB2	1:B:523:LEU:HB3	2.04	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	407/422 (96%)	392 (96%)	13 (3%)	2 (0%)	29	15
1	B	411/422 (97%)	403 (98%)	8 (2%)	0	100	100
All	All	818/844 (97%)	795 (97%)	21 (3%)	2 (0%)	47	33

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	514	ARG
1	A	716	TRP

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	366/377 (97%)	353 (96%)	13 (4%)	35	18
1	B	370/377 (98%)	358 (97%)	12 (3%)	39	22
All	All	736/754 (98%)	711 (97%)	25 (3%)	39	19

All (25) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	328	GLU
1	A	381	LEU
1	A	390	SER
1	A	476	ASN
1	A	523	LEU
1	A	547	ARG
1	A	550	LYS
1	A	555	LYS
1	A	569	ASN
1	A	601	ASN
1	A	645	LYS
1	A	713	THR
1	A	715	VAL
1	B	337	LEU
1	B	353	GLN
1	B	382	GLU
1	B	423	LYS
1	B	540	LEU
1	B	547	ARG
1	B	569	ASN
1	B	602[A]	SER
1	B	602[B]	SER

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	668[A]	CYS
1	B	668[B]	CYS
1	B	699	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	569	ASN
1	B	353	GLN
1	B	527	ASN
1	B	569	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 9 ligands modelled in this entry, 1 is monoatomic - leaving 8 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
5	ACT	A	860	-	1,3,3	1.16	0	0,3,3	0.00	-

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	W64	A	800	-	25,25,25	0.38	0	33,34,34	2.00	10 (30%)
4	W64	B	800	-	25,25,25	0.43	0	33,34,34	2.08	9 (27%)
5	ACT	B	860	-	1,3,3	1.13	0	0,3,3	0.00	-
3	H4B	B	760	-	16,18,18	1.02	1 (6%)	11,26,26	2.57	5 (45%)
2	HEM	B	750	1	27,50,50	2.11	6 (22%)	17,82,82	1.57	3 (17%)
2	HEM	A	750	1	27,50,50	2.18	7 (25%)	17,82,82	1.98	5 (29%)
3	H4B	A	760	-	16,18,18	0.80	0	11,26,26	2.65	5 (45%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	W64	A	800	-	-	4/9/19/19	0/3/3/3
4	W64	B	800	-	-	2/9/19/19	0/3/3/3
3	H4B	B	760	-	-	0/8/17/17	0/2/2/2
2	HEM	B	750	1	-	1/6/54/54	-
2	HEM	A	750	1	-	1/6/54/54	-
3	H4B	A	760	-	-	0/8/17/17	0/2/2/2

All (14) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	750	HEM	C3D-C2D	5.13	1.52	1.37
2	B	750	HEM	C3D-C2D	4.99	1.52	1.37
2	A	750	HEM	C3B-C2B	-4.87	1.33	1.40
2	B	750	HEM	C3B-C2B	-4.56	1.34	1.40
2	A	750	HEM	C3B-CAB	3.72	1.55	1.47
2	A	750	HEM	C3C-CAC	3.71	1.55	1.47
2	A	750	HEM	C3C-C2C	-3.71	1.35	1.40
2	B	750	HEM	C3B-CAB	3.64	1.55	1.47
2	B	750	HEM	C3C-CAC	3.58	1.55	1.47
2	B	750	HEM	C3C-C2C	-3.31	1.35	1.40
3	B	760	H4B	C4-C4A	-2.45	1.38	1.41
2	A	750	HEM	C1D-ND	2.21	1.40	1.36
2	B	750	HEM	C4B-NB	2.16	1.40	1.36
2	A	750	HEM	CAA-C2A	2.02	1.55	1.52

All (37) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	760	H4B	C4-C4A-C8A	5.85	119.77	114.57
4	B	800	W64	C02-N01-C06	5.64	122.38	118.10
2	A	750	HEM	CBD-CAD-C3D	-5.52	102.31	112.48
3	A	760	H4B	C4-C4A-C8A	5.40	119.36	114.57
4	A	800	W64	C02-N01-C06	4.69	121.65	118.10
4	A	800	W64	C22-N21-C15	-4.36	106.31	118.09
4	B	800	W64	C22-N21-C15	-4.35	106.35	118.09
4	B	800	W64	C26-N21-C15	-4.01	107.26	118.09
4	A	800	W64	C26-N21-C15	-3.82	107.78	118.09
4	B	800	W64	C08-C06-N01	3.81	121.62	115.95
4	B	800	W64	C05-C06-N01	-3.65	119.03	122.90
2	A	750	HEM	C1D-C2D-C3D	-3.46	104.59	107.00
3	B	760	H4B	N3-C2-N1	-3.29	120.27	125.42
3	A	760	H4B	N3-C2-N1	-3.25	120.32	125.42
4	A	800	W64	C22-C23-N24	-3.24	107.15	110.80
4	A	800	W64	C05-C06-N01	-3.15	119.56	122.90
4	A	800	W64	C08-C06-N01	3.11	120.59	115.95
3	B	760	H4B	C4-N3-C2	3.08	120.83	115.93
3	A	760	H4B	C2-N1-C8A	3.02	121.31	114.54
2	B	750	HEM	CBD-CAD-C3D	-3.01	106.94	112.48
3	A	760	H4B	C4-C4A-N5	2.99	121.63	119.12
3	A	760	H4B	C4-N3-C2	2.98	120.66	115.93
2	B	750	HEM	CAA-CBA-CGA	-2.93	107.76	112.67
4	B	800	W64	C25-N24-C23	-2.71	105.73	109.52
3	B	760	H4B	C2-N1-C8A	2.68	120.54	114.54
4	A	800	W64	C25-N24-C23	-2.65	105.82	109.52
4	A	800	W64	C16-N11-C12	2.63	121.07	117.48
4	A	800	W64	C14-C13-C12	2.61	119.23	116.71
2	A	750	HEM	CBA-CAA-C2A	-2.37	108.11	112.49
4	A	800	W64	N02-C02-N01	2.19	119.95	116.49
4	B	800	W64	C16-N11-C12	2.18	120.46	117.48
2	A	750	HEM	CAD-CBD-CGD	-2.17	109.04	112.67
4	B	800	W64	C22-C23-N24	-2.16	108.37	110.80
4	B	800	W64	C09-C13-C12	-2.11	118.43	121.83
2	B	750	HEM	C1D-C2D-C3D	-2.09	105.55	107.00
2	A	750	HEM	C4C-C3C-C2C	2.08	108.35	106.90
3	B	760	H4B	N2-C2-N3	2.03	120.41	117.25

There are no chirality outliers.

All (8) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	750	HEM	C2A-CAA-CBA-CGA

*Continued on next page...*

*Continued from previous page...*


Mol	Chain	Res	Type	Atoms
2	A	750	HEM	C2A-CAA-CBA-CGA
4	A	800	W64	C16-C15-N21-C22
4	A	800	W64	C14-C15-N21-C22
4	A	800	W64	C08-C09-C13-C12
4	B	800	W64	C08-C09-C13-C14
4	B	800	W64	C08-C09-C13-C12
4	A	800	W64	C08-C09-C13-C14


There are no ring outliers.


3 monomers are involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	760	H4B	1	0
2	B	750	HEM	4	0
2	A	750	HEM	3	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	408/422 (96%)	0.32	15 (3%) 41 39	16, 41, 83, 117	1 (0%)
1	B	411/422 (97%)	0.12	11 (2%) 54 53	15, 32, 60, 94	0
All	All	819/844 (97%)	0.22	26 (3%) 47 45	15, 36, 78, 117	1 (0%)

All (26) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	717	LYS	16.6
1	A	715	VAL	5.8
1	A	716	TRP	4.5
1	A	355	PHE	3.8
1	A	554	PHE	3.8
1	B	352	ASP	3.8
1	A	352	ASP	3.6
1	B	718	GLY	3.6
1	B	300	PHE	3.6
1	B	348	VAL	3.6
1	B	350	THR	3.5
1	A	551	PHE	3.2
1	B	349	ARG	2.7
1	A	321	THR	2.6
1	A	338	PRO	2.5
1	B	389	GLU	2.5
1	A	351	LYS	2.4
1	A	300	PHE	2.4
1	A	550	LYS	2.3
1	A	322	LEU	2.2
1	A	713	THR	2.2
1	B	385	ASN	2.1
1	A	552	ASP	2.1
1	B	351	LYS	2.1

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	668[A]	CYS	2.1
1	B	355	PHE	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
5	ACT	B	860	4/4	0.87	0.19	47,49,49,49	0
4	W64	B	800	23/23	0.88	0.28	18,79,97,99	0
3	H4B	B	760	17/17	0.89	0.11	22,28,32,34	0
3	H4B	A	760	17/17	0.91	0.12	27,32,40,40	0
5	ACT	A	860	4/4	0.93	0.34	50,51,52,53	0
4	W64	A	800	23/23	0.93	0.27	19,78,99,100	0
2	HEM	A	750	43/43	0.98	0.12	7,21,48,55	0
2	HEM	B	750	43/43	0.98	0.13	12,21,50,65	0
6	ZN	A	900	1/1	0.98	0.09	25,25,25,25	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

**Electron density around W64 B 800:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)




**Electron density around W64 A 800:**


$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



**Electron density around HEM A 750:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





## 6.5 Other polymers [i](#)

There are no such residues in this entry.