



# Full wwPDB X-ray Structure Validation Report ⓘ

May 16, 2020 – 12:04 pm BST

PDB ID : 5G1V  
Title : Linalool Dehydratase Isomerase: Selenomethionine Derivative  
Authors : Chambers, S.; Hau, A.; Man, H.; Omar, M.; Turkenburg, J.P.; Grogan, G.  
Deposited on : 2016-03-30  
Resolution : 2.68 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.68 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	3863 (2.70-2.66)
Clashscore	141614	4210 (2.70-2.66)
Ramachandran outliers	138981	4141 (2.70-2.66)
Sidechain outliers	138945	4141 (2.70-2.66)
RSRZ outliers	127900	3780 (2.70-2.66)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	372	 90% 6% . .
1	B	372	 90% 6% . . .
1	C	372	 90% 6% . .
1	D	372	 91% 5% . .
1	E	372	 90% 5% . . .

## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 15049 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called LINALOOL DEHYDRATASE ISOMERASE.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	S	Se			
1	A	362	Total 2875	C 1859	N 480	O 523	S 4	Se 9	0	0	0
1	B	363	Total 2877	C 1860	N 479	O 525	S 4	Se 9	0	0	0
1	C	363	Total 2891	C 1868	N 482	O 528	S 4	Se 9	0	0	0
1	D	362	Total 2866	C 1855	N 475	O 523	S 4	Se 9	0	0	0
1	E	362	Total 2885	C 1864	N 485	O 523	S 4	Se 9	0	0	0

There are 5 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MSE	-	expression tag	UNP E1XUJ2
B	1	MSE	-	expression tag	UNP E1XUJ2
C	1	MSE	-	expression tag	UNP E1XUJ2
D	1	MSE	-	expression tag	UNP E1XUJ2
E	1	MSE	-	expression tag	UNP E1XUJ2


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	123	Total 123	O 123	0	0
2	B	131	Total 131	O 131	0	0
2	C	147	Total 147	O 147	0	0
2	D	123	Total 123	O 123	0	0
2	E	131	Total 131	O 131	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: LINALOOL DEHYDRATASE ISOMERASE




- Molecule 1: LINALOOL DEHYDRATASE ISOMERASE



- Molecule 1: LINALOOL DEHYDRATASE ISOMERASE




- Molecule 1: LINALOOL DEHYDRATASE ISOMERASE



- Molecule 1: LINALOOL DEHYDRATASE ISOMERASE





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	100.51Å 106.72Å 222.81Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	111.41 – 2.68 96.25 – 2.68	Depositor EDS
% Data completeness (in resolution range)	99.8 (111.41-2.68) 99.8 (96.25-2.68)	Depositor EDS
$R_{merge}$	0.24	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.58 (at 2.69Å)	Xtrriage
Refinement program	REFMAC 5.8.0131	Depositor
R, $R_{free}$	0.168 , 0.203 0.180 , 0.218	Depositor DCC
$R_{free}$ test set	3290 reflections (4.84%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	25.9	Xtrriage
Anisotropy	1.012	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 24.6	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	15049	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	27.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.65% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup> Intensities estimated from amplitudes.

<sup>2</sup> Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality i

### 5.1 Standard geometry i

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.72	0/2952	0.87	7/4003 (0.2%)
1	B	0.72	1/2954 (0.0%)	0.89	10/4007 (0.2%)
1	C	0.73	0/2968	0.85	4/4023 (0.1%)
1	D	0.72	0/2943	0.85	5/3993 (0.1%)
1	E	0.72	0/2962	0.87	9/4015 (0.2%)
All	All	0.72	1/14779 (0.0%)	0.87	35/20041 (0.2%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	122	GLU	CD-OE2	-5.25	1.19	1.25

All (35) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	E	8	ARG	NE-CZ-NH2	-8.86	115.87	120.30
1	A	8	ARG	NE-CZ-NH2	-8.79	115.91	120.30
1	D	8	ARG	NE-CZ-NH2	-8.55	116.02	120.30
1	B	8	ARG	NE-CZ-NH2	-7.86	116.37	120.30
1	A	8	ARG	NE-CZ-NH1	7.73	124.16	120.30
1	E	8	ARG	NE-CZ-NH1	7.37	123.99	120.30
1	D	8	ARG	NE-CZ-NH1	6.96	123.78	120.30
1	E	8	ARG	CG-CD-NE	-6.84	97.43	111.80
1	E	155	ARG	NE-CZ-NH2	-6.76	116.92	120.30
1	B	219	ARG	NE-CZ-NH1	6.72	123.66	120.30
1	A	214	LEU	CB-CG-CD2	6.54	122.11	111.00
1	E	214	LEU	CB-CG-CD2	6.42	121.91	111.00
1	C	327	SER	O-C-N	6.36	132.87	122.70
1	B	219	ARG	NE-CZ-NH2	-6.30	117.15	120.30
1	D	8	ARG	CG-CD-NE	-6.23	98.71	111.80
1	A	8	ARG	CG-CD-NE	-6.10	99.00	111.80
1	C	61	GLN	CA-CB-CG	6.09	126.81	113.40
1	B	155	ARG	NE-CZ-NH2	-6.08	117.26	120.30

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	8	ARG	NE-CZ-NH1	6.04	123.32	120.30
1	A	61	GLN	CA-CB-CG	6.00	126.59	113.40
1	B	61	GLN	CA-CB-CG	5.98	126.55	113.40
1	E	273	ASP	CB-CG-OD1	-5.89	113.00	118.30
1	D	61	GLN	CA-CB-CG	5.87	126.31	113.40
1	E	61	GLN	CA-CB-CG	5.59	125.70	113.40
1	D	248	MSE	CA-CB-CG	-5.58	103.81	113.30
1	B	39	ASP	C-N-CA	5.40	135.21	121.70
1	B	8	ARG	CG-CD-NE	-5.36	100.55	111.80
1	C	248	MSE	CA-CB-CG	-5.32	104.25	113.30
1	E	248	MSE	CA-CB-CG	-5.31	104.28	113.30
1	C	179	GLN	CB-CA-C	-5.22	99.97	110.40
1	B	4	LEU	N-CA-C	5.14	124.87	111.00
1	A	331	ALA	N-CA-C	-5.07	97.31	111.00
1	A	193	ARG	CG-CD-NE	-5.04	101.23	111.80
1	B	279	ARG	NE-CZ-NH1	5.03	122.82	120.30
1	E	179	GLN	CB-CA-C	-5.00	100.39	110.40

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2875	0	2767	11	0
1	B	2877	0	2758	8	0
1	C	2891	0	2786	14	0
1	D	2866	0	2744	9	0
1	E	2885	0	2782	13	0
2	A	123	0	0	3	0
2	B	131	0	0	2	0
2	C	147	0	0	6	0
2	D	123	0	0	1	0
2	E	131	0	0	5	0
All	All	15049	0	13837	52	0



The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (52) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:121:LYS:HG2	1:E:122:GLU:HG2	1.35	1.06
1:A:172:GLU:OE2	2:A:2034:HOH:O	1.82	0.95
1:D:329:VAL:O	2:D:2116:HOH:O	1.83	0.94
1:E:121:LYS:HG2	1:E:122:GLU:CG	2.12	0.77
1:E:121:LYS:C	1:E:122:GLU:HG2	2.05	0.77
1:E:125:MSE:HE2	2:E:2032:HOH:O	1.89	0.72
1:A:216:ASP:OD2	1:A:219:ARG:NH1	2.24	0.70
1:C:121:LYS:O	2:C:2054:HOH:O	2.10	0.69
1:A:337:HIS:HB3	2:A:2118:HOH:O	1.96	0.66
1:B:291:GLY:HA3	1:D:330:SER:OG	1.98	0.63
1:C:125:MSE:HE2	2:C:2036:HOH:O	2.02	0.59
1:D:79:ALA:HB3	1:D:362:MSE:HE1	1.86	0.58
1:C:121:LYS:HG3	1:C:122:GLU:HG2	1.87	0.56
1:E:121:LYS:CG	1:E:122:GLU:HG2	2.22	0.56
1:C:8:ARG:HD2	2:C:2001:HOH:O	2.06	0.55
1:E:193:ARG:NH2	1:E:362:MSE:O	2.39	0.55
1:D:193:ARG:NH2	1:D:362:MSE:O	2.39	0.55
1:A:193:ARG:NH2	1:A:362:MSE:O	2.38	0.55
1:C:8:ARG:NH2	1:C:314:ASP:OD1	2.39	0.55
1:A:125:MSE:HE1	2:A:2032:HOH:O	2.08	0.54
1:C:193:ARG:NH2	1:C:362:MSE:O	2.41	0.53
1:C:121:LYS:C	1:C:122:GLU:HG2	2.28	0.53
1:C:250:HIS:HD2	1:C:306:GLU:OE2	1.91	0.53
1:A:219:ARG:HD2	1:A:237:ILE:HD13	1.92	0.52
1:D:250:HIS:HD2	1:D:306:GLU:OE2	1.92	0.52
1:E:174:ASP:N	2:E:2069:HOH:O	2.42	0.52
1:E:250:HIS:HD2	1:E:306:GLU:OE2	1.93	0.52
1:B:121:LYS:C	1:B:122:GLU:HG2	2.28	0.52
1:B:250:HIS:HD2	1:B:306:GLU:OE2	1.93	0.51
1:C:155:ARG:HG2	2:C:2072:HOH:O	2.09	0.51
1:A:250:HIS:HD2	1:A:306:GLU:OE2	1.94	0.51
1:C:337:HIS:HB3	2:C:2142:HOH:O	2.11	0.51
1:B:125:MSE:HE2	2:B:2038:HOH:O	2.10	0.50
1:A:91:HIS:CE1	1:D:227:HIS:CD2	3.00	0.50
1:E:327:SER:HB3	2:E:2122:HOH:O	2.12	0.49
1:C:125:MSE:HE1	2:C:2034:HOH:O	2.14	0.46
1:E:8:ARG:NH2	1:E:314:ASP:OD1	2.49	0.46

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:8:ARG:NH2	1:D:314:ASP:OD1	2.49	0.45
1:C:121:LYS:HG3	1:C:122:GLU:CG	2.45	0.45
1:B:8:ARG:NH2	1:B:314:ASP:OD1	2.50	0.45
1:A:227:HIS:CD2	1:C:91:HIS:CE1	3.06	0.44
1:E:266:GLN:HB2	2:E:2094:HOH:O	2.17	0.44
1:D:79:ALA:CB	1:D:362:MSE:HE1	2.46	0.44
1:E:8:ARG:HD2	2:E:2002:HOH:O	2.17	0.44
1:B:125:MSE:HE1	2:B:2036:HOH:O	2.19	0.43
1:B:325:LYS:N	1:B:326:PRO:CD	2.83	0.41
1:B:214:LEU:HD11	1:B:226:TYR:HB2	2.01	0.41
1:E:325:LYS:N	1:E:326:PRO:CD	2.83	0.41
1:A:8:ARG:NH2	1:A:314:ASP:OD1	2.51	0.41
1:A:39:ASP:HA	1:A:45:TYR:CE1	2.56	0.41
1:D:18:GLN:HE21	1:D:24:VAL:HA	1.86	0.41
1:C:18:GLN:HE21	1:C:25:THR:H	1.69	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	360/372 (97%)	350 (97%)	8 (2%)	2 (1%)	25 47
1	B	361/372 (97%)	350 (97%)	11 (3%)	0	100 100
1	C	361/372 (97%)	352 (98%)	7 (2%)	2 (1%)	25 47
1	D	360/372 (97%)	351 (98%)	8 (2%)	1 (0%)	41 64
1	E	360/372 (97%)	351 (98%)	6 (2%)	3 (1%)	19 40
All	All	1802/1860 (97%)	1754 (97%)	40 (2%)	8 (0%)	34 58

All (8) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	E	174	ASP
1	A	212	LYS
1	C	122	GLU
1	D	212	LYS
1	E	212	LYS
1	A	330	SER
1	C	212	LYS
1	E	122	GLU

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	293/293 (100%)	280 (96%)	13 (4%)	28	53
1	B	292/293 (100%)	278 (95%)	14 (5%)	25	49
1	C	296/293 (101%)	284 (96%)	12 (4%)	30	56
1	D	290/293 (99%)	280 (97%)	10 (3%)	37	63
1	E	294/293 (100%)	280 (95%)	14 (5%)	25	49
All	All	1465/1465 (100%)	1402 (96%)	63 (4%)	29	54

All (63) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	4	LEU
1	A	8	ARG
1	A	46	SER
1	A	61	GLN
1	A	80	LEU
1	A	98	SER
1	A	121	LYS
1	A	179	GLN
1	A	204	ARG
1	A	214	LEU
1	A	219	ARG
1	A	240	TYR

*Continued on next page...*

*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Res</b>	<b>Type</b>
1	A	330	SER
1	B	4	LEU
1	B	8	ARG
1	B	40	PHE
1	B	46	SER
1	B	61	GLN
1	B	80	LEU
1	B	98	SER
1	B	122	GLU
1	B	179	GLN
1	B	214	LEU
1	B	218	GLU
1	B	240	TYR
1	B	327	SER
1	B	330	SER
1	C	4	LEU
1	C	8	ARG
1	C	46	SER
1	C	61	GLN
1	C	80	LEU
1	C	98	SER
1	C	122	GLU
1	C	179	GLN
1	C	214	LEU
1	C	240	TYR
1	C	328	ILE
1	C	330	SER
1	D	4	LEU
1	D	8	ARG
1	D	46	SER
1	D	61	GLN
1	D	98	SER
1	D	121	LYS
1	D	179	GLN
1	D	214	LEU
1	D	240	TYR
1	D	330	SER
1	E	4	LEU
1	E	8	ARG
1	E	46	SER
1	E	61	GLN
1	E	80	LEU

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	E	98	SER
1	E	121	LYS
1	E	122	GLU
1	E	155	ARG
1	E	179	GLN
1	E	204	ARG
1	E	214	LEU
1	E	240	TYR
1	E	330	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (29) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	18	GLN
1	A	61	GLN
1	A	211	GLN
1	A	227	HIS
1	A	250	HIS
1	A	310	GLN
1	B	18	GLN
1	B	61	GLN
1	B	211	GLN
1	B	227	HIS
1	B	250	HIS
1	B	310	GLN
1	C	18	GLN
1	C	61	GLN
1	C	211	GLN
1	C	227	HIS
1	C	250	HIS
1	D	18	GLN
1	D	61	GLN
1	D	211	GLN
1	D	227	HIS
1	D	250	HIS
1	D	310	GLN
1	E	18	GLN
1	E	58	HIS
1	E	61	GLN
1	E	227	HIS
1	E	250	HIS
1	E	310	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

### 5.7 Other polymers [i](#)

There are no such residues in this entry.

### 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	353/372 (94%)	-0.30	0 <a href="#">100</a>   <a href="#">100</a>	18, 26, 44, 64	0
1	B	354/372 (95%)	-0.31	1 (0%) <a href="#">94</a>   <a href="#">95</a>	17, 25, 44, 70	0
1	C	354/372 (95%)	-0.39	0 <a href="#">100</a>   <a href="#">100</a>	16, 25, 42, 66	0
1	D	353/372 (94%)	-0.26	0 <a href="#">100</a>   <a href="#">100</a>	16, 26, 40, 54	0
1	E	353/372 (94%)	-0.33	1 (0%) <a href="#">94</a>   <a href="#">95</a>	16, 27, 43, 67	0
All	All	1767/1860 (95%)	-0.32	2 (0%) <a href="#">95</a>   <a href="#">96</a>	16, 26, 43, 70	0

All (2) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	365	PRO	2.2
1	E	4	LEU	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers

There are no such residues in this entry.