Full wwPDB X-ray Structure Validation Report (i)

Sep 6, 2023 – 07:22 PM EDT

PDB ID	:	4G8T
Title	:	Crystal structure of a glucarate dehydratase related protein, from actinobacillus succinogenes, target EFI-502312, with sodium and sulfate bound, ordered loop
Authors	:	Vetting, M.W.; Toro, R.; Bhosle, R.; Al Obaidi, N.F.; Morisco, L.L.; Wasserman, S.R.; Sojitra, S.; Washington, E.; Scott Glenn, A.; Chowdhury, S.; Evans, B.; Hammonds, J.; Hillerich, B.; Love, J.; Seidel, R.D.; Imker, H.J.; Gerlt, J.A.; Almo, S.C.; Enzyme Function Initiative (EFI)
Deposited on	:	2012-07-23
Resolution	:	1.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.35
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 1.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4298 (1.70-1.70)
Clashscore	141614	4695 (1.70-1.70)
Ramachandran outliers	138981	4610 (1.70-1.70)
Sidechain outliers	138945	4610 (1.70-1.70)
RSRZ outliers	127900	4222 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	464	% 93% • 5%
1	B	464	% 90% 5% 5%
1	C	464	2% 92% • 5%

Continued on next page...

Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	D	464	2% 91% 5%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
6	DTT	D	503	-	-	X	-

2 Entry composition (i)

There are 7 unique types of molecules in this entry. The entry contains 15933 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Glucarate dehydratase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	441	3448	2176	598	653	21	0	3	0
1	B	441	3474	2191	601	661	21	0	6	0
1	C	441	3458	2181	601	655	21	0	4	0
1	D	441	3482	2196	602	663	21	0	7	0

There are 88 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-21	MET	-	initiating methionine	UNP A6VQF1
A	-20	HIS	-	expression tag	UNP A6VQF1
A	-19	HIS	-	expression tag	UNP A6VQF1
A	-18	HIS	-	expression tag	UNP A6VQF1
A	-17	HIS	-	expression tag	UNP A6VQF1
A	-16	HIS	-	expression tag	UNP A6VQF1
A	-15	HIS	-	expression tag	UNP A6VQF1
A	-14	SER	-	expression tag	UNP A6VQF1
A	-13	SER	-	expression tag	UNP A6VQF1
A	-12	GLY	-	expression tag	UNP A6VQF1
A	-11	VAL	-	expression tag	UNP A6VQF1
A	-10	ASP	-	expression tag	UNP A6VQF1
A	-9	LEU	-	expression tag	UNP A6VQF1
A	-8	GLY	-	expression tag	UNP A6VQF1
A	-7	THR	-	expression tag	UNP A6VQF1
A	-6	GLU	-	expression tag	UNP A6VQF1
A	-5	ASN	-	expression tag	UNP A6VQF1
A	-4	LEU	-	expression tag	UNP A6VQF1
A	-3	TYR	-	expression tag	UNP A6VQF1
A	-2	PHE	-	expression tag	UNP A6VQF1
A	-1	GLN	-	expression tag	UNP A6VQF1

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	SER	-	expression tag	UNP A6VQF1
B	-21	MET	-	initiating methionine	UNP A6VQF1
B	-20	HIS	-	expression tag	UNP A6VQF1
B	-19	HIS	-	expression tag	UNP A6VQF1
B	-18	HIS	-	expression tag	UNP A6VQF1
B	-17	HIS	-	expression tag	UNP A6VQF1
B	-16	HIS	-	expression tag	UNP A6VQF1
B	-15	HIS	-	expression tag	UNP A6VQF1
B	-14	SER	-	expression tag	UNP A6VQF1
B	-13	SER	-	expression tag	UNP A6VQF1
B	-12	GLY	-	expression tag	UNP A6VQF1
B	-11	VAL	-	expression tag	UNP A6VQF1
B	-10	ASP	-	expression tag	UNP A6VQF1
B	-9	LEU	-	expression tag	UNP A6VQF1
B	-8	GLY	-	expression tag	UNP A6VQF1
B	-7	THR	-	expression tag	UNP A6VQF1
B	-6	GLU	-	expression tag	UNP A6VQF1
B	-5	ASN	-	expression tag	UNP A6VQF1
B	-4	LEU	-	expression tag	UNP A6VQF1
B	-3	TYR	-	expression tag	UNP A6VQF1
B	-2	PHE	-	expression tag	UNP A6VQF1
B	-1	GLN	-	expression tag	UNP A6VQF1
B	0	SER	-	expression tag	UNP A6VQF1
C	-21	MET	-	initiating methionine	UNP A6VQF1
C	-20	HIS	-	expression tag	UNP A6VQF1
C	-19	HIS	-	expression tag	UNP A6VQF1
C	-18	HIS	-	expression tag	UNP A6VQF1
C	-17	HIS	-	expression tag	UNP A6VQF1
C	-16	HIS	-	expression tag	UNP A6VQF1
C	-15	HIS	-	expression tag	UNP A6VQF1
C	-14	SER	-	expression tag	UNP A6VQF1
C	-13	SER	-	expression tag	UNP A6VQF1
C	-12	GLY	-	expression tag	UNP A6VQF1
C	-11	VAL	-	expression tag	UNP A6VQF1
C	-10	ASP	-	expression tag	UNP A6VQF1
C	-9	LEU	-	expression tag	UNP A6VQF1
C	-8	GLY	-	expression tag	UNP A6VQF1
C	-7	THR	-	expression tag	UNP A6VQF1
C	-6	GLU	-	expression tag	UNP A6VQF1
C	-5	ASN	-	expression tag	UNP A6VQF1
C	-4	LEU	-	expression tag	UNP A6VQF1
C	-3	TYR	-	expression tag	UNP A6VQF1

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	-2	PHE	-	expression tag	UNP A6VQF1
C	-1	GLN	-	expression tag	UNP A6VQF1
C	0	SER	-	expression tag	UNP A6VQF1
D	-21	MET	-	initiating methionine	UNP A6VQF1
D	-20	HIS	-	expression tag	UNP A6VQF1
D	-19	HIS	-	expression tag	UNP A6VQF1
D	-18	HIS	-	expression tag	UNP A6VQF1
D	-17	HIS	-	expression tag	UNP A6VQF1
D	-16	HIS	-	expression tag	UNP A6VQF1
D	-15	HIS	-	expression tag	UNP A6VQF1
D	-14	SER	-	expression tag	UNP A6VQF1
D	-13	SER	-	expression tag	UNP A6VQF1
D	-12	GLY	-	expression tag	UNP A6VQF1
D	-11	VAL	-	expression tag	UNP A6VQF1
D	-10	ASP	-	expression tag	UNP A6VQF1
D	-9	LEU	-	expression tag	UNP A6VQF1
D	-8	GLY	-	expression tag	UNP A6VQF1
D	-7	THR	-	expression tag	UNP A6VQF1
D	-6	GLU	-	expression tag	UNP A6VQF1
D	-5	ASN	-	expression tag	UNP A6VQF1
D	-4	LEU	-	expression tag	UNP A6VQF1
D	-3	TYR	-	expression tag	UNP A6VQF1
D	-2	PHE	-	expression tag	UNP A6VQF1
D	-1	GLN	-	expression tag	UNP A6VQF1
D	0	SER	-	expression tag	UNP A6VQF1

- Molecule 2 is SULFATE ION (three-letter code: SO4) (formula: O_4S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total O S 5 4 1	0	0
2	B	1	Total O S 5 4 1	0	0
2	C	1	Total O S 5 4 1	0	0
2	D	1	Total O S 5 4 1	0	0

- Molecule 3 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total Na 1 1	0	0
3	B	1	Total Na 1 1	0	0
3	C	1	Total Na 1 1	0	0
3	D	1	Total Na 1 1	0	0

- Molecule 4 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C O 6 3 3	0	0
4	A	1	Total C O 6 3 3	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	O			
4	B	1	6	3	3		0	0
			Total	C	O			
4	B	1	6	3	3		0	0
			Total	C	O			
4	B	1	6	3	3		0	0
			Total	C	O			
4	B	1	6	3	3		0	0
			Total	C	O			
4	C	1	6	3	3		0	0

- Molecule 5 is (2R,3S)-1,4-DIMERCAPTOBUTANE-2,3-DIOL (three-letter code: DTU) (formula: $C_4H_{10}O_2S_2$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	O	S		
5	A	1	8	4	2	2	0	0

- Molecule 6 is 2,3-DIHYDROXY-1,4-DITHIOBUTANE (three-letter code: DTT) (formula: $C_4H_{10}O_2S_2$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	O	S		
6	B	1	8	4	2	2	0	0
			Total	C	O	S		
6	C	1	8	4	2	2	0	0
			Total	C	O	S		
6	D	1	8	4	2	2	0	0

- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
7	A	525	525	525	0	0
			Total	O		
7	B	503	503	503	0	0
			Total	O		
7	C	459	459	459	0	0
			Total	O		
7	D	486	486	486	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Glucarate dehydratase

Chain A:





• Molecule 1: Glucarate dehydratase

Chain B:





• Molecule 1: Glucarate dehydratase

Chain C:





• Molecule 1: Glucarate dehydratase

Chain D:





4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α, β, γ	110.84Å 124.85Å 139.09Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	28.48 – 1.70 28.48 – 1.70	Depositor EDS
% Data completeness (in resolution range)	99.4 (28.48-1.70) 99.5 (28.48-1.70)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	0.11	Depositor
$< I/\sigma(I) >$ [1]	2.62 (at 1.70Å)	Xtriage
Refinement program	PHENIX 1.7.3_928	Depositor
R, R_{free}	0.152 , 0.179 0.151 , 0.179	Depositor DCC
R_{free} test set	10582 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å^2)	17.7	Xtriage
Anisotropy	0.760	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.36 , 50.9	EDS
L-test for twinning[2]	$< \|L\| > = 0.50$, $< L^2 > = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o,F_c correlation	0.97	EDS
Total number of atoms	15933	wwPDB-VP
Average B, all atoms (Å^2)	22.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.85% of the height of the origin peak. No significant pseudotranslation is detected.*

[1]Intensities estimated from amplitudes.

[2]Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: NA, DTT, SO4, DTU, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\|>5$	RMSZ	$\#\|Z\|>5$
1	A	0.36	0/3523	0.52	0/4771
1	B	0.34	0/3549	0.51	0/4805
1	C	0.34	0/3533	0.51	0/4784
1	D	0.35	0/3557	0.52	0/4816
All	All	0.35	0/14162	0.51	0/19176

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3448	0	3379	7	0
1	B	3474	0	3398	15	0
1	C	3458	0	3387	10	0
1	D	3482	0	3406	16	0
2	A	5	0	0	0	0
2	B	5	0	0	0	0
2	C	5	0	0	0	0
2	D	5	0	0	0	0
3	A	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	B	1	0	0	0	0
3	C	1	0	0	0	0
3	D	1	0	0	0	0
4	A	12	0	16	0	0
4	B	24	0	32	3	1
4	C	6	0	8	1	0
5	A	8	0	9	0	0
6	B	8	0	10	2	0
6	C	8	0	9	0	0
6	D	8	0	10	5	0
7	A	525	0	0	0	0
7	B	503	0	0	1	0
7	C	459	0	0	2	1
7	D	486	0	0	1	0
All	All	15933	0	13664	44	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (44) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:168:CYS:SG	6:D:503:DTT:S4	2.39	1.15
1:B:168:CYS:SG	6:B:507:DTT:S1	2.44	1.05
1:D:168:CYS:HG	6:D:503:DTT:C4	1.98	0.76
1:D:168:CYS:SG	6:D:503:DTT:C4	2.75	0.74
1:D:168:CYS:HG	6:D:503:DTT:HS2	0.72	0.70
1:C:327[B]:GLU:OE2	7:C:788:HOH:O	2.10	0.68
1:C:320:ARG:NH2	1:D:327[A]:GLU:OE2	2.33	0.61
1:D:168:CYS:SG	6:D:503:DTT:H41	2.41	0.60
1:C:320:ARG:HH22	1:D:327[A]:GLU:CD	2.04	0.60
1:D:149:PHE:HB3	1:D:179:MET:HE1	1.82	0.60
1:C:327[A]:GLU:OE2	1:D:320:ARG:NH2	2.35	0.59
1:C:327[A]:GLU:CD	1:D:320:ARG:HH22	2.06	0.59
1:B:182:GLU:O	1:B:186:ARG:HG2	2.05	0.56
1:B:168:CYS:CB	6:B:507:DTT:HS1	2.17	0.53
1:C:281:PRO:HA	4:C:503:GOL:H12	1.90	0.53
1:D:89:ASP:OD1	1:D:89:ASP:N	2.42	0.53
1:D:249:LYS:O	7:D:1078:HOH:O	2.18	0.52
1:C:249:LYS:O	7:C:764:HOH:O	2.19	0.51
1:A:134:ASP:OD1	4:B:503:GOL:H2	2.11	0.50

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:201:LYS:NZ	1:D:362:ASP:OD2	2.41	0.50
1:A:376:LYS:NZ	1:A:397[A]:GLU:OE2	2.42	0.49
1:A:89:ASP:OD1	1:A:89:ASP:N	2.46	0.49
1:B:6:ILE:HD13	1:B:39:LEU:HD13	1.96	0.48
1:B:390:LYS:NZ	7:B:1089:HOH:O	2.47	0.48
1:B:216:VAL:HG11	1:B:230:LEU:HB2	1.96	0.47
1:B:341:ASP:OD1	1:B:341:ASP:N	2.43	0.46
1:B:8:GLU:HB2	1:B:40:LYS:HB2	1.97	0.46
1:D:106:ILE:HG12	1:D:312:PHE:O	2.16	0.46
1:B:89:ASP:OD1	1:B:89:ASP:N	2.49	0.45
1:B:91:ASP:CG	1:B:104:THR:H	2.18	0.45
1:C:229:THR:HA	1:C:252:LEU:HD22	1.97	0.45
1:B:91:ASP:OD1	1:B:104:THR:N	2.41	0.44
1:A:376:LYS:HD2	1:A:397[A]:GLU:CD	2.39	0.43
1:B:30:PRO:HG3	1:B:158:LEU:HB3	2.00	0.42
1:B:334:SER:O	1:B:362:ASP:HB2	2.20	0.42
1:C:376:LYS:NZ	1:C:397:GLU:OE2	2.35	0.42
1:B:304:ASP:OD1	4:B:505:GOL:H31	2.20	0.42
1:A:341:ASP:OD1	1:A:341:ASP:N	2.49	0.42
1:A:221:LYS:HB2	1:A:221:LYS:HE3	1.85	0.42
1:B:281:PRO:HA	4:B:505:GOL:H12	2.02	0.42
1:C:341:ASP:OD1	1:C:341:ASP:N	2.49	0.41
1:A:200:PHE:O	1:A:228:ILE:HA	2.21	0.41
1:D:106:ILE:HD11	1:D:314:THR:HG23	2.03	0.40
1:D:181:PRO:HB3	1:D:215:ALA:HA	2.03	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:B:506:GOL:O1	7:C:760:HOH:O[2_555]	2.12	0.08

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	442/464 (95%)	434 (98%)	8 (2%)	0	100	100
1	B	445/464 (96%)	438 (98%)	7 (2%)	0	100	100
1	C	443/464 (96%)	433 (98%)	10 (2%)	0	100	100
1	D	446/464 (96%)	439 (98%)	7 (2%)	0	100	100
All	All	1776/1856 (96%)	1744 (98%)	32 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	362/380 (95%)	360 (99%)	2 (1%)	86	80
1	B	365/380 (96%)	363 (100%)	2 (0%)	88	83
1	C	363/380 (96%)	360 (99%)	3 (1%)	81	74
1	D	366/380 (96%)	362 (99%)	4 (1%)	73	63
All	All	1456/1520 (96%)	1445 (99%)	11 (1%)	81	74

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	146	TYR
1	A	401	ASP
1	B	23	ASN
1	B	146	TYR
1	C	23	ASN
1	C	90	HIS
1	C	146	TYR
1	D	23	ASN
1	D	89	ASP
1	D	146	TYR

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	D	186	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	380	GLN
1	B	90	HIS
1	C	247	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

Of 19 ligands modelled in this entry, 4 are monoatomic - leaving 15 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
4	GOL	C	503	-	5,5,5	0.34	0	5,5,5	0.40	0
6	DTT	D	503	-	7,7,7	0.55	0	4,8,8	0.43	0
4	GOL	A	503	-	5,5,5	0.32	0	5,5,5	0.48	0
2	SO4	A	501	-	4,4,4	0.26	0	6,6,6	0.49	0
2	SO4	D	501	-	4,4,4	0.20	0	6,6,6	0.42	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
4	GOL	A	504	-	5,5,5	0.36	0	5,5,5	0.40	0
2	SO4	C	501	-	4,4,4	0.19	0	6,6,6	0.46	0
6	DTT	C	504	-	7,7,7	0.32	0	4,8,8	0.49	0
4	GOL	B	504	-	5,5,5	0.34	0	5,5,5	0.47	0
4	GOL	B	506	-	5,5,5	0.34	0	5,5,5	0.44	0
5	DTU	A	505	-	7,7,7	0.55	0	4,8,8	0.77	0
4	GOL	B	505	-	5,5,5	0.36	0	5,5,5	0.48	0
2	SO4	B	501	-	4,4,4	0.09	0	6,6,6	0.50	0
6	DTT	B	507	-	7,7,7	0.52	0	4,8,8	0.51	0
4	GOL	B	503	-	5,5,5	0.32	0	5,5,5	0.61	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GOL	C	503	-	-	2/4/4/4	-
6	DTT	D	503	-	-	4/8/8/8	-
4	GOL	A	503	-	-	0/4/4/4	-
4	GOL	A	504	-	-	2/4/4/4	-
6	DTT	C	504	-	-	2/8/8/8	-
4	GOL	B	504	-	-	1/4/4/4	-
4	GOL	B	506	-	-	2/4/4/4	-
5	DTU	A	505	-	-	8/8/8/8	-
4	GOL	B	505	-	-	2/4/4/4	-
6	DTT	B	507	-	-	8/8/8/8	-
4	GOL	B	503	-	-	3/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (34) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	504	GOL	O1-C1-C2-C3
4	B	503	GOL	C1-C2-C3-O3
4	B	505	GOL	C1-C2-C3-O3
4	B	506	GOL	O1-C1-C2-C3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
4	C	503	GOL	C1-C2-C3-O3
5	A	505	DTU	S1-C1-C2-O2
5	A	505	DTU	S1-C1-C2-C3
5	A	505	DTU	C1-C2-C3-O3
5	A	505	DTU	C1-C2-C3-C4
5	A	505	DTU	O2-C2-C3-O3
5	A	505	DTU	O2-C2-C3-C4
5	A	505	DTU	C2-C3-C4-S4
5	A	505	DTU	O3-C3-C4-S4
6	B	507	DTT	C1-C2-C3-O3
6	B	507	DTT	C1-C2-C3-C4
6	B	507	DTT	O2-C2-C3-O3
6	B	507	DTT	O2-C2-C3-C4
6	B	507	DTT	C2-C3-C4-S4
6	B	507	DTT	O3-C3-C4-S4
6	C	504	DTT	C2-C3-C4-S4
6	C	504	DTT	O3-C3-C4-S4
6	D	503	DTT	C1-C2-C3-O3
6	D	503	DTT	C1-C2-C3-C4
6	D	503	DTT	O2-C2-C3-O3
6	D	503	DTT	O2-C2-C3-C4
4	A	504	GOL	O1-C1-C2-O2
4	C	503	GOL	O2-C2-C3-O3
4	B	505	GOL	O2-C2-C3-O3
4	B	506	GOL	O1-C1-C2-O2
4	B	503	GOL	O2-C2-C3-O3
4	B	503	GOL	O1-C1-C2-O2
6	B	507	DTT	S1-C1-C2-O2
6	B	507	DTT	S1-C1-C2-C3
4	B	504	GOL	O1-C1-C2-C3

There are no ring outliers.

6 monomers are involved in 12 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	C	503	GOL	1	0
6	D	503	DTT	5	0
4	B	506	GOL	0	1
4	B	505	GOL	2	0
6	B	507	DTT	2	0
4	B	503	GOL	1	0

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å^2)	Q<0.9
1	A	441/464 (95%)	-0.45	4 (0%)	84	87	11, 17, 32, 56	0
1	B	441/464 (95%)	-0.41	4 (0%)	84	87	12, 19, 35, 60	0
1	C	441/464 (95%)	-0.25	7 (1%)	72	76	11, 21, 37, 80	0
1	D	441/464 (95%)	-0.27	8 (1%)	68	72	11, 19, 33, 78	0
All	All	1764/1856 (95%)	-0.34	23 (1%)	77	81	11, 19, 35, 80	0

All (23) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	92	ALA	5.9
1	D	92	ALA	5.4
1	C	89	ASP	4.8
1	D	88	ASN	4.6
1	C	88	ASN	4.5
1	D	89	ASP	4.5
1	C	93	GLY	4.3
1	A	92	ALA	4.2
1	D	90	HIS	4.2
1	D	93	GLY	4.0
1	C	2	SER	3.7
1	A	89	ASP	3.7
1	C	90	HIS	3.3
1	B	92	ALA	2.9
1	D	94	GLY	2.8
1	B	89	ASP	2.6
1	A	165	ASN	2.6
1	B	2	SER	2.3
1	A	93	GLY	2.3
1	D	2	SER	2.2
1	D	91	ASP	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	167	PRO	2.0
1	B	165	ASN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
6	DTT	B	507	8/8	0.68	0.31	56,74,75,115	0
4	GOL	A	504	6/6	0.79	0.27	30,40,46,49	0
4	GOL	C	503	6/6	0.83	0.20	27,34,38,40	0
5	DTU	A	505	8/8	0.84	0.28	43,76,82,115	0
4	GOL	B	506	6/6	0.84	0.37	38,48,53,55	0
6	DTT	C	504	8/8	0.87	0.29	46,59,71,124	0
4	GOL	B	505	6/6	0.89	0.16	30,34,36,45	0
4	GOL	B	504	6/6	0.89	0.15	18,30,38,39	0
6	DTT	D	503	8/8	0.90	0.29	43,72,77,116	0
3	NA	C	502	1/1	0.91	0.09	33,33,33,33	0
4	GOL	A	503	6/6	0.92	0.10	21,27,34,34	0
4	GOL	B	503	6/6	0.92	0.12	21,25,35,36	0
3	NA	A	502	1/1	0.94	0.10	22,22,22,22	0
3	NA	D	502	1/1	0.95	0.07	40,40,40,40	0
3	NA	B	502	1/1	0.95	0.08	26,26,26,26	0
2	SO4	C	501	5/5	0.95	0.09	19,22,29,30	0
2	SO4	B	501	5/5	0.98	0.08	18,21,22,23	0
2	SO4	A	501	5/5	0.98	0.07	15,16,21,22	0
2	SO4	D	501	5/5	0.98	0.07	18,18,29,32	0

6.5 Other polymers (i)

There are no such residues in this entry.