



Full wwPDB X-ray Structure Validation Report ⓘ

Jul 3, 2023 – 05:03 PM EDT

PDB ID : 7G8P
Title : ARHGEF2 PanDDA analysis group deposition – ARHGEF2 and RhoA in complex with Z1269184613
Authors : Bradshaw, W.J.; Katis, V.L.; Bountra, C.; von Delft, F.; Brennan, P.E.
Deposited on : 2023-06-22
Resolution : 2.21 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.34
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.34


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION



The reported resolution of this entry is 2.21 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4898 (2.20-2.20)
Clashscore	141614	5594 (2.20-2.20)
Ramachandran outliers	138981	5503 (2.20-2.20)
Sidechain outliers	138945	5504 (2.20-2.20)
RSRZ outliers	127900	4800 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	185	
2	B	245	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	FMT	A	205	-	-	X	-
5	FMT	B	504	-	-	X	-

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 3714 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Transforming protein RhoA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	179	1439	914	241	274	10	0	4	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	SER	-	expression tag	UNP P61586


- Molecule 2 is a protein called Rho guanine nucleotide exchange factor 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	244	2062	1313	361	376	12	0	15	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	204	SER	-	expression tag	UNP Q92974
B	205	MET	-	expression tag	UNP Q92974

- Molecule 3 is 1,3-benzothiazole-6-sulfonamide (three-letter code: Z2F) (formula: C₇H₆N₂O₂S₂) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	S		
3	A	1	13	7	2	2	2	0	0

- Molecule 4 is DIMETHYL SULFOXIDE (three-letter code: DMS) (formula: C₂H₆OS).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	
			Total	C	O			S
4	A	1	4	2	1	1	0	0
4	B	1	4	2	1	1	0	0
4	B	1	4	2	1	1	0	0

- Molecule 5 is FORMIC ACID (three-letter code: FMT) (formula: CH₂O₂).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total C O 3 1 2	0	0
5	A	1	Total C O 3 1 2	0	0
5	A	1	Total C O 3 1 2	0	0
5	A	1	Total C O 3 1 2	0	0
5	B	1	Total C O 3 1 2	0	0
5	B	1	Total C O 3 1 2	0	0
5	B	1	Total C O 3 1 2	0	0


- Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	65	Total O 65 65	0	0
6	B	101	Total O 102 102	0	2


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Transforming protein RhoA



- Molecule 2: Rho guanine nucleotide exchange factor 2



4 Data and refinement statistics

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, α , β , γ	70.57Å 70.57Å 197.00Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	33.22 – 2.21 33.22 – 2.21	Depositor EDS
% Data completeness (in resolution range)	89.2 (33.22-2.21) 89.3 (33.22-2.21)	Depositor EDS
R_{merge}	0.43	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.09 (at 2.22Å)	Xtrriage
Refinement program	BUSTER 2.10.4 (17-FEB-2023)	Depositor
R, R_{free}	0.259 , 0.324 0.247 , 0.322	Depositor DCC
R_{free} test set	1172 reflections (5.08%)	wwPDB-VP
Wilson B-factor (Å ²)	45.6	Xtrriage
Anisotropy	0.552	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 77.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	3714	wwPDB-VP
Average B, all atoms (Å ²)	77.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.38% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: DMS, Z2F, FMT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.43	0/1479	0.60	0/2001
2	B	0.38	0/2149	0.54	0/2888
All	All	0.40	0/3628	0.57	0/4889

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1439	0	1444	11	0
2	B	2062	0	2127	30	0
3	A	13	0	0	0	0
4	A	4	0	6	0	0
4	B	8	0	12	0	0
5	A	12	0	7	3	0
5	B	9	0	3	4	0
6	A	65	0	0	0	0
6	B	102	0	0	0	0
All	All	3714	0	3599	41	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 6.

All (41) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:231:LYS:HG2	2:B:235:LYS:HZ3	1.56	0.71
2:B:340:GLU:HG3	2:B:344:ARG:HE	1.57	0.69
2:B:265:MET:HG2	2:B:269:LEU:HD12	1.81	0.63
2:B:231:LYS:HD3	2:B:235:LYS:NZ	2.14	0.62
2:B:437:GLU:HB3	2:B:440:ALA:HB2	1.81	0.62
2:B:231:LYS:HG2	2:B:235:LYS:NZ	2.15	0.62
1:A:145:ARG:HE	5:A:203:FMT:H	1.69	0.58
5:A:205:FMT:C	2:B:376:LYS:HD3	2.35	0.57
2:B:231:LYS:CG	2:B:235:LYS:HZ3	2.17	0.56
2:B:323:ILE:HD11	2:B:417:THR:HB	1.88	0.56
2:B:262:ARG:HG2	2:B:276:VAL:HG12	1.89	0.55
1:A:16:CYS:HA	1:A:85:SER:HB2	1.90	0.53
1:A:118:LYS:HB3	1:A:121:LEU:HD12	1.90	0.53
2:B:319:GLY:HA3	2:B:414:ASP:HB3	1.90	0.53
2:B:296:GLN:HB3	2:B:321:LEU:HD11	1.91	0.52
2:B:226:LEU:HD11	2:B:234:MET:HG3	1.92	0.52
2:B:345:HIS:CD2	2:B:349:LEU:HD11	2.44	0.51
2:B:423:LYS:NZ	5:B:503:FMT:H	2.26	0.51
1:A:43:VAL:O	5:A:205:FMT:H	2.10	0.50
2:B:231:LYS:CD	2:B:235:LYS:HZ3	2.24	0.49
2:B:231:LYS:CG	2:B:235:LYS:NZ	2.73	0.49
2:B:258:THR:HG23	2:B:283[B]:VAL:HG22	1.93	0.49
2:B:231:LYS:HD3	2:B:235:LYS:HZ1	1.77	0.49
2:B:418:ALA:O	2:B:422:VAL:HG23	2.13	0.48
2:B:345:HIS:NE2	2:B:349:LEU:HD11	2.31	0.46
2:B:336:LYS:NZ	5:B:504:FMT:C	2.78	0.46
2:B:353:LYS:NZ	2:B:381:GLN:HE22	2.13	0.46
2:B:231:LYS:CD	2:B:235:LYS:NZ	2.80	0.45
2:B:336:LYS:HZ3	5:B:504:FMT:C	2.29	0.45
1:A:10[B]:ILE:HG12	1:A:81:LEU:HB2	1.98	0.45
1:A:119:LYS:HD3	1:A:158:GLU:HB3	1.99	0.44
2:B:231:LYS:HD3	2:B:235:LYS:HZ3	1.80	0.43
2:B:336:LYS:CE	5:B:504:FMT:C	2.96	0.43
1:A:108:PRO:O	1:A:109:ASN:HB2	2.18	0.43
1:A:4:ILE:HG23	1:A:53:VAL:HG22	2.01	0.43
2:B:353:LYS:HZ3	2:B:381:GLN:HE22	1.69	0.41
1:A:85:SER:HB3	1:A:88:SER:HB3	2.02	0.41
2:B:372:PRO:HD2	2:B:375:LEU:HD23	2.03	0.40

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:71:PRO:HA	1:A:74:TYR:CD2	2.56	0.40
2:B:345:HIS:HE2	2:B:349:LEU:HD11	1.86	0.40
1:A:31:PRO:HB2	1:A:35:VAL:HG22	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	181/185 (98%)	174 (96%)	7 (4%)	0	100	100
2	B	257/245 (105%)	250 (97%)	7 (3%)	0	100	100
All	All	438/430 (102%)	424 (97%)	14 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	160/160 (100%)	155 (97%)	5 (3%)	40	51
2	B	239/224 (107%)	233 (98%)	6 (2%)	47	60
All	All	399/384 (104%)	388 (97%)	11 (3%)	42	56

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	4	ILE
1	A	28	ASP
1	A	29	GLN
1	A	85	SER
1	A	150	ARG
2	B	271	LEU
2	B	340	GLU
2	B	375	LEU
2	B	410	GLU
2	B	438	LYS
2	B	446	TYR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	63	GLN
1	A	123	ASN
1	A	149	ASN
2	B	381	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

11 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond

length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	DMS	B	501	-	3,3,3	0.66	0	3,3,3	0.28	0
5	FMT	A	205	-	2,2,2	1.10	0	1,1,1	0.15	0
3	Z2F	A	201	-	12,14,14	0.69	0	15,21,21	1.08	1 (6%)
5	FMT	A	206	-	2,2,2	1.25	0	1,1,1	0.14	0
4	DMS	A	202	-	3,3,3	0.65	0	3,3,3	0.38	0
5	FMT	B	503	-	2,2,2	1.11	0	1,1,1	0.18	0
5	FMT	A	204	-	2,2,2	1.27	0	1,1,1	0.13	0
4	DMS	B	502	-	3,3,3	0.68	0	3,3,3	0.29	0
5	FMT	B	505	-	2,2,2	0.95	0	1,1,1	0.26	0
5	FMT	B	504	-	2,2,2	0.80	0	1,1,1	0.29	0
5	FMT	A	203	-	2,2,2	1.32	0	1,1,1	0.10	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	Z2F	A	201	-	-	0/6/6/6	0/2/2/2

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	201	Z2F	C5-C3-N1	3.13	114.83	107.87

There are no chirality outliers.


There are no torsion outliers.

There are no ring outliers.

4 monomers are involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	205	FMT	2	0
5	B	503	FMT	1	0
5	B	504	FMT	3	0
5	A	203	FMT	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	179/185 (96%)	0.79	28 (15%) 2 1	48, 71, 128, 140	0
2	B	244/245 (99%)	0.64	27 (11%) 5 4	50, 78, 110, 119	0
All	All	423/430 (98%)	0.70	55 (13%) 3 3	48, 74, 113, 140	0

All (55) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	134	MET	8.3
2	B	269	LEU	5.8
1	A	135	LYS	5.4
1	A	30	PHE	4.5
2	B	270	HIS	4.4
2	B	226	LEU	4.3
2	B	298	LEU	4.1
1	A	157	MET	4.0
2	B	273	PRO	3.7
1	A	127	THR	3.7
1	A	181	ALA	3.7
2	B	406	HIS	3.7
2	B	386	LEU	3.7
1	A	31	PRO	3.7
1	A	33	VAL	3.6
1	A	124	ASP	3.6
1	A	133	LYS	3.4
1	A	123	ASN	3.4
1	A	126	HIS	3.3
2	B	357	ALA	3.3
1	A	129	ARG	3.3
2	B	329	PRO	3.3
2	B	384	ILE	3.3
1	A	29	GLN	3.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	130	GLU	3.0
2	B	407	GLY	3.0
2	B	408[A]	ILE	2.9
2	B	385	LEU	2.8
1	A	10[A]	ILE	2.8
2	B	403	GLN	2.7
2	B	444	GLU	2.7
2	B	388	THR	2.6
1	A	156	TYR	2.5
2	B	297	LEU	2.4
1	A	28	ASP	2.4
2	B	254	LEU	2.3
1	A	125	GLU	2.3
2	B	412	ARG	2.3
1	A	170	VAL	2.3
1	A	8	LEU	2.2
2	B	434	TYR	2.2
2	B	430	ASP	2.2
2	B	360	LYS	2.1
1	A	11	VAL	2.1
1	A	128	ARG	2.1
2	B	359	ASP	2.1
2	B	355	LEU	2.1
1	A	9	VAL	2.1
2	B	389	GLN	2.1
1	A	115	VAL	2.1
1	A	108	PRO	2.1
2	B	275	VAL	2.0
1	A	34	TYR	2.0
1	A	113	ILE	2.0
2	B	383	CYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no monosaccharides in this entry.

6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	FMT	A	204	3/3	0.61	0.31	89,89,89,89	0
5	FMT	A	205	3/3	0.80	0.38	85,85,85,85	0
4	DMS	B	502	4/4	0.83	0.19	86,86,86,86	4
5	FMT	A	203	3/3	0.85	0.21	74,74,75,75	0
5	FMT	B	503	3/3	0.90	0.25	80,80,80,80	0
5	FMT	A	206	3/3	0.92	0.14	77,77,78,78	0
5	FMT	B	505	3/3	0.92	0.19	82,82,83,83	0
3	Z2F	A	201	13/13	0.93	0.10	61,62,64,64	13
5	FMT	B	504	3/3	0.94	0.11	67,67,68,68	0
4	DMS	B	501	4/4	0.96	0.13	84,84,84,84	4
4	DMS	A	202	4/4	0.98	0.32	69,69,69,69	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.