

Full wwPDB X-ray Structure Validation Report (i)

Sep 6, 2023 – 02:21 PM EDT

PDB ID : 4GE0

> Title : Schizosaccharomyces pombe DJ-1 T114P mutant

Authors Madzelan, P.; Labunska, T.; Wilson, M.A.

2012-08-01 Deposited on

1.45 Å(reported) Resolution

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

> The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity 4.02b-467

> 1.8.5 (274361), CSD as541be (2020) Mogul

Xtriage (Phenix) 1.13

EDS 2.35

20191225.v01 (using entries in the PDB archive December 25th 2019) Percentile statistics

> Refmac 5.8.0158

CCP4 7.0.044 (Gargrove)

Ideal geometry (proteins) Engh & Huber (2001) Ideal geometry (DNA, RNA) Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) 2.35

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 1.45 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive	Similar resolution
Metric	$(\# ext{Entries})$	$(\# ext{Entries}, ext{ resolution range}(ext{Å}))$
R_{free}	130704	1156 (1.46-1.46)
Clashscore	141614	1202 (1.46-1.46)
Ramachandran outliers	138981	1178 (1.46-1.46)
Sidechain outliers	138945	1178 (1.46-1.46)
RSRZ outliers	127900	1139 (1.46-1.46)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain	
1	A	194	90%	8% ••
1	В	194	87%	10%
1	С	194	2%	10% •
1	D	194	89%	9% •

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard



residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	EDO	В	201	-	-	X	-



2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 7200 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Uncharacterized protein C22E12.03c.


Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace		
1	Λ	190	Total	С	N	О	S	0	9	0
1	A	190	1557	1001	256	291	9	0	9	
1	В	190	Total	С	N	О	S	0	10	0
1			1561	1003	255	295	8	U		
1	С	190	Total	С	N	О	S	0	15	0
1		190	1607	1030	263	305	9	0	19	
1	1 D	190	Total	С	N	О	S	0	7	0
1			1537	987	251	292	7	U	1	U

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	GLY	-	expression tag	UNP Q10356
A	-1	SER	-	expression tag	UNP Q10356
A	0	HIS	-	expression tag	UNP Q10356
A	114	PRO	THR	engineered mutation	UNP Q10356
В	-2	GLY	-	expression tag	UNP Q10356
В	-1	SER	-	expression tag	UNP Q10356
В	0	HIS	-	expression tag	UNP Q10356
В	114	PRO	THR	engineered mutation	UNP Q10356
С	-2	GLY	-	expression tag	UNP Q10356
С	-1	SER	-	expression tag	UNP Q10356
С	0	HIS	-	expression tag	UNP Q10356
С	114	PRO	THR	engineered mutation	UNP Q10356
D	-2	GLY	-	expression tag	UNP Q10356
D	-1	SER	-	expression tag	UNP Q10356
D	0	HIS	-	expression tag	UNP Q10356
D	114	PRO	THR	engineered mutation	UNP Q10356

• Molecule 2 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: $C_2H_6O_2$).





Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	В	1	Total C O 4 2 2	0	0
2	С	1	Total C O 4 2 2	0	0

• Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	С	2	Total Mg 2 2	0	0

• Molecule 4 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	237	Total O 237 237	0	0
4	В	223	Total O 223 223	0	0
4	C	292	Total O 292 292	0	0
4	D	176	Total O 176 176	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Uncharacterized protein C22E12.03c





4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1	Depositor
Cell constants	44.49\AA 51.74Å 82.75Å	Donositon
a, b, c, α , β , γ	89.10° 89.08° 66.86°	Depositor
Resolution (Å)	36.51 - 1.45	Depositor
Resolution (A)	35.50 - 1.45	EDS
% Data completeness	94.3 (36.51-1.45)	Depositor
(in resolution range)	94.3 (35.50-1.45)	EDS
R_{merge}	0.07	Depositor
$R_{sym} < I/\sigma(I) > 1$	(Not available)	Depositor
$< I/\sigma(I) > 1$	4.44 (at 1.45Å)	Xtriage
Refinement program	REFMAC 5.6.0116	Depositor
D D.	0.179 , 0.208	Depositor
R, R_{free}	0.173 , 0.202	DCC
R_{free} test set	5669 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	17.0	Xtriage
Anisotropy	0.228	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.33 , 44.8	EDS
L-test for twinning ²	$< L > = 0.50, < L^2> = 0.33$	Xtriage
	0.010 for h,h-k,-l	
Estimated twinning fraction	0.004 for -h,-k,l	Xtriage
	0.003 for -h,-h+k,-l	
F_o, F_c correlation	0.97	EDS
Total number of atoms	7200	wwPDB-VP
Average B, all atoms (\mathring{A}^2)	19.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 6.28% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of <|L|>, $<L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: CSD, EDO, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond angles	
IVIOI		RMSZ	# Z > 5	RMSZ	# Z > 5
1	A	0.43	0/1583	0.57	0/2137
1	В	0.42	0/1588	0.56	0/2145
1	С	0.45	0/1634	0.59	0/2204
1	D	0.44	0/1564	0.61	0/2114
All	All	0.44	0/6369	0.58	0/8600

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a maintenain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	В	0	1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	В	102	LYS	Peptide

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.



Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1557	0	1575	20	0
1	В	1561	0	1570	25	1
1	С	1607	0	1610	24	0
1	D	1537	0	1541	14	0
2	В	4	0	6	8	0
2	С	4	0	6	1	0
3	С	2	0	0	0	0
4	A	237	0	0	10	2
4	В	223	0	0	9	1
4	С	292	0	0	16	2
4	D	176	0	0	8	2
All	All	7200	0	6308	84	4

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (84) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:170[B]:LYS:NZ	4:C:579:HOH:O	1.64	1.29
1:A:106[A]:TRP:CH2	4:A:421:HOH:O	1.67	1.27
1:C:106[A]:TRP:HH2	4:C:526:HOH:O	1.21	1.21
1:C:106[A]:TRP:CH2	4:C:526:HOH:O	1.94	1.06
1:A:102:LYS:HE3	4:A:400:HOH:O	1.61	0.99
1:A:106[A]:TRP:HH2	4:A:421:HOH:O	1.10	0.99
1:B:46[B]:MET:SD	4:B:461:HOH:O	2.22	0.96
1:D:104[A]:ASN:HB2	4:D:363:HOH:O	1.65	0.94
1:B:106[A]:TRP:HZ3	4:B:508:HOH:O	1.51	0.94
1:A:46[B]:MET:SD	1:A:52[B]:MET:HE3	2.10	0.90
1:D:147:GLN:CB	4:D:343:HOH:O	2.22	0.86
1:B:45[B]:LYS:HZ3	2:B:201:EDO:H12	1.39	0.86
1:B:45[A]:LYS:HE3	4:B:364:HOH:O	1.76	0.84
1:A:46[B]:MET:SD	1:A:52[B]:MET:CE	2.65	0.84
1:C:66[B]:ASP:OD1	4:C:397:HOH:O	1.96	0.84
1:D:103[A]:PRO:O	4:D:363:HOH:O	1.93	0.84
1:A:106[A]:TRP:CZ3	4:A:421:HOH:O	2.03	0.83
1:D:147:GLN:HB3	4:D:343:HOH:O	1.78	0.83
1:B:124:ASN:HB2	1:B:152:GLU:HG2	1.61	0.81
1:B:45[B]:LYS:NZ	2:B:201:EDO:H12	1.97	0.80
1:C:10[A]:ASP:OD2	4:C:527:HOH:O	2.00	0.80
1:C:93:GLN:O	1:C:97[B]:LYS:HG3	1.87	0.74
1:C:18[B]:SER:OG	4:C:508:HOH:O	2.05	0.73



 $Continued\ from\ previous\ page...$

Atom-1	Atom-2	Interatomic	Clash
Atom-1	Atom-2	${ m distance} \; ({ m \AA})$	overlap(Å)
1:D:147:GLN:HB2	4:D:343:HOH:O	1.88	0.70
1:A:3[A]:LYS:HD2	4:A:365:HOH:O	1.91	0.68
1:B:45[B]:LYS:CE	2:B:201:EDO:H12	2.25	0.67
1:C:39:ASN:HB2	4:C:468:HOH:O	1.95	0.66
1:C:97[B]:LYS:HD3	1:C:120:SER:O	1.94	0.66
1:B:104[A]:ASN:O	1:B:106[A]:TRP:CZ3	2.49	0.66
1:D:38[B]:GLU:HG2	4:D:367:HOH:O	1.96	0.65
1:D:51:GLU:OE1	4:D:375:HOH:O	2.15	0.65
1:B:80:GLY:O	1:B:84:LYS:HG2	1.98	0.63
1:C:37:GLY:O	4:C:417:HOH:O	2.16	0.61
1:B:124:ASN:CB	1:B:152:GLU:HG2	2.32	0.60
1:A:106[A]:TRP:HZ3	4:A:270:HOH:O	1.87	0.56
1:C:106[A]:TRP:CZ3	4:C:526:HOH:O	2.43	0.56
1:C:3:LYS:HD3	4:C:360:HOH:O	2.05	0.56
1:B:106[B]:TRP:CZ3	1:B:154:ASN:HA	2.41	0.55
2:C:201:EDO:O1	1:D:18[A]:SER:OG	2.24	0.54
1:A:84:LYS:NZ	4:A:295:HOH:O	2.40	0.54
1:A:106[A]:TRP:CZ3	1:A:175:VAL:HG12	2.44	0.53
1:C:41:ASP:HB2	4:C:376:HOH:O	2.08	0.53
1:C:97[B]:LYS:HG2	1:C:120:SER:HB3	1.92	0.52
1:B:81[B]:LEU:HD12	4:B:336:HOH:O	2.09	0.52
1:B:106[A]:TRP:CZ3	4:B:508:HOH:O	2.36	0.52
1:B:45[B]:LYS:HE2	2:B:201:EDO:C1	2.40	0.51
1:B:103[A]:PRO:C	1:B:104[A]:ASN:CG	2.69	0.51
1:A:18[B]:SER:OG	4:A:364:HOH:O	2.18	0.50
1:C:97[A]:LYS:HD2	4:C:401:HOH:O	2.11	0.49
1:D:106[A]:TRP:HZ3	4:D:353:HOH:O	1.96	0.49
1:B:179[B]:ASP:CG	4:B:397:HOH:O	2.51	0.48
1:D:3:LYS:HG3	1:D:30:PRO:HB2	1.95	0.48
1:C:52[B]:MET:HG3	1:C:52[B]:MET:O	2.14	0.48
1:A:3[A]:LYS:CD	4:A:365:HOH:O	2.57	0.47
2:B:201:EDO:H21	4:B:336:HOH:O	2.14	0.47
1:A:46[B]:MET:SD	1:A:52[B]:MET:HE1	2.50	0.47
1:B:45[B]:LYS:CE	2:B:201:EDO:C1	2.92	0.47
1:D:76:ILE:HD13	1:D:116:THR:HG21	1.96	0.47
1:B:45[B]:LYS:HE2	2:B:201:EDO:H11	1.96	0.46
1:A:156[A]:ILE:HD12	1:A:156[A]:ILE:N	2.31	0.46
1:C:97[A]:LYS:CD	4:C:401:HOH:O	2.62	0.46
1:A:156[B]:ILE:HD13	1:A:170[B]:LYS:HG3	1.97	0.46
1:A:43:LEU:HD12	1:A:52[B]:MET:O	2.17	0.45
1:D:156[A]:ILE:HD12	1:D:156[A]:ILE:N	2.32	0.44



Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:102:LYS:HA	1:B:103[A]:PRO:HD2	1.69	0.43
1:B:134:ARG:HD2	4:B:388:HOH:O	2.17	
			0.43
1:D:97:LYS:HE2	1:D:120:SER:O	2.19	0.43
1:C:97[B]:LYS:HE3	1:C:120:SER:HA	2.00	0.43
1:A:180:LYS:HE2	4:A:259:HOH:O	2.18	0.43
1:B:134:ARG:HG3	1:B:144:TYR:CG	2.54	0.43
1:B:179[B]:ASP:OD1	4:B:397:HOH:O	2.21	0.43
1:B:45[B]:LYS:HE2	2:B:201:EDO:H12	2.01	0.42
1:C:186:LYS:HE3	4:C:501:HOH:O	2.20	0.42
1:B:109:MET:O	1:B:157:THR:HA	2.20	0.42
1:A:98:GLU:O	1:A:102:LYS:HG3	2.20	0.42
1:C:66[B]:ASP:CG	4:C:397:HOH:O	2.52	0.41
1:A:100:TYR:CE2	1:A:101:LYS:HG3	2.54	0.41
1:C:180:LYS:HE3	4:C:478:HOH:O	2.20	0.41
1:A:27:ALA:O	1:A:180:LYS:HE3	2.20	0.41
1:D:109:MET:O	1:D:157:THR:HA	2.20	0.41
1:C:111:CSD:O	1:C:114:PRO:HD2	2.20	0.40
1:C:97[A]:LYS:HE2	1:C:120:SER:O	2.22	0.40
1:B:100:TYR:CE2	1:B:123:PRO:HD2	2.57	0.40
1:C:156[B]:ILE:HD12	1:C:171:LEU:HD23	2.04	0.40

All (4) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	$egin{aligned} & & & & & & & & & \\ & & & & & & & & & $	Clash overlap (Å)
		uistance (A)	overiap (A)
4:A:343:HOH:O	4:B:497:HOH:O[1_645]	1.98	0.22
4:C:330:HOH:O	4:D:342:HOH:O[1_655]	2.16	0.04
1:B:104[A]:ASN:ND2	4:D:343:HOH:O[1_556]	2.17	0.03
4:A:412:HOH:O	4:C:472:HOH:O[1_645]	2.19	0.01

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.



Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	A	$196/194\ (101\%)$	192 (98%)	4 (2%)	0	100	100
1	В	$197/194\ (102\%)$	194 (98%)	3 (2%)	0	100	100
1	C	$202/194\ (104\%)$	198 (98%)	4 (2%)	0	100	100
1	D	194/194 (100%)	190 (98%)	4 (2%)	0	100	100
All	All	789/776~(102%)	774 (98%)	15 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Perce	ntiles
1	A	168/162 (104%)	167 (99%)	1 (1%)	86	69
1	В	169/162 (104%)	167 (99%)	2 (1%)	71	43
1	С	174/162 (107%)	174 (100%)	0	100	100
1	D	166/162 (102%)	165 (99%)	1 (1%)	86	69
All	All	677/648 (104%)	673 (99%)	4 (1%)	86	69

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	180	LYS
1	В	40	LYS
1	В	178	LYS
1	D	139	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	126	GLN



5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Tuno	Chain	Res	Link	Bond lengths			Е	ond ang	gles
MIOI	Type	Chain	nes	Lilik	Counts	RMSZ	# Z >2	Counts	RMSZ	# Z > 2
1	CSD	A	111	1	3,7,8	1.26	0	1,8,10	0.96	0
1	CSD	В	111	1	3,7,8	1.12	0	1,8,10	1.41	0
1	CSD	D	111	1	3,7,8	1.11	0	1,8,10	0.06	0
1	CSD	С	111	1	3,7,8	1.14	0	1,8,10	1.56	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CSD	A	111	1	-	2/2/6/8	-
1	CSD	В	111	1	-	2/2/6/8	-
1	CSD	D	111	1	-	2/2/6/8	-
1	CSD	С	111	1	-	2/2/6/8	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (8) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	111	CSD	N-CA-CB-SG
1	A	111	CSD	CA-CB-SG-OD1



Continued from previous page...

Mol	Chain	Res	Type	Atoms
1	В	111	CSD	CA-CB-SG-OD1
1	С	111	CSD	N-CA-CB-SG
1	С	111	CSD	CA-CB-SG-OD1
1	D	111	CSD	N-CA-CB-SG
1	D	111	CSD	CA-CB-SG-OD1
1	В	111	CSD	N-CA-CB-SG

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	С	111	CSD	1	0

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

Of 4 ligands modelled in this entry, 2 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type Chain		$_{ m nain} \mid { m Res} \mid$	Link	\mathbf{B}_{0}	Bond lengths			Bond angles		
	Туре	Chain	rtes	LIIIK	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2	
2	EDO	С	201	_	3,3,3	0.57	0	2,2,2	0.08	0	
2	EDO	В	201	-	3,3,3	0.43	0	2,2,2	0.07	0	

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	EDO	С	201	-	-	0/1/1/1	-



Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	EDO	В	201	-	-	1/1/1/1	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	В	201	EDO	O1-C1-C2-O2

There are no ring outliers.

2 monomers are involved in 9 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	С	201	EDO	1	0
2	В	201	EDO	8	0

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ $>$	$\#\mathrm{RSRZ}{>}2$	$OWAB(A^2)$	Q < 0.9
1	A	189/194 (97%)	0.35	5 (2%) 56 58	10, 16, 32, 40	0
1	В	189/194 (97%)	0.61	19 (10%) 7 8	11, 20, 41, 56	0
1	С	189/194 (97%)	0.22	4 (2%) 63 65	9, 14, 23, 35	0
1	D	189/194 (97%)	0.26	4 (2%) 63 65	9, 15, 28, 44	0
All	All	756/776 (97%)	0.36	32 (4%) 36 39	9, 16, 33, 56	0

All (32) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	В	103[A]	PRO	5.2
1	В	139	GLU	4.7
1	D	103[A]	PRO	3.6
1	В	142	TYR	3.3
1	В	131	PRO	3.2
1	В	123	PRO	3.0
1	В	121	GLY	2.9
1	В	135	GLY	2.9
1	В	133	VAL	2.9
1	В	137	LEU	2.9
1	В	145	LEU	2.9
1	С	40	LYS	2.8
1	В	119	THR	2.7
1	A	123	PRO	2.6
1	A	140	GLY	2.5
1	В	126	GLN	2.5
1	A	40	LYS	2.4
1	С	121	GLY	2.4
1	В	136	GLN	2.4
1	A	2	VAL	2.4
1	В	144	TYR	2.4



Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	С	75	ILE	2.4
1	D	121	GLY	2.3
1	A	106[A]	TRP	2.2
1	В	50	VAL	2.2
1	С	178	LYS	2.2
1	В	179[A]	ASP	2.2
1	В	106[A]	TRP	2.1
1	В	125	LYS	2.1
1	В	124	ASN	2.1
1	D	50	VAL	2.1
1	D	104[A]	ASN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-}\mathbf{factors}(\mathbf{\mathring{A}}^2)$	Q < 0.9
1	CSD	В	111	8/9	0.91	0.13	17,18,21,21	0
1	CSD	A	111	8/9	0.93	0.12	14,16,18,19	0
1	CSD	D	111	8/9	0.95	0.09	13,15,17,17	0
1	CSD	С	111	8/9	0.96	0.08	12,12,15,16	0

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-}\mathbf{factors}(\mathbf{\mathring{A}}^2)$	Q<0.9
2	EDO	С	201	4/4	0.90	0.16	16,19,20,23	0
2	EDO	В	201	4/4	0.94	0.18	21,22,25,35	0
3	MG	С	202	1/1	0.96	0.23	32,32,32,32	0



Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-}\mathbf{factors}(\mathbf{\mathring{A}}^2)$	Q < 0.9
3	MG	С	203	1/1	0.96	0.33	28,28,28,28	0

6.5 Other polymers (i)

There are no such residues in this entry.

