



Full wwPDB X-ray Structure Validation Report i

Sep 7, 2023 – 01:07 AM EDT

PDB ID : 4GJ4
Title : The Crystal Structure of the soluble Guanylate Cyclase PAS alpha domain from *Manduca sexta*
Authors : Purohit, R.; Montfort, W.R.; Weichsel, A.
Deposited on : 2012-08-09
Resolution : 1.80 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>
with specific help available everywhere you see the i symbol.

The types of validation reports are described at
<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references](#) i) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.35
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5950 (1.80-1.80)
Clashscore	141614	6793 (1.80-1.80)
Ramachandran outliers	138981	6697 (1.80-1.80)
Sidechain outliers	138945	6696 (1.80-1.80)
RSRZ outliers	127900	5850 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.



2 Entry composition (i)

There are 3 unique types of molecules in this entry. The entry contains 3844 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Soluble guanylyl cyclase alpha-1 subunit.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
1	A	113	Total	C 908	N 595	O 158	S 151	4	0	3	0
1	B	112	Total	C 897	N 589	O 154	S 150	4	0	3	0
1	C	113	Total	C 894	N 588	O 153	S 149	4	0	2	0
1	D	117	Total	C 907	N 595	O 153	S 155	4	0	0	0

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	285	ALA	CYS	engineered mutation	UNP O77105
A	352	ALA	CYS	engineered mutation	UNP O77105
A	374	ALA	CYS	engineered mutation	UNP O77105
B	285	ALA	CYS	engineered mutation	UNP O77105
B	352	ALA	CYS	engineered mutation	UNP O77105
B	374	ALA	CYS	engineered mutation	UNP O77105
C	285	ALA	CYS	engineered mutation	UNP O77105
C	352	ALA	CYS	engineered mutation	UNP O77105
C	374	ALA	CYS	engineered mutation	UNP O77105
D	285	ALA	CYS	engineered mutation	UNP O77105
D	352	ALA	CYS	engineered mutation	UNP O77105
D	374	ALA	CYS	engineered mutation	UNP O77105

- Molecule 2 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total O S 10 8 2	0	1
2	B	1	Total O S 5 4 1	0	0
2	B	1	Total O S 5 4 1	0	0
2	B	1	Total O S 10 8 2	0	1
2	C	1	Total O S 5 4 1	0	0
2	C	1	Total O S 5 4 1	0	0
2	C	1	Total O S 10 8 2	0	1
2	C	1	Total O S 5 4 1	0	0
2	D	1	Total O S 5 4 1	0	0
2	D	1	Total O S 5 4 1	0	0
2	D	1	Total O S 5 4 1	0	0

- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	47	Total O 47 47	0	0

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	B	46	Total O 47 47	0	1
3	C	42	Total O 42 42	0	0
3	D	30	Total O 32 32	0	2


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Soluble guanylyl cyclase alpha-1 subunit




- Molecule 1: Soluble guanylyl cyclase alpha-1 subunit



- Molecule 1: Soluble guanylyl cyclase alpha-1 subunit



- Molecule 1: Soluble guanylyl cyclase alpha-1 subunit



4 Data and refinement statistics i

Property	Value	Source
Space group	H 3 2	Depositor
Cell constants a, b, c, α , β , γ	95.42Å 95.42Å 317.69Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	23.15 – 1.80 22.27 – 1.80	Depositor EDS
% Data completeness (in resolution range)	99.9 (23.15-1.80) 100.0 (22.27-1.80)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ ¹	2.66 (at 1.80Å)	Xtriage
Refinement program	REFMAC 5.6.0117	Depositor
R , R_{free}	0.196 , 0.242 0.208 , 0.250	Depositor DCC
R_{free} test set	2658 reflections (5.11%)	wwPDB-VP
Wilson B-factor (Å ²)	37.4	Xtriage
Anisotropy	0.075	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 58.9	EDS
L-test for twinning ²	$< L > = 0.52$, $< L^2 > = 0.36$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	3844	wwPDB-VP
Average B, all atoms (Å ²)	58.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.53% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [\(i\)](#)

5.1 Standard geometry [\(i\)](#)

Bond lengths and bond angles in the following residue types are not validated in this section:
SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.65	0/931	0.75	0/1254
1	B	0.68	0/921	0.83	2/1240 (0.2%)
1	C	0.65	0/920	0.80	1/1240 (0.1%)
1	D	0.61	0/930	0.75	1/1254 (0.1%)
All	All	0.65	0/3702	0.78	4/4988 (0.1%)

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	C	295	ASP	CB-CG-OD1	5.76	123.48	118.30
1	B	339	ARG	CG-CD-NE	-5.42	100.41	111.80
1	B	339	ARG	NE-CZ-NH2	-5.33	117.63	120.30
1	D	295	ASP	CB-CG-OD1	5.11	122.90	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	908	0	929	11	0
1	B	897	0	909	12	0
1	C	894	0	918	11	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	D	907	0	924	11	0
2	A	10	0	0	0	0
2	B	20	0	0	1	0
2	C	25	0	0	1	0
2	D	15	0	0	0	0
3	A	47	0	0	0	0
3	B	47	0	0	0	0
3	C	42	0	0	0	0
3	D	32	0	0	0	0
All	All	3844	0	3680	38	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (38) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:359:THR:HG21	1:B:279:ILE:HG22	1.56	0.88
1:B:359:THR:HB	1:B:362:ALA:HB3	1.68	0.75
1:D:344:ARG:O	1:D:347:THR:HG22	1.87	0.75
1:A:344[A]:ARG:HH21	1:A:347:THR:HG21	1.52	0.74
1:A:281:VAL:HG22	1:D:361:LEU:HD21	1.73	0.70
1:D:355:MET:HG3	1:D:362:ALA:HB3	1.77	0.66
1:C:353:LEU:HD13	1:C:367:ILE:HD11	1.78	0.66
1:C:353:LEU:HD13	1:C:367:ILE:CD1	2.25	0.66
1:C:329:LEU:HD11	1:C:352:ALA:HB2	1.82	0.60
1:C:355:MET:HG3	1:C:362:ALA:HB3	1.84	0.60
1:A:355:MET:HG3	1:A:362:ALA:HB3	1.85	0.58
1:C:353:LEU:CD1	1:C:367:ILE:CD1	2.82	0.58
1:B:344:ARG:HH11	1:B:347:THR:HG21	1.71	0.55
1:D:342:LEU:HD21	1:D:373:PHE:CG	2.42	0.55
1:D:313:THR:HG23	1:D:314:HIS:ND1	2.23	0.54
1:C:353:LEU:CD1	1:C:367:ILE:HD11	2.39	0.53
1:D:360:ALA:HB1	1:D:394:LEU:HD11	1.92	0.51
1:B:280:GLY:N	2:B:502:SO4:O4	2.42	0.50
1:A:359:THR:HG1	1:B:279:ILE:N	2.10	0.49
1:B:355:MET:H	1:B:363:GLU:HG2	1.78	0.48
1:A:314:HIS:CD2	1:A:356:PRO:HB2	2.49	0.48
1:C:290:TRP:CD1	1:C:386:PRO:HD3	2.49	0.48
1:A:359:THR:CG2	1:B:279:ILE:HG22	2.36	0.48
1:C:330:ARG:HD2	2:C:501:SO4:O1	2.14	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:329:LEU:HD11	1:B:352:ALA:HB2	1.96	0.47
1:D:290:TRP:CD1	1:D:386:PRO:HD3	2.49	0.47
1:A:329:LEU:HD11	1:A:352:ALA:HB2	1.97	0.46
1:A:290:TRP:CD1	1:A:386:PRO:HD3	2.52	0.45
1:B:367:ILE:HD13	1:B:386:PRO:HA	1.98	0.45
1:B:309[B]:ARG:NH1	1:D:286:LYS:HD3	2.32	0.45
1:C:314:HIS:CD2	1:C:356:PRO:HB2	2.53	0.44
1:D:285:ALA:HB1	1:D:291:HIS:CD2	2.54	0.43
1:A:361:LEU:HD21	1:B:281:VAL:HA	2.00	0.42
1:D:282:ALA:HB1	1:D:302:GLN:HE22	1.85	0.41
1:C:329:LEU:HD12	1:C:350:MET:HE3	2.03	0.41
1:C:285:ALA:HB1	1:C:291:HIS:CD2	2.56	0.41
1:A:388:LEU:O	1:B:370:GLN:HG3	2.21	0.40
1:D:313:THR:HG23	1:D:314:HIS:CE1	2.56	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [\(i\)](#)

5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	114/126 (90%)	111 (97%)	3 (3%)	0	100 100
1	B	113/126 (90%)	109 (96%)	4 (4%)	0	100 100
1	C	113/126 (90%)	108 (96%)	5 (4%)	0	100 100
1	D	115/126 (91%)	112 (97%)	3 (3%)	0	100 100
All	All	455/504 (90%)	440 (97%)	15 (3%)	0	100 100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	94/101 (93%)	94 (100%)	0	100 100
1	B	92/101 (91%)	92 (100%)	0	100 100
1	C	93/101 (92%)	91 (98%)	2 (2%)	52 39
1	D	94/101 (93%)	91 (97%)	3 (3%)	39 25
All	All	373/404 (92%)	368 (99%)	5 (1%)	73 62

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	309[A]	ARG
1	C	309[B]	ARG
1	D	345	VAL
1	D	361	LEU
1	D	389	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	302	GLN
1	B	302	GLN
1	C	302	GLN
1	C	314	HIS
1	C	346	ASN
1	D	302	GLN

5.3.3 RNA [\(i\)](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [\(i\)](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [\(i\)](#)

14 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	SO4	B	502	-	4,4,4	0.41	0	6,6,6	0.22	0
2	SO4	B	503[A]	-	4,4,4	0.33	0	6,6,6	0.19	0
2	SO4	A	501[B]	-	4,4,4	0.34	0	6,6,6	0.23	0
2	SO4	A	501[A]	-	4,4,4	0.34	0	6,6,6	0.09	0
2	SO4	B	503[B]	-	4,4,4	0.33	0	6,6,6	0.40	0
2	SO4	C	502	-	4,4,4	0.45	0	6,6,6	0.10	0
2	SO4	D	503	-	4,4,4	0.35	0	6,6,6	0.26	0
2	SO4	C	503[B]	-	4,4,4	0.32	0	6,6,6	0.34	0
2	SO4	C	503[A]	-	4,4,4	0.27	0	6,6,6	0.31	0
2	SO4	B	501	-	4,4,4	0.32	0	6,6,6	0.17	0
2	SO4	C	504	-	4,4,4	0.20	0	6,6,6	0.54	0
2	SO4	D	501	-	4,4,4	0.36	0	6,6,6	0.27	0
2	SO4	D	502	-	4,4,4	0.43	0	6,6,6	0.16	0
2	SO4	C	501	-	4,4,4	0.45	0	6,6,6	0.78	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	502	SO4	1	0
2	C	501	SO4	1	0

5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	113/126 (89%)	0.54	18 (15%) 1 1	39, 51, 93, 117	0
1	B	112/126 (88%)	0.32	10 (8%) 9 7	38, 49, 104, 134	0
1	C	113/126 (89%)	0.44	14 (12%) 4 3	37, 49, 95, 122	0
1	D	117/126 (92%)	0.65	16 (13%) 3 2	42, 56, 111, 135	0
All	All	455/504 (90%)	0.49	58 (12%) 3 2	37, 52, 102, 135	0

All (58) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	360	ALA	9.2
1	A	358	SER	8.9
1	C	358	SER	8.7
1	B	359	THR	8.6
1	D	357	GLY	8.4
1	D	358	SER	7.9
1	A	360	ALA	7.6
1	C	360	ALA	7.5
1	D	361	LEU	7.4
1	B	361	LEU	6.9
1	D	395	THR	6.2
1	C	359	THR	5.7
1	C	390	GLY	5.1
1	D	359	THR	4.4
1	B	390	GLY	4.4
1	A	359	THR	4.4
1	B	363	GLU	4.3
1	D	360	ALA	4.3
1	B	358	SER	4.0
1	C	357	GLY	4.0
1	A	391	LEU	3.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	389	ASP	3.9
1	A	387	PHE	3.8
1	A	361	LEU	3.8
1	C	361	LEU	3.6
1	A	382	PHE	3.6
1	C	389	ASP	3.4
1	A	357	GLY	3.3
1	D	292	PHE	3.3
1	B	362	ALA	3.2
1	A	339	ARG	3.1
1	D	382	PHE	3.1
1	B	389	ASP	3.1
1	C	382	PHE	3.1
1	A	390	GLY	3.0
1	D	346	ASN	3.0
1	B	387	PHE	2.9
1	D	332	ARG	2.9
1	A	381	LEU	2.8
1	A	375	ALA	2.8
1	A	363	GLU	2.8
1	C	292	PHE	2.8
1	D	376	GLU	2.6
1	A	292	PHE	2.5
1	C	346	ASN	2.4
1	C	363	GLU	2.4
1	A	380	LEU	2.3
1	D	392	GLU	2.3
1	C	339	ARG	2.3
1	D	339	ARG	2.3
1	D	363	GLU	2.2
1	D	394	LEU	2.2
1	D	393	GLY	2.2
1	C	381	LEU	2.2
1	B	292	PHE	2.1
1	C	391	LEU	2.1
1	A	376	GLU	2.0
1	A	383[A]	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [\(i\)](#)

There are no monosaccharides in this entry.

6.4 Ligands [\(i\)](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	SO4	D	501	5/5	0.70	0.25	108,113,125,137	0
2	SO4	D	503	5/5	0.81	0.18	126,141,146,152	0
2	SO4	C	503[A]	5/5	0.85	0.15	72,82,85,90	5
2	SO4	C	503[B]	5/5	0.85	0.15	77,78,85,86	5
2	SO4	A	501[A]	5/5	0.88	0.12	72,72,77,80	5
2	SO4	A	501[B]	5/5	0.88	0.12	84,85,88,90	5
2	SO4	B	503[A]	5/5	0.88	0.16	67,69,72,76	5
2	SO4	B	503[B]	5/5	0.88	0.16	73,81,87,89	5
2	SO4	D	502	5/5	0.90	0.37	100,105,107,111	0
2	SO4	C	502	5/5	0.90	0.26	116,118,123,124	0
2	SO4	C	504	5/5	0.96	0.12	81,82,101,104	0
2	SO4	B	502	5/5	0.96	0.13	94,98,100,107	0
2	SO4	B	501	5/5	0.98	0.26	96,98,100,100	0
2	SO4	C	501	5/5	0.99	0.08	73,75,77,79	0

6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.