



# Full wwPDB X-ray Structure Validation Report ⓘ

May 30, 2022 – 04:16 pm BST

PDB ID : 6GJ2  
Title : PURPLE ACID PHYTASE FROM WHEAT ISOFORM B2 - COMPLEX  
WITH INOSITOL HEXASULPHATE  
Authors : Faba-Rodriguez, R.; Brearley, C.A.; Hemmings, A.M.  
Deposited on : 2018-05-15  
Resolution : 1.68 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.4, CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.28.1  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0267  
CCP4 : 7.1.010 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.28.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*



The reported resolution of this entry is 1.68 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	6780 (1.70-1.66)
Clashscore	141614	7310 (1.70-1.66)
Ramachandran outliers	138981	7173 (1.70-1.66)
Sidechain outliers	138945	7172 (1.70-1.66)
RSRZ outliers	127900	6661 (1.70-1.66)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	516	
2	B	2	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
9	EDO	A	622	-	-	X	-

## 2 Entry composition [i](#)

There are 11 unique types of molecules in this entry. The entry contains 4748 atoms, of which 251 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Purple acid phosphatase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	504	3966	2522	670	750	24	0	7	0

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	511	HIS	-	expression tag	UNP C4PKL0
A	512	HIS	-	expression tag	UNP C4PKL0
A	513	HIS	-	expression tag	UNP C4PKL0
A	514	HIS	-	expression tag	UNP C4PKL0
A	515	HIS	-	expression tag	UNP C4PKL0
A	516	HIS	-	expression tag	UNP C4PKL0

- Molecule 2 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	H	N	O			
2	B	2	55	16	27	2	10	0	0	0

- Molecule 3 is FE (III) ION (three-letter code: FE) (formula: Fe).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	2	Total	Fe	0	0
			2	2		

- Molecule 4 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: C<sub>8</sub>H<sub>15</sub>NO<sub>6</sub>).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
			Total	C	H	N			O
4	A	1	Total	C	H	N	O	0	0
			28	8	14	1	5		
4	A	1	Total	C	H	N	O	0	0
			28	8	14	1	5		
4	A	1	Total	C	H	N	O	0	0
			28	8	14	1	5		
4	A	1	Total	C	H	N	O	0	0
			28	8	14	1	5		

- Molecule 5 is DI(HYDROXYETHYL)ETHER (three-letter code: PEG) (formula: C<sub>4</sub>H<sub>10</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	1	Total	C	H	O	0	0
			17	4	10	3		
5	A	1	Total	C	H	O	0	0
			17	4	10	3		
5	A	1	Total	C	H	O	0	0
			17	4	10	3		
5	A	1	Total	C	H	O	0	0
			17	4	10	3		

- Molecule 6 is TRIETHYLENE GLYCOL (three-letter code: PGE) (formula: C<sub>6</sub>H<sub>14</sub>O<sub>4</sub>).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
6	A	1	Total	C	H	O	0	0
			24	6	14	4		
6	A	1	Total	C	H	O	0	0
			24	6	14	4		
6	A	1	Total	C	H	O	0	0
			24	6	14	4		
6	A	1	Total	C	H	O	0	0
			24	6	14	4		

- Molecule 7 is PHOSPHATE ION (three-letter code: PO4) (formula: O<sub>4</sub>P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	A	1	Total	O	P	0	0
			5	4	1		

- Molecule 8 is TETRAETHYLENE GLYCOL (three-letter code: PG4) (formula:  $C_8H_{18}O_5$ ).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
8	A	1	Total	C	H	O	0	0
			31	8	18	5		

- Molecule 9 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula:  $C_2H_6O_2$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
9	A	1	Total	C	H	O	0	0
			10	2	6	2		
9	A	1	Total	C	H	O	0	0
			10	2	6	2		
9	A	1	Total	C	H	O	0	0
			10	2	6	2		
9	A	1	Total	C	H	O	0	0
			10	2	6	2		

- Molecule 10 is D-MYO-INOSITOL-HEXASULPHATE (three-letter code: IHS) (formula:  $C_6H_{12}O_{24}S_6$ ).





Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	H	O	S		
10	A	1	42	6	6	24	6	0	0


- Molecule 11 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
11	A	286	Total	O	0	0
			286	286		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Purple acid phosphatase



- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



## 4 Data and refinement statistics

Property	Value	Source
Space group	H 3	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	126.02Å 126.02Å 105.91Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	38.44 – 1.68 38.44 – 1.68	Depositor EDS
% Data completeness (in resolution range)	99.9 (38.44-1.68) 99.9 (38.44-1.68)	Depositor EDS
$R_{merge}$	0.06	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.56 (at 1.68Å)	Xtrriage
Refinement program	PHENIX	Depositor
R, $R_{free}$	0.134 , 0.176 0.136 , 0.132	Depositor DCC
$R_{free}$ test set	3791 reflections (5.31%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	26.2	Xtrriage
Anisotropy	0.240	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	(Not available) , (Not available)	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	0.020 for h,-h-k,-l	Xtrriage
$F_o, F_c$ correlation	0.98	EDS
Total number of atoms	4748	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	37.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.15% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: IHS, PG4, PO4, EDO, PEG, PGE, FE, NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.34	0/4111	0.52	0/5609

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3966	0	3722	42	0
2	B	28	27	25	0	0
3	A	2	0	0	0	0
4	A	70	70	65	2	0
5	A	35	50	50	8	0
6	A	40	56	56	7	0
7	A	5	0	0	0	0
8	A	13	18	18	0	0
9	A	16	24	24	4	0
10	A	36	6	12	1	0
11	A	286	0	0	1	0
All	All	4497	251	3972	43	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 5.

All (43) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:213:THR:OG1	4:A:605:NAG:N2	2.18	0.74
1:A:134:LYS:HE2	6:A:616:PGE:H62	1.73	0.70
1:A:446:ALA:HA	5:A:613:PEG:H22	1.76	0.68
1:A:315:GLU:HG3	9:A:622:EDO:H11	1.78	0.65
1:A:64:GLY:HA2	6:A:617:PGE:H3	1.78	0.64
1:A:219:SER:HB3	10:A:625:IHS:H3	1.81	0.62
1:A:319:TRP:HB2	9:A:622:EDO:H12	1.80	0.62
1:A:315:GLU:HG3	9:A:622:EDO:C1	2.31	0.60
1:A:95:ARG:HH21	5:A:614:PEG:H22	1.66	0.60
1:A:213:THR:HG1	4:A:605:NAG:HN2	1.46	0.59
1:A:429:PRO:CG	1:A:436:PHE:HD1	2.16	0.58
1:A:163:ARG:NH2	5:A:612:PEG:H31	2.20	0.56
1:A:219:SER:HA	1:A:223:ALA:HB2	1.88	0.56
1:A:105:SER:H	5:A:611:PEG:H11	1.71	0.54
1:A:138[B]:GLN:HG3	1:A:148:MET:CE	2.37	0.54
1:A:134:LYS:HE2	6:A:616:PGE:C6	2.37	0.53
1:A:476[A]:GLU:HG3	1:A:477:THR:HG23	1.91	0.53
1:A:429:PRO:HG3	1:A:436:PHE:HD1	1.72	0.52
1:A:138[A]:GLN:HG2	1:A:149:SER:O	2.10	0.51
1:A:95:ARG:HB3	5:A:614:PEG:H32	1.92	0.51
1:A:318:ARG:HG2	9:A:622:EDO:H22	1.92	0.51
1:A:370:LEU:HD21	1:A:373:VAL:HG22	1.93	0.51
1:A:469:GLY:HA3	1:A:481:TRP:CH2	2.45	0.51
1:A:318:ARG:HE	1:A:322:LYS:HZ1	1.60	0.49
1:A:452:TRP:HA	5:A:613:PEG:H21	1.94	0.48
5:A:610:PEG:H12	6:A:616:PGE:H12	1.96	0.46
1:A:43:PRO:HB3	1:A:61:TRP:CD1	2.51	0.46
1:A:326:LYS:HD2	6:A:615:PGE:H52	1.99	0.45
1:A:138[B]:GLN:HG3	1:A:148:MET:HE2	1.98	0.45
1:A:36[B]:ARG:NH2	1:A:40:GLY:O	2.49	0.45
1:A:379:HIS:HE1	1:A:409:GLU:OE1	2.00	0.45
1:A:105:SER:H	5:A:611:PEG:C1	2.31	0.44
1:A:134:LYS:NZ	6:A:616:PGE:H6	2.32	0.44
1:A:429:PRO:HG3	1:A:436:PHE:CD1	2.53	0.43
1:A:339:TRP:O	1:A:375:THR:HA	2.19	0.43
1:A:138[B]:GLN:HG3	1:A:148:MET:HE3	2.01	0.42
1:A:326:LYS:NZ	6:A:615:PGE:H52	2.35	0.41
1:A:428:THR:N	1:A:429:PRO:CD	2.83	0.41

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:426:MET:CE	1:A:426:MET:HA	2.50	0.41
1:A:505:GLU:HG3	11:A:797:HOH:O	2.20	0.41
1:A:318:ARG:NE	1:A:322:LYS:HZ1	2.19	0.41
1:A:429:PRO:HG2	1:A:436:PHE:HD1	1.85	0.40
1:A:26:ASP:OD1	1:A:178:THR:HG22	2.21	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	507/516 (98%)	488 (96%)	18 (4%)	1 (0%)	47 29

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	70	GLY

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	417/425 (98%)	413 (99%)	4 (1%)	76 65

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	4	SER
1	A	36[A]	ARG
1	A	36[B]	ARG
1	A	401	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

2 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	NAG	B	1	2,1	14,14,15	1.77	2 (14%)	17,19,21	1.02	1 (5%)
2	NAG	B	2	2	14,14,15	1.97	4 (28%)	17,19,21	1.12	2 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	B	1	2,1	-	0/6/23/26	0/1/1/1
2	NAG	B	2	2	-	0/6/23/26	0/1/1/1

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	2	NAG	O5-C1	4.29	1.50	1.43
2	B	1	NAG	O5-C1	4.15	1.50	1.43
2	B	2	NAG	C7-N2	3.64	1.46	1.34
2	B	1	NAG	C7-N2	3.38	1.46	1.34
2	B	2	NAG	C2-N2	2.57	1.50	1.46
2	B	2	NAG	O5-C5	2.00	1.47	1.43

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	2	NAG	C8-C7-N2	3.01	121.19	116.10
2	B	1	NAG	C1-C2-N2	-2.66	105.95	110.49
2	B	2	NAG	C2-N2-C7	-2.00	120.05	122.90

There are no chirality outliers.


There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





## 5.6 Ligand geometry [i](#)

Of 23 ligands modelled in this entry, 2 are monoatomic - leaving 21 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
4	NAG	A	606	1	14,14,15	1.94	3 (21%)	17,19,21	1.07	1 (5%)
9	EDO	A	623	-	3,3,3	0.48	0	2,2,2	0.29	0
5	PEG	A	611	-	6,6,6	0.47	0	5,5,5	0.76	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
9	EDO	A	624	-	3,3,3	0.47	0	2,2,2	0.28	0
6	PGE	A	615	-	9,9,9	0.52	0	8,8,8	0.35	0
9	EDO	A	621	-	3,3,3	0.47	0	2,2,2	0.31	0
5	PEG	A	614	-	6,6,6	0.49	0	5,5,5	0.24	0
6	PGE	A	616	-	9,9,9	0.53	0	8,8,8	0.88	0
6	PGE	A	618	-	9,9,9	0.51	0	8,8,8	0.50	0
7	PO4	A	619	3	4,4,4	0.90	0	6,6,6	0.59	0
4	NAG	A	604	1	14,14,15	1.66	2 (14%)	17,19,21	0.93	1 (5%)
5	PEG	A	613	-	6,6,6	0.46	0	5,5,5	0.34	0
8	PG4	A	620	-	12,12,12	0.52	0	11,11,11	0.35	0
9	EDO	A	622	-	3,3,3	0.45	0	2,2,2	0.27	0
10	IHS	A	625	-	36,36,36	1.84	12 (33%)	42,60,60	1.31	6 (14%)
5	PEG	A	610	-	6,6,6	0.50	0	5,5,5	0.33	0
4	NAG	A	603	1	14,14,15	1.77	2 (14%)	17,19,21	1.09	1 (5%)
4	NAG	A	607	1	14,14,15	1.84	2 (14%)	17,19,21	1.18	2 (11%)
5	PEG	A	612	-	6,6,6	0.51	0	5,5,5	0.48	0
6	PGE	A	617	-	9,9,9	0.51	0	8,8,8	0.87	0
4	NAG	A	605	1	14,14,15	1.88	3 (21%)	17,19,21	1.27	3 (17%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	A	606	1	-	0/6/23/26	0/1/1/1
9	EDO	A	623	-	-	0/1/1/1	-
5	PEG	A	611	-	-	1/4/4/4	-
9	EDO	A	624	-	-	1/1/1/1	-
6	PGE	A	615	-	-	3/7/7/7	-
9	EDO	A	621	-	-	1/1/1/1	-
5	PEG	A	614	-	-	1/4/4/4	-
6	PGE	A	616	-	-	0/7/7/7	-
6	PGE	A	618	-	-	4/7/7/7	-
4	NAG	A	604	1	-	0/6/23/26	0/1/1/1
5	PEG	A	613	-	-	0/4/4/4	-
8	PG4	A	620	-	-	1/10/10/10	-
9	EDO	A	622	-	-	0/1/1/1	-
10	IHS	A	625	-	-	2/30/54/54	0/1/1/1
5	PEG	A	610	-	-	0/4/4/4	-
4	NAG	A	603	1	-	0/6/23/26	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	A	607	1	-	0/6/23/26	0/1/1/1
5	PEG	A	612	-	-	0/4/4/4	-
6	PGE	A	617	-	-	4/7/7/7	-
4	NAG	A	605	1	-	0/6/23/26	0/1/1/1

All (24) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	606	NAG	O5-C1	4.62	1.51	1.43
4	A	607	NAG	O5-C1	4.24	1.50	1.43
4	A	603	NAG	O5-C1	4.11	1.50	1.43
4	A	605	NAG	O5-C1	3.99	1.50	1.43
4	A	604	NAG	O5-C1	3.75	1.49	1.43
4	A	605	NAG	C7-N2	3.64	1.46	1.34
4	A	606	NAG	C7-N2	3.43	1.46	1.34
4	A	607	NAG	C7-N2	3.43	1.46	1.34
4	A	603	NAG	C7-N2	3.34	1.45	1.34
4	A	604	NAG	C7-N2	3.26	1.45	1.34
10	A	625	IHS	O1-S1	3.16	1.66	1.57
10	A	625	IHS	O15-S5	3.11	1.66	1.57
10	A	625	IHS	O12-S2	2.98	1.66	1.57
10	A	625	IHS	O16-S6	2.91	1.65	1.57
10	A	625	IHS	O13-S3	2.87	1.65	1.57
10	A	625	IHS	O14-S4	2.78	1.65	1.57
4	A	605	NAG	C2-N2	2.67	1.50	1.46
10	A	625	IHS	O14-C4	-2.63	1.41	1.46
10	A	625	IHS	O13-C3	-2.62	1.41	1.46
10	A	625	IHS	O12-C2	-2.58	1.41	1.46
10	A	625	IHS	O16-C6	-2.48	1.41	1.46
10	A	625	IHS	O15-C5	-2.46	1.41	1.46
10	A	625	IHS	O1-C1	-2.30	1.41	1.46
4	A	606	NAG	C2-N2	2.12	1.49	1.46

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	607	NAG	C2-N2-C7	-2.95	118.70	122.90
10	A	625	IHS	O4-S1-O3	-2.91	100.55	112.22
10	A	625	IHS	O36-S6-O26	-2.86	100.76	112.22
10	A	625	IHS	O42-S2-O22	-2.84	100.81	112.22
10	A	625	IHS	O43-S3-O23	-2.77	101.08	112.22
10	A	625	IHS	O34-S4-O24	-2.71	101.36	112.22

Continued on next page...

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	605	NAG	C8-C7-N2	2.65	120.58	116.10
4	A	603	NAG	C2-N2-C7	-2.55	119.28	122.90
4	A	607	NAG	C8-C7-N2	2.44	120.23	116.10
4	A	605	NAG	C1-C2-N2	-2.17	106.78	110.49
4	A	604	NAG	C2-N2-C7	-2.16	119.83	122.90
10	A	625	IHS	O35-S5-O25	-2.15	101.02	108.49
4	A	606	NAG	C1-O5-C5	-2.14	109.30	112.19
4	A	605	NAG	O5-C1-C2	2.04	114.50	111.29

There are no chirality outliers.

All (18) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
6	A	617	PGE	C4-C3-O2-C2
6	A	615	PGE	O2-C3-C4-O3
6	A	617	PGE	O2-C3-C4-O3
6	A	618	PGE	C4-C3-O2-C2
6	A	615	PGE	O3-C5-C6-O4
5	A	611	PEG	C1-C2-O2-C3
5	A	614	PEG	C4-C3-O2-C2
6	A	617	PGE	C6-C5-O3-C4
6	A	615	PGE	C3-C4-O3-C5
6	A	618	PGE	C3-C4-O3-C5
10	A	625	IHS	C1-C6-O16-S6
6	A	617	PGE	C3-C4-O3-C5
9	A	621	EDO	O1-C1-C2-O2
9	A	624	EDO	O1-C1-C2-O2
10	A	625	IHS	C6-O16-S6-O46
8	A	620	PG4	O3-C5-C6-O4
6	A	618	PGE	C6-C5-O3-C4
6	A	618	PGE	O2-C3-C4-O3

There are no ring outliers.

11 monomers are involved in 21 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	611	PEG	2	0
6	A	615	PGE	2	0
5	A	614	PEG	2	0
6	A	616	PGE	4	0
5	A	613	PEG	2	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Clashes	Symm-Clashes
9	A	622	EDO	4	0
10	A	625	IHS	1	0
5	A	610	PEG	1	0
5	A	612	PEG	1	0
6	A	617	PGE	1	0
4	A	605	NAG	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	504/516 (97%)	-0.13	23 (4%) <span style="border: 1px solid red; padding: 2px;">32</span> <span style="border: 1px solid red; padding: 2px;">35</span>	20, 29, 52, 76	0

All (23) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	17	LEU	7.2
1	A	220	CYS	6.0
1	A	223	ALA	4.8
1	A	222	PHE	4.7
1	A	1	GLU	4.3
1	A	2	PRO	3.5
1	A	419	PRO	3.3
1	A	18	ARG	3.1
1	A	145	PRO	3.1
1	A	221	SER	3.0
1	A	225	SER	2.9
1	A	109	PRO	2.9
1	A	110	PHE	2.8
1	A	224	LYS	2.6
1	A	421	ARG	2.6
1	A	144	ILE	2.4
1	A	16	PRO	2.4
1	A	265	ILE	2.3
1	A	41	TRP	2.2
1	A	420	GLY	2.2
1	A	199	LEU	2.1
1	A	79	THR	2.1
1	A	227	PRO	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)


There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	NAG	B	2	14/15	0.77	0.27	57,69,87,90	0
2	NAG	B	1	14/15	0.95	0.13	37,53,65,65	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.



## 6.4 Ligands [i](#)


In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,



median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
9	EDO	A	623	4/4	0.73	0.16	53,65,73,78	0
5	PEG	A	613	7/7	0.78	0.26	70,84,89,90	0
4	NAG	A	605	14/15	0.78	0.18	66,79,92,94	0
5	PEG	A	614	7/7	0.79	0.13	59,71,76,78	0
5	PEG	A	612	7/7	0.81	0.20	59,70,84,86	0
4	NAG	A	606	14/15	0.81	0.23	54,67,80,83	28
10	IHS	A	625	36/36	0.81	0.15	95,116,139,146	0
9	EDO	A	624	4/4	0.83	0.12	71,85,90,91	0
6	PGE	A	617	10/10	0.83	0.27	60,75,90,92	0
6	PGE	A	618	10/10	0.84	0.13	66,81,92,92	0
8	PG4	A	620	13/13	0.86	0.11	58,72,87,88	0
6	PGE	A	616	10/10	0.86	0.14	67,80,88,90	0
5	PEG	A	610	7/7	0.87	0.15	62,74,86,89	0
9	EDO	A	621	4/4	0.88	0.17	69,83,84,85	0
9	EDO	A	622	4/4	0.89	0.16	66,80,80,81	0
5	PEG	A	611	7/7	0.89	0.21	66,79,80,80	0
6	PGE	A	615	10/10	0.90	0.12	51,64,78,78	0
3	FE	A	601	1/1	0.95	0.09	56,56,56,56	1
4	NAG	A	607	14/15	0.96	0.20	46,58,72,72	0
4	NAG	A	603	14/15	0.96	0.18	38,48,57,61	0
4	NAG	A	604	14/15	0.98	0.06	29,37,46,48	0
7	PO4	A	619	5/5	0.98	0.10	38,46,52,63	0
3	FE	A	602	1/1	1.00	0.13	20,20,20,20	1

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



## 6.5 Other polymers [i](#)

There are no such residues in this entry.