



wwPDB X-ray Structure Validation Summary Report ⓘ

Nov 5, 2023 – 02:55 PM EST

PDB ID : 4GK7
Title : yeast 20S proteasome in complex with the Syringolin-Glidobactin chimera
Authors : Groll, M.; Stein, M.L.; Bachmann, A.
Deposited on : 2012-08-10
Resolution : 2.80 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3140 (2.80-2.80)
Clashscore	141614	3569 (2.80-2.80)
Ramachandran outliers	138981	3498 (2.80-2.80)
Sidechain outliers	138945	3500 (2.80-2.80)
RSRZ outliers	127900	3078 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	250	 71% 27% 2% 0%
1	O	250	 69% 29% 2% 0%
2	B	244	 64% 29% 7% 3%
2	P	244	 65% 28% 7% 3%
3	C	241	 59% 38% 3% 3%





Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	Q	241	 5% 59% 38% .
4	D	242	 3% 71% 26% .
4	R	242	 3% 70% 28% .
5	E	233	 3% 58% 36% 7%
5	S	233	 2% 57% 36% 7%
6	F	244	 3% 65% 31% .
6	T	244	 2% 64% 32% .
7	G	243	 % 62% 33% 5%
7	U	243	 % 63% 32% 5%
8	H	222	 % 69% 28% .
8	V	222	 % 69% 28% .
9	I	204	 % 69% 29% .
9	W	204	 % 68% 30% .
10	J	198	 2% 67% 31% .
10	X	198	 2% 65% 33% .
11	K	212	 % 70% 26% .
11	Y	212	 % 70% 26% .
12	L	222	 % 65% 30% .
12	Z	222	 % 67% 28% 5%
13	M	233	 % 67% 30% .
13	a	233	 % 94% 6%
14	N	196	 % 73% 23% .
14	b	196	 % 95% 5%
15	1	4	 25% 50% 25%
15	2	4	 50% 50%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
15	3	4	 25% 25% 50%
15	4	4	 75% 25%
15	5	4	 25% 75%
15	6	4	 25% 75%

2 Entry composition

There are 16 unique types of molecules in this entry. The entry contains 51109 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Proteasome component Y7.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	250	Total	C	N	O	S	0	0	0
			1915	1219	315	377	4			
1	O	250	Total	C	N	O	S	0	0	0
			1915	1219	315	377	4			

- Molecule 2 is a protein called Proteasome component Y13.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	244	Total	C	N	O	S	0	0	0
			1905	1201	321	380	3			
2	P	244	Total	C	N	O	S	0	0	0
			1905	1201	321	380	3			

- Molecule 3 is a protein called Proteasome component PRE6.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	C	241	Total	C	N	O	S	0	0	0
			1891	1181	331	375	4			
3	Q	241	Total	C	N	O	S	0	0	0
			1891	1181	331	375	4			

- Molecule 4 is a protein called Proteasome component PUP2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
4	D	242	Total	C	N	O	S	0	0	0
			1862	1162	314	379	7			
4	R	242	Total	C	N	O	S	0	0	0
			1862	1162	314	379	7			

- Molecule 5 is a protein called Proteasome component PRE5.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
5	E	233	Total	C	N	O	S	0	0	0
			1795	1129	312	350	4			
5	S	233	Total	C	N	O	S	0	0	0
			1795	1129	312	350	4			

- Molecule 6 is a protein called Proteasome component C1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
6	F	244	Total	C	N	O	S	0	0	0
			1897	1205	330	358	4			
6	T	244	Total	C	N	O	S	0	0	0
			1897	1205	330	358	4			

- Molecule 7 is a protein called Proteasome component C7-alpha.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
7	G	243	Total	C	N	O	S	0	0	0
			1921	1221	322	370	8			
7	U	243	Total	C	N	O	S	0	0	0
			1921	1221	322	370	8			

- Molecule 8 is a protein called Proteasome component PUP1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
8	H	222	Total	C	N	O	S	0	0	0
			1685	1061	293	324	7			
8	V	222	Total	C	N	O	S	0	0	0
			1685	1061	293	324	7			

- Molecule 9 is a protein called Proteasome component PUP3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
9	I	204	Total	C	N	O	S	0	0	0
			1581	1010	258	305	8			
9	W	204	Total	C	N	O	S	0	0	0
			1581	1010	258	305	8			

- Molecule 10 is a protein called Proteasome component C11.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
10	J	198	Total	C	N	O	S	0	0	0
			1585	1005	269	305	6			

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
10	X	198	Total	C	N	O	S	0	0	0
			1585	1005	269	305	6			

- Molecule 11 is a protein called Proteasome component PRE2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
11	K	212	Total	C	N	O	S	0	0	0
			1644	1045	280	312	7			
11	Y	212	Total	C	N	O	S	0	0	0
			1644	1045	280	312	7			

- Molecule 12 is a protein called Proteasome component C5.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
12	L	222	Total	C	N	O	S	0	0	0
			1757	1115	303	335	4			
12	Z	222	Total	C	N	O	S	0	0	0
			1757	1115	303	335	4			

- Molecule 13 is a protein called Proteasome component PRE4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
13	M	233	Total	C	N	O	S	0	0	0
			1824	1154	312	351	7			
13	a	233	Total	C	N	O	S	0	0	0
			1824	1154	312	351	7			

- Molecule 14 is a protein called Proteasome component PRE3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
14	N	196	Total	C	N	O	S	0	0	0
			1512	955	250	300	7			
14	b	196	Total	C	N	O	S	0	0	0
			1512	955	250	300	7			

- Molecule 15 is a protein called SYRINGOLIN-GLIDOBACTIN CHIMERA.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
15	1	4	Total	C	N	O	0	0	0
			38	29	4	5			
15	2	4	Total	C	N	O	0	0	0
			38	29	4	5			

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
15	3	4	Total	C	N	O	0	0	0
			38	29	4	5			
15	4	4	Total	C	N	O	0	0	0
			38	29	4	5			
15	5	4	Total	C	N	O	0	0	0
			38	29	4	5			
15	6	4	Total	C	N	O	0	0	0
			38	29	4	5			

- Molecule 16 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
16	A	59	Total	O	0	0
			59	59		
16	B	35	Total	O	0	0
			35	35		
16	C	44	Total	O	0	0
			44	44		
16	D	39	Total	O	0	0
			39	39		
16	E	23	Total	O	0	0
			23	23		
16	F	49	Total	O	0	0
			49	49		
16	G	60	Total	O	0	0
			60	60		
16	H	49	Total	O	0	0
			49	49		
16	I	64	Total	O	0	0
			64	64		
16	J	53	Total	O	0	0
			53	53		
16	K	42	Total	O	0	0
			42	42		
16	L	55	Total	O	0	0
			55	55		
16	M	75	Total	O	0	0
			75	75		
16	N	56	Total	O	0	0
			56	56		
16	O	34	Total	O	0	0
			34	34		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
16	P	28	Total 28	O 28	0	0
16	Q	29	Total 29	O 29	0	0
16	R	33	Total 33	O 33	0	0
16	S	21	Total 21	O 21	0	0
16	T	39	Total 39	O 39	0	0
16	U	63	Total 63	O 63	0	0
16	V	47	Total 47	O 47	0	0
16	W	55	Total 55	O 55	0	0
16	X	46	Total 46	O 46	0	0
16	Y	50	Total 50	O 50	0	0
16	Z	51	Total 51	O 51	0	0
16	a	73	Total 73	O 73	0	0
16	b	58	Total 58	O 58	0	0
16	1	1	Total 1	O 1	0	0
16	2	1	Total 1	O 1	0	0
16	3	1	Total 1	O 1	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Proteasome component Y7




- Molecule 1: Proteasome component Y7






- Molecule 2: Proteasome component Y13









- Molecule 2: Proteasome component Y13



- Molecule 3: Proteasome component PRE6




- Molecule 3: Proteasome component PRE6




- Molecule 4: Proteasome component PUP2







- Molecule 4: Proteasome component PUP2




- Molecule 5: Proteasome component PRE5




- Molecule 5: Proteasome component PRE5




- Molecule 6: Proteasome component C1




• Molecule 6: Proteasome component C1




• Molecule 7: Proteasome component C7-alpha




• Molecule 7: Proteasome component C7-alpha







- Molecule 8: Proteasome component PUP1




- Molecule 8: Proteasome component PUP1




- Molecule 9: Proteasome component PUP3




- Molecule 9: Proteasome component PUP3




- Molecule 10: Proteasome component C11




- Molecule 10: Proteasome component C11




- Molecule 11: Proteasome component PRE2



- Molecule 11: Proteasome component PRE2



- Molecule 12: Proteasome component C5



K210
R211
D212

- Molecule 12: Proteasome component C5

Chain Z:  67% 28% 5%

Q-9 F-8 N-7 P-6 Y-5 N-2 G-1 T1 T2 L3 A6 G7 E8 D9 F10 L13 R18 M19 I20 S24 I25 R28 K32 V33 F34 N39 I40 V41 M42 N45 R57 N60 S61 N70 D71 A80 R81 R82 I83 Q84 K90 F93 P94 Y95

V96 V97 H98 T99 I100 I101 K109 V112 F115 D116 F117 V118 G119 S120 I136 M137 L140 D141 N142 F146 Q149 K163 Y164 L165 S166 V167 E168 E169 V170 I171 S177 A181 H185 I186 Q187 D190 V197 T198 K199 D200 I83 Q201 V202 K210 R211 D212

- Molecule 13: Proteasome component PRE4

Chain M:  67% 30%

T-8 P-5 I-4 V-3 T0 S1 V2 M5 K6 Y7 D8 N9 G10 V11 I12 D16 M17 L18 L25 R32 V36 N39 T40 V41 S45 I48 I49 Q82 D60 L61 N69 A72 E79 P80 S81 Y82 I83 F84 E85 Y86 L87 A88 M91 R95

W102 M103 A104 I105 I106 Q111 L118 R119 V120 Y121 L133 A134 T135 R147 V150 D151 R152 E153 I156 P157 T160 V161 Q162 E165 I168 V169 M170 R173 S183 F186 I191 D192 T195 G196 L197 Q204 I216 K217 G218 Q222 K223 I224

- Molecule 13: Proteasome component PRE4

Chain a:  94% 6%

T-8 I-4 S1 N39 D60 L61 R95 L101 W102 I106 V150 D151 R152 Q162 I191 I216 K217 I224

- Molecule 14: Proteasome component PRE3

Chain N:  73% 23%

T1 S2 I3 F8 K9 D10 I13 L14 G15 A16 R19 T20 T22 I26 T31 R36 D39 K40 W42 W43 C44 I55 V59 Q60 Y66 P73 K83 E84 E88 L93 I98 D103 D104 K105 N106 K107 G108 P114 L115 G116 G117

S118 E144 M145 M146 S147 E150 T151 L159 I163 G170 R174 M175 V176 V177 L178 L186 Q195 L196

- Molecule 14: Proteasome component PRE3

Chain b:  95% 5%

T1 I26 R36 I41 R83 E88 K105 L115 L178 Q195 L196

- Molecule 15: SYRINGOLIN-GLIDOBACTIN CHIMERA

Chain 1:  25% 50% 25%




● Molecule 15: SYRINGOLIN-GLIDOBACTIN CHIMERA

Chain 2:  50% 50%




● Molecule 15: SYRINGOLIN-GLIDOBACTIN CHIMERA

Chain 3:  25% 25% 50%




● Molecule 15: SYRINGOLIN-GLIDOBACTIN CHIMERA

Chain 4:  75% 25%




● Molecule 15: SYRINGOLIN-GLIDOBACTIN CHIMERA

Chain 5:  25% 75%



● Molecule 15: SYRINGOLIN-GLIDOBACTIN CHIMERA

Chain 6:  25% 75%



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	135.85Å 298.45Å 145.25Å 90.00° 112.63° 90.00°	Depositor
Resolution (Å)	15.00 – 2.80 29.84 – 2.80	Depositor EDS
% Data completeness (in resolution range)	97.8 (15.00-2.80) 97.9 (29.84-2.80)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.94 (at 2.80Å)	Xtrriage
Refinement program	CNS	Depositor
R, R_{free}	0.210 , 0.241 0.206 , 0.205	Depositor DCC
R_{free} test set	12775 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	50.3	Xtrriage
Anisotropy	0.648	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 57.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.29$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	51109	wwPDB-VP
Average B, all atoms (Å ²)	56.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.16% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: LYH, MH9, OJT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.37	0/1952	0.64	0/2642
1	O	0.38	0/1952	0.63	0/2642
2	B	0.38	0/1935	0.64	0/2618
2	P	0.39	0/1935	0.64	0/2618
3	C	0.36	0/1920	0.62	0/2598
3	Q	0.38	0/1920	0.62	0/2598
4	D	0.36	0/1887	0.64	0/2541
4	R	0.36	0/1887	0.64	0/2541
5	E	0.36	0/1823	0.63	0/2463
5	S	0.36	0/1823	0.63	0/2463
6	F	0.39	0/1937	0.62	0/2614
6	T	0.39	0/1937	0.62	0/2614
7	G	0.41	0/1959	0.64	0/2652
7	U	0.39	0/1959	0.63	0/2652
8	H	0.41	0/1716	0.66	0/2326
8	V	0.39	0/1716	0.66	0/2326
9	I	0.39	0/1611	0.67	0/2174
9	W	0.41	0/1611	0.67	0/2174
10	J	0.40	0/1613	0.65	0/2173
10	X	0.40	0/1613	0.65	0/2173
11	K	0.41	0/1681	0.66	1/2274 (0.0%)
11	Y	0.39	0/1681	0.66	1/2274 (0.0%)
12	L	0.40	0/1795	0.69	3/2420 (0.1%)
12	Z	0.39	0/1795	0.69	3/2420 (0.1%)
13	M	0.39	0/1855	0.67	0/2514
13	a	0.38	0/1855	0.67	1/2514 (0.0%)
14	N	0.40	0/1541	0.64	0/2087
14	b	0.39	0/1541	0.64	0/2087
15	1	1.68	0/6	1.40	0/7
15	2	1.23	0/6	1.89	0/7
15	3	1.71	0/6	1.64	0/7
15	4	1.75	0/6	1.44	0/7

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
15	5	1.31	0/6	1.84	0/7
15	6	1.69	0/6	1.68	0/7
All	All	0.39	0/50486	0.65	9/68234 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
12	L	0	1
12	Z	0	1
All	All	0	2

There are no bond length outliers.

The worst 5 of 9 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
12	L	164	TYR	CA-CB-CG	-5.36	103.22	113.40
11	Y	4	LEU	CA-CB-CG	5.32	127.53	115.30
11	K	4	LEU	CA-CB-CG	5.29	127.48	115.30
12	Z	98	HIS	N-CA-C	-5.27	96.77	111.00
12	Z	164	TYR	CA-CB-CG	-5.27	103.39	113.40

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
12	L	164	TYR	Sidechain
12	Z	164	TYR	Sidechain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1915	0	1926	66	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	O	1915	0	1926	68	0
2	B	1905	0	1901	78	0
2	P	1905	0	1901	84	0
3	C	1891	0	1900	91	0
3	Q	1891	0	1900	88	0
4	D	1862	0	1836	52	0
4	R	1862	0	1836	55	0
5	E	1795	0	1797	95	0
5	S	1795	0	1797	97	0
6	F	1897	0	1886	77	0
6	T	1897	0	1886	79	0
7	G	1921	0	1910	84	0
7	U	1921	0	1910	87	0
8	H	1685	0	1687	57	0
8	V	1685	0	1687	57	0
9	I	1581	0	1574	59	0
9	W	1581	0	1574	59	0
10	J	1585	0	1590	73	0
10	X	1585	0	1590	73	0
11	K	1644	0	1594	54	0
11	Y	1644	0	1594	57	0
12	L	1757	0	1711	64	0
12	Z	1757	0	1711	60	0
13	M	1824	0	1832	60	0
13	a	1824	0	1832	0	0
14	N	1512	0	1480	45	0
14	b	1512	0	1480	0	0
15	1	38	0	34	4	0
15	2	38	0	34	0	0
15	3	38	0	34	2	0
15	4	38	0	34	4	0
15	5	38	0	34	1	0
15	6	38	0	34	0	0
16	1	1	0	0	0	0
16	2	1	0	0	0	0
16	3	1	0	0	0	0
16	A	59	0	0	1	0
16	B	35	0	0	0	0
16	C	44	0	0	5	0
16	D	39	0	0	4	0
16	E	23	0	0	2	0
16	F	49	0	0	2	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
16	G	60	0	0	6	0
16	H	49	0	0	3	0
16	I	64	0	0	2	0
16	J	53	0	0	2	0
16	K	42	0	0	3	0
16	L	55	0	0	4	0
16	M	75	0	0	3	0
16	N	56	0	0	3	0
16	O	34	0	0	1	0
16	P	28	0	0	1	0
16	Q	29	0	0	2	0
16	R	33	0	0	2	0
16	S	21	0	0	0	0
16	T	39	0	0	2	0
16	U	63	0	0	2	0
16	V	47	0	0	5	0
16	W	55	0	0	2	0
16	X	46	0	0	4	0
16	Y	50	0	0	5	0
16	Z	51	0	0	2	0
16	a	73	0	0	0	0
16	b	58	0	0	0	0
All	All	51109	0	49452	1642	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 18.

The worst 5 of 1642 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:P:203:THR:HG22	2:P:205:SER:H	1.08	1.13
11:K:107:LYS:H	11:K:107:LYS:HD2	1.03	1.11
11:Y:107:LYS:H	11:Y:107:LYS:HD2	1.04	1.11
2:B:203:THR:HG22	2:B:205:SER:H	1.11	1.06
2:P:72:ASN:ND2	2:P:73:ASP:H	1.56	1.04

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	248/250 (99%)	235 (95%)	10 (4%)	3 (1%)	13	39
1	O	248/250 (99%)	236 (95%)	9 (4%)	3 (1%)	13	39
2	B	242/244 (99%)	214 (88%)	22 (9%)	6 (2%)	5	19
2	P	242/244 (99%)	215 (89%)	21 (9%)	6 (2%)	5	19
3	C	239/241 (99%)	220 (92%)	14 (6%)	5 (2%)	7	23
3	Q	239/241 (99%)	221 (92%)	13 (5%)	5 (2%)	7	23
4	D	240/242 (99%)	219 (91%)	16 (7%)	5 (2%)	7	23
4	R	240/242 (99%)	221 (92%)	14 (6%)	5 (2%)	7	23
5	E	231/233 (99%)	207 (90%)	20 (9%)	4 (2%)	9	29
5	S	231/233 (99%)	207 (90%)	20 (9%)	4 (2%)	9	29
6	F	242/244 (99%)	226 (93%)	13 (5%)	3 (1%)	13	39
6	T	242/244 (99%)	226 (93%)	13 (5%)	3 (1%)	13	39
7	G	241/243 (99%)	228 (95%)	12 (5%)	1 (0%)	34	66
7	U	241/243 (99%)	228 (95%)	12 (5%)	1 (0%)	34	66
8	H	220/222 (99%)	206 (94%)	10 (4%)	4 (2%)	8	28
8	V	220/222 (99%)	207 (94%)	9 (4%)	4 (2%)	8	28
9	I	202/204 (99%)	195 (96%)	7 (4%)	0	100	100
9	W	202/204 (99%)	195 (96%)	7 (4%)	0	100	100
10	J	196/198 (99%)	184 (94%)	9 (5%)	3 (2%)	10	33
10	X	196/198 (99%)	184 (94%)	9 (5%)	3 (2%)	10	33
11	K	210/212 (99%)	199 (95%)	10 (5%)	1 (0%)	29	61
11	Y	210/212 (99%)	199 (95%)	10 (5%)	1 (0%)	29	61
12	L	220/222 (99%)	206 (94%)	13 (6%)	1 (0%)	29	61
12	Z	220/222 (99%)	208 (94%)	10 (4%)	2 (1%)	17	46
13	M	231/233 (99%)	215 (93%)	14 (6%)	2 (1%)	17	46

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
13	a	231/233 (99%)	215 (93%)	14 (6%)	2 (1%)	17	46
14	N	194/196 (99%)	187 (96%)	7 (4%)	0	100	100
14	b	194/196 (99%)	187 (96%)	7 (4%)	0	100	100
15	1	1/4 (25%)	0	1 (100%)	0	100	100
15	2	1/4 (25%)	1 (100%)	0	0	100	100
15	3	1/4 (25%)	1 (100%)	0	0	100	100
15	4	1/4 (25%)	0	1 (100%)	0	100	100
15	5	1/4 (25%)	1 (100%)	0	0	100	100
15	6	1/4 (25%)	1 (100%)	0	0	100	100
All	All	6318/6392 (99%)	5894 (93%)	347 (6%)	77 (1%)	13	39

5 of 77 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	5	THR
1	A	169	LYS
2	B	186	MET
2	B	224	ASP
3	C	58	LEU

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	209/209 (100%)	204 (98%)	5 (2%)	49	81
1	O	209/209 (100%)	204 (98%)	5 (2%)	49	81
2	B	203/203 (100%)	186 (92%)	17 (8%)	11	31
2	P	203/203 (100%)	188 (93%)	15 (7%)	13	37
3	C	213/213 (100%)	201 (94%)	12 (6%)	21	51
3	Q	213/213 (100%)	201 (94%)	12 (6%)	21	51
4	D	198/198 (100%)	188 (95%)	10 (5%)	24	55

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
4	R	198/198 (100%)	188 (95%)	10 (5%)	24	55
5	E	192/192 (100%)	173 (90%)	19 (10%)	8	23
5	S	192/192 (100%)	173 (90%)	19 (10%)	8	23
6	F	201/201 (100%)	185 (92%)	16 (8%)	12	34
6	T	201/201 (100%)	185 (92%)	16 (8%)	12	34
7	G	207/207 (100%)	195 (94%)	12 (6%)	20	50
7	U	207/207 (100%)	195 (94%)	12 (6%)	20	50
8	H	181/181 (100%)	175 (97%)	6 (3%)	38	72
8	V	181/181 (100%)	175 (97%)	6 (3%)	38	72
9	I	172/172 (100%)	167 (97%)	5 (3%)	42	76
9	W	172/172 (100%)	168 (98%)	4 (2%)	50	82
10	J	175/175 (100%)	165 (94%)	10 (6%)	20	50
10	X	175/175 (100%)	165 (94%)	10 (6%)	20	50
11	K	169/169 (100%)	157 (93%)	12 (7%)	14	39
11	Y	169/169 (100%)	158 (94%)	11 (6%)	17	44
12	L	185/185 (100%)	170 (92%)	15 (8%)	11	33
12	Z	185/185 (100%)	171 (92%)	14 (8%)	13	36
13	M	199/199 (100%)	188 (94%)	11 (6%)	21	52
13	a	199/199 (100%)	187 (94%)	12 (6%)	19	48
14	N	162/162 (100%)	154 (95%)	8 (5%)	25	57
14	b	162/162 (100%)	153 (94%)	9 (6%)	21	51
15	1	1/1 (100%)	0	1 (100%)	0	0
15	2	1/1 (100%)	1 (100%)	0	100	100
15	3	1/1 (100%)	0	1 (100%)	0	0
15	4	1/1 (100%)	0	1 (100%)	0	0
15	5	1/1 (100%)	1 (100%)	0	100	100
15	6	1/1 (100%)	0	1 (100%)	0	0
All	All	5338/5338 (100%)	5021 (94%)	317 (6%)	19	49

5 of 317 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
6	T	216	ILE

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
12	Z	70	ASN
7	U	88	ASN
10	X	51	THR
13	a	162	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 202 such sidechains are listed below:

Mol	Chain	Res	Type
3	Q	126	GLN
6	T	127	ASN
14	b	145	ASN
3	Q	247	GLN
5	S	71	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

12 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
15	LYH	4	2	15	7,8,9	1.16	1 (14%)	4,8,10	1.03	0
15	LYH	1	2	15	7,8,9	1.14	0	4,8,10	0.98	0
15	LYH	3	2	15	7,8,9	1.17	0	4,8,10	1.45	1 (25%)
15	OJT	4	3	15,8	8,8,9	1.34	1 (12%)	7,9,11	1.02	0
15	OJT	1	3	15,8	8,8,9	1.35	2 (25%)	7,9,11	1.04	0
15	OJT	5	3	15,11	8,8,9	1.65	2 (25%)	7,9,11	1.16	1 (14%)
15	LYH	2	2	15	7,8,9	1.13	0	4,8,10	1.36	1 (25%)
15	LYH	5	2	15	7,8,9	1.13	1 (14%)	4,8,10	1.41	1 (25%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
15	OJT	2	3	15,11	8,8,9	1.51	1 (12%)	7,9,11	1.14	0
15	OJT	3	3	14,15	8,8,9	1.60	2 (25%)	7,9,11	1.06	0
15	OJT	6	3	14,15	8,8,9	1.70	2 (25%)	7,9,11	1.00	0
15	LYH	6	2	15	7,8,9	1.08	0	4,8,10	1.46	1 (25%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
15	LYH	4	2	15	-	1/3/7/9	-
15	LYH	1	2	15	-	1/3/7/9	-
15	LYH	3	2	15	-	1/3/7/9	-
15	OJT	4	3	15,8	-	5/7/8/9	-
15	OJT	1	3	15,8	-	5/7/8/9	-
15	OJT	5	3	15,11	-	2/7/8/9	-
15	LYH	2	2	15	-	1/3/7/9	-
15	LYH	5	2	15	-	1/3/7/9	-
15	OJT	2	3	15,11	-	2/7/8/9	-
15	OJT	3	3	14,15	-	2/7/8/9	-
15	OJT	6	3	14,15	-	2/7/8/9	-
15	LYH	6	2	15	-	1/3/7/9	-

The worst 5 of 12 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
15	6	3	OJT	C19-CA	3.69	1.58	1.53
15	5	3	OJT	C19-CA	3.60	1.58	1.53
15	3	3	OJT	C19-CA	3.42	1.58	1.53
15	2	3	OJT	C19-CA	3.34	1.58	1.53
15	4	3	OJT	C19-CA	2.81	1.57	1.53

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
15	5	2	LYH	CE-CD-CG	-2.34	107.75	113.06
15	6	2	LYH	CE-CD-CG	-2.33	107.77	113.06
15	3	2	LYH	CE-CD-CG	-2.29	107.85	113.06
15	2	2	LYH	CE-CD-CG	-2.20	108.06	113.06

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
15	5	3	0JT	C17-C16-C	2.01	118.02	112.29

There are no chirality outliers.

5 of 24 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
15	2	2	LYH	CG-CD-CE-NZ
15	3	2	LYH	CG-CD-CE-NZ
15	5	2	LYH	CG-CD-CE-NZ
15	6	2	LYH	CG-CD-CE-NZ
15	1	3	0JT	C-C16-C17-CA

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
15	3	2	LYH	1	0

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	250/250 (100%)	-0.45	3 (1%) 79 73	35, 55, 84, 105	0
1	O	250/250 (100%)	-0.43	2 (0%) 86 81	36, 55, 84, 105	0
2	B	244/244 (100%)	-0.25	7 (2%) 51 41	39, 56, 94, 118	0
2	P	244/244 (100%)	-0.27	8 (3%) 46 36	39, 57, 95, 118	0
3	C	241/241 (100%)	-0.21	7 (2%) 51 41	40, 59, 110, 129	0
3	Q	241/241 (100%)	-0.21	11 (4%) 32 22	42, 61, 110, 129	0
4	D	242/242 (100%)	-0.24	7 (2%) 51 41	39, 60, 92, 125	0
4	R	242/242 (100%)	-0.19	8 (3%) 46 36	41, 61, 93, 126	0
5	E	233/233 (100%)	-0.30	6 (2%) 56 46	42, 62, 87, 111	0
5	S	233/233 (100%)	-0.27	4 (1%) 70 63	42, 62, 87, 111	0
6	F	244/244 (100%)	-0.36	8 (3%) 46 36	39, 56, 92, 106	0
6	T	244/244 (100%)	-0.26	5 (2%) 65 56	39, 56, 93, 105	0
7	G	243/243 (100%)	-0.47	3 (1%) 79 73	37, 53, 80, 112	0
7	U	243/243 (100%)	-0.43	3 (1%) 79 73	36, 53, 79, 112	0
8	H	222/222 (100%)	-0.55	1 (0%) 91 88	31, 48, 68, 98	0
8	V	222/222 (100%)	-0.50	3 (1%) 75 70	33, 48, 67, 98	0
9	I	204/204 (100%)	-0.56	1 (0%) 91 88	35, 50, 67, 84	0
9	W	204/204 (100%)	-0.53	1 (0%) 91 88	37, 50, 68, 84	0
10	J	198/198 (100%)	-0.45	3 (1%) 73 68	37, 51, 68, 124	0
10	X	198/198 (100%)	-0.45	4 (2%) 65 56	38, 51, 70, 124	0
11	K	212/212 (100%)	-0.46	1 (0%) 91 88	34, 51, 75, 82	0
11	Y	212/212 (100%)	-0.50	1 (0%) 91 88	35, 51, 74, 82	0
12	L	222/222 (100%)	-0.59	2 (0%) 84 80	33, 50, 71, 96	0
12	Z	222/222 (100%)	-0.55	2 (0%) 84 80	33, 50, 71, 95	0

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
13	M	233/233 (100%)	-0.61	1 (0%) 92 91	35, 49, 64, 69	0
13	a	233/233 (100%)	-0.55	1 (0%) 92 91	34, 48, 64, 69	0
14	N	196/196 (100%)	-0.54	0 100 100	33, 45, 67, 77	0
14	b	196/196 (100%)	-0.53	1 (0%) 91 88	33, 46, 68, 77	0
15	1	1/4 (25%)	-0.31	0 100 100	50, 50, 50, 50	0
15	2	1/4 (25%)	-0.20	0 100 100	52, 52, 52, 52	0
15	3	1/4 (25%)	-0.23	0 100 100	57, 57, 57, 57	0
15	4	1/4 (25%)	-0.39	0 100 100	51, 51, 51, 51	0
15	5	1/4 (25%)	0.04	0 100 100	51, 51, 51, 51	0
15	6	1/4 (25%)	0.06	0 100 100	57, 57, 57, 57	0
All	All	6374/6392 (99%)	-0.41	104 (1%) 72 66	31, 53, 84, 129	0

The worst 5 of 104 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
4	D	126	GLY	9.6
3	C	55	THR	8.8
4	D	128	SER	8.5
4	R	128	SER	8.4
10	X	196	GLN	7.8

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
15	LYH	1	2	9/10	0.88	0.16	47,48,50,50	0
15	LYH	6	2	9/10	0.89	0.19	46,48,50,50	0
15	LYH	3	2	9/10	0.92	0.22	46,50,52,52	0
15	LYH	4	2	9/10	0.93	0.20	47,48,50,50	0
15	LYH	5	2	9/10	0.94	0.18	46,47,49,49	0
15	LYH	2	2	9/10	0.94	0.16	45,46,47,47	0
15	OJT	1	3	9/10	0.94	0.14	40,45,48,49	0
15	OJT	6	3	9/10	0.95	0.14	38,44,47,50	0
15	OJT	4	3	9/10	0.96	0.17	40,43,48,49	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
15	OJT	3	3	9/10	0.96	0.14	39,45,50,52	0
15	OJT	5	3	9/10	0.97	0.13	42,44,47,47	0
15	OJT	2	3	9/10	0.98	0.13	39,43,46,46	0

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.