

wwPDB X-ray Structure Validation Summary Report (i)

Feb 12, 2024 – 05:12 AM EST

PDB ID : 3GLF

Title : Crystal Structure of the Ecoli Clamp Loader Bound to Primer-Template DNA

Authors: Simonetta, K.R.; Kuriyan, J.

Deposited on : 2009-03-12

Resolution : 3.39 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity : 4.02b-467

Mogul : 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : 1.13

EDS : 2.36

buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

 $Refmac \quad : \quad 5.8.0158$

CCP4 : 7.0.044 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.36

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 3.39 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive $(\# \mathrm{Entries})$	$\begin{array}{c} {\rm Similar\ resolution} \\ (\#{\rm Entries,\ resolution\ range(\mathring{A})}) \end{array}$
R_{free}	130704	1691 (3.46-3.30)
Clashscore	141614	1762 (3.46-3.30)
Ramachandran outliers	138981	1732 (3.46-3.30)
Sidechain outliers	138945	1731 (3.46-3.30)
RSRZ outliers	127900	1635 (3.46-3.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Qualit	y of chain	
			15%		
1	A	343	48%	44%	5% •
			7%		
1	F	343	50%	42%	5% •
			4%		
2	В	395	54%	35%	• 8%
			3%		
2	С	395	52%	33%	7% • 8%
			3%		
2	D	395	50%	37%	• 8%



Continued from previous page...

Mol	Chain	Length	Quality	of chain
2	G	395	59%	34% • •
2	Н	395	50%	36% 6% 8%
2	I	395	49%	38% • 8%
3	Е	334	63%	34%
3	J	334	66%	30% •
4	K	15	13%	80% 7%
4	M	15	7% 13%	73% 7%
5	L	10	40%	60%
5	N	10	60%	40%



2 Entry composition (i)

There are 9 unique types of molecules in this entry. The entry contains 28758 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called DNA polymerase III subunit delta.

\mathbf{Mol}	Chain	Residues		\mathbf{At}	oms		ZeroOcc	AltConf	Trace	
1	A	333	Total 2650	C 1678	N 482	O 480	S 10	0	0	0
1	F	333	Total 2650	C 1678	N 482	O 480	S 10	0	0	0

• Molecule 2 is a protein called DNA polymerase III subunit tau.

Mol	Chain	Residues		At	oms			ZeroOcc	AltConf	Trace
2	В	364	Total	С	N	О	S	0	0	0
	D	304	2829	1779	511	523	16	O	0	
2	С	365	Total	С	N	О	S	0	0	0
2		300	2838	1784	513	525	16	U	U	
2	D	362	Total	С	N	О	S	0	0	0
	ש	302	2818	1770	510	522	16	U	0	
2	G	378	Total	С	N	О	S	0	0	0
2	G	310	2941	1852	529	543	17	U	U	
2	Н	365	Total	С	N	О	S	0	0	0
2	11	303	2838	1784	513	525	16	U	0	
2	Т	362	Total	С	N	О	S	0	0	0
	1	302	2818	1770	510	522	16	U	U	

There are 132 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
В	-21	MET	-	expression tag	UNP P06710
В	-20	GLY	-	expression tag	UNP P06710
В	-19	SER	-	expression tag	UNP P06710
В	-18	SER	-	expression tag	UNP P06710
В	-17	HIS	-	expression tag	UNP P06710
В	-16	HIS	-	expression tag	UNP P06710
В	-15	HIS	-	expression tag	UNP P06710
В	-14	HIS	-	expression tag	UNP P06710



 $Continued\ from\ previous\ page...$

Chain	Residue	Modelled Modelled	Actual	Comment	Reference
В	-13	HIS	-	expression tag	UNP P06710
В	-12	HIS	-	expression tag	UNP P06710
В	-11	SER	-	expression tag	UNP P06710
В	-10	SER	-	expression tag	UNP P06710
В	-9	GLY	-	expression tag	UNP P06710
В	-8	LEU	-	expression tag	UNP P06710
В	-7	GLU	-	expression tag	UNP P06710
В	-6	VAL	-	expression tag	UNP P06710
В	-5	LEU	-	expression tag	UNP P06710
В	-4	PHE	-	expression tag	UNP P06710
В	-3	GLN	-	expression tag	UNP P06710
В	-2	GLY	-	expression tag	UNP P06710
В	-1	PRO	-	expression tag	UNP P06710
В	0	HIS	-	expression tag	UNP P06710
С	-21	MET	-	expression tag	UNP P06710
С	-20	GLY	-	expression tag	UNP P06710
С	-19	SER	-	expression tag	UNP P06710
С	-18	SER	-	expression tag	UNP P06710
С	-17	HIS	-	expression tag	UNP P06710
С	-16	HIS	-	expression tag	UNP P06710
С	-15	HIS	-	expression tag	UNP P06710
С	-14	HIS	-	expression tag	UNP P06710
С	-13	HIS	-	expression tag	UNP P06710
С	-12	HIS	-	expression tag	UNP P06710
С	-11	SER	-	expression tag	UNP P06710
С	-10	SER	-	expression tag	UNP P06710
С	-9	GLY	-	expression tag	UNP P06710
С	-8	LEU	-	expression tag	UNP P06710
С	-7	GLU	-	expression tag	UNP P06710
С	-6	VAL	-	expression tag	UNP P06710
С	-5	LEU	-	expression tag	UNP P06710
С	-4	PHE	-	expression tag	UNP P06710
С	-3	GLN	-	expression tag	UNP P06710
С	-2	GLY	-	expression tag	UNP P06710
С	-1	PRO	_	expression tag	UNP P06710
С	0	HIS	-	expression tag	UNP P06710
D	-21	MET	-	expression tag	UNP P06710
D	-20	GLY	-	expression tag	UNP P06710
D	-19	SER	-	expression tag	UNP P06710
D	-18	SER	-	expression tag	UNP P06710
D	-17	HIS	-	expression tag	UNP P06710
D	-16	HIS	-	expression tag	UNP P06710



 $Continued\ from\ previous\ page...$

Chain	Residue	Modelled	Actual	Comment	Reference
D	-15	HIS	-	expression tag	UNP P06710
D	-14	HIS	_	expression tag	UNP P06710
D	-13	HIS	-	expression tag	UNP P06710
D	-12	HIS	_	expression tag	UNP P06710
D	-11	SER	-	expression tag	UNP P06710
D	-10	SER	_	expression tag	UNP P06710
D	-9	GLY	-	expression tag	UNP P06710
D	-8	LEU	-	expression tag	UNP P06710
D	-7	GLU	-	expression tag	UNP P06710
D	-6	VAL	-	expression tag	UNP P06710
D	-5	LEU	-	expression tag	UNP P06710
D	-4	PHE	-	expression tag	UNP P06710
D	-3	GLN	-	expression tag	UNP P06710
D	-2	GLY	_	expression tag	UNP P06710
D	-1	PRO	-	expression tag	UNP P06710
D	0	HIS	_	expression tag	UNP P06710
G	-21	MET	-	expression tag	UNP P06710
G	-20	GLY	-	expression tag	UNP P06710
G	-19	SER	_	expression tag	UNP P06710
G	-18	SER	-	expression tag	UNP P06710
G	-17	HIS	_	expression tag	UNP P06710
G	-16	HIS	-	expression tag	UNP P06710
G	-15	HIS	-	expression tag	UNP P06710
G	-14	HIS	-	expression tag	UNP P06710
G	-13	HIS	-	expression tag	UNP P06710
G	-12	HIS	-	expression tag	UNP P06710
G	-11	SER	-	expression tag	UNP P06710
G	-10	SER	-	expression tag	UNP P06710
G	-9	GLY	-	expression tag	UNP P06710
G	-8	LEU	-	expression tag	UNP P06710
G	-7	GLU	-	expression tag	UNP P06710
G	-6	VAL	-	expression tag	UNP P06710
G	-5	LEU	-	expression tag	UNP P06710
G	-4	PHE	-	expression tag	UNP P06710
G	-3	GLN	-	expression tag	UNP P06710
G	-2	GLY	-	expression tag	UNP P06710
G	-1	PRO	-	expression tag	UNP P06710
G	0	HIS	-	expression tag	UNP P06710
Н	-21	MET		expression tag	UNP P06710
Н	-20	GLY	-	expression tag	UNP P06710
Н	-19	SER	-	expression tag	UNP P06710
Н	-18	SER	-	expression tag	UNP P06710



 $Continued\ from\ previous\ page...$

Chain	Residue	Modelled Modelled	Actual	Comment	Reference
Н	-17	HIS	-	expression tag	UNP P06710
Н	-16	HIS	-	expression tag	UNP P06710
Н	-15	HIS	-	expression tag	UNP P06710
Н	-14	HIS	-	expression tag	UNP P06710
Н	-13	HIS	-	expression tag	UNP P06710
Н	-12	HIS	-	expression tag	UNP P06710
Н	-11	SER	-	expression tag	UNP P06710
Н	-10	SER	-	expression tag	UNP P06710
Н	-9	GLY	-	expression tag	UNP P06710
Н	-8	LEU	-	expression tag	UNP P06710
Н	-7	GLU	-	expression tag	UNP P06710
Н	-6	VAL	-	expression tag	UNP P06710
Н	-5	LEU	-	expression tag	UNP P06710
Н	-4	PHE	-	expression tag	UNP P06710
Н	-3	GLN	-	expression tag	UNP P06710
Н	-2	GLY	-	expression tag	UNP P06710
Н	-1	PRO	-	expression tag	UNP P06710
Н	0	HIS	-	expression tag	UNP P06710
I	-21	MET	-	expression tag	UNP P06710
I	-20	GLY	-	expression tag	UNP P06710
I	-19	SER	-	expression tag	UNP P06710
I	-18	SER	-	expression tag	UNP P06710
I	-17	HIS	_	expression tag	UNP P06710
I	-16	HIS	-	expression tag	UNP P06710
I	-15	HIS	-	expression tag	UNP P06710
I	-14	HIS	-	expression tag	UNP P06710
I	-13	HIS	-	expression tag	UNP P06710
I	-12	HIS	-	expression tag	UNP P06710
I	-11	SER	-	expression tag	UNP P06710
I	-10	SER	-	expression tag	UNP P06710
I	-9	GLY	-	expression tag	UNP P06710
I	-8	LEU	-	expression tag	UNP P06710
I	-7	GLU	-	expression tag	UNP P06710
I	-6	VAL	-	expression tag	UNP P06710
I	-5	LEU	-	expression tag	UNP P06710
I	-4	PHE	-	expression tag	UNP P06710
I	-3	GLN	-	expression tag	UNP P06710
I	-2	GLY	-	expression tag	UNP P06710
I	-1	PRO	-	expression tag	UNP P06710
I	0	HIS	-	expression tag	UNP P06710

 \bullet Molecule 3 is a protein called DNA polymerase III subunit delta'.



Mol	Chain	Residues		At	oms		ZeroOcc	AltConf	Trace	
3	Е	334	Total 2601	C 1655	N 468	O 465	S 13	0	0	0
3	J	334	Total 2601	C 1655	N 468	O 465	S 13	0	0	0

• Molecule 4 is a DNA chain called DNA (5'-D(*TP*TP*TP*TP*TP*TP*AP*TP*AP*GP*GP*CP*CP*AP*G)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	I/	1.4	Total	С	N	О	Р	0	0	0
4	4 K	14	287	138	48	87	14	U	U	U
1	М	1.4	Total	С	N	О	Р	0	0	0
4	4 M	14	287	138	48	87	14		U	U

 \bullet Molecule 5 is a DNA chain called DNA (5'-D(*CP*TP*GP*GP*CP*TP*AP*TP*A)-3').


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
5	Т	10	Total	С	N	О	Р	0	0	0
9 L	10	200	97	35	59	9	U	U		
5	N	10	Total	С	N	О	Р	0	0	0
5 N	10	200	97	35	59	9	U			

• Molecule 6 is ADENOSINE-5'-DIPHOSPHATE (three-letter code: ADP) (formula: $C_{10}H_{15}N_5O_{10}P_2$).



Mol	Chain	Residues		Ato	oms			ZeroOcc	AltConf
6	В	1	Total	С	N	О	Р	0	0
0	Ъ	1	27	10	5	10	2	0	0
6	С	1	Total	С	N	О	Р	0	0
0		1	27	10	5	10	2	0	0
6	D	1	Total	С	N	О	Р	0	0
0	D	1	27	10	5	10	2	U	
6	G	1	Total	С	N	О	Р	0	0
0	G	1	27	10	5	10	2	U	0
6	Н	1	Total	С	N	О	Р	0	0
0	11	1	27	10	5	10	2	U	0
6	Т	1	Total	С	N	О	Р	0	0
	1	1	27	10	5	10	2	U	

 \bullet Molecule 7 is BERYLLIUM TRIFLUORIDE ION (three-letter code: BEF) (formula: BeF3).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	В	1	Total Be F 4 1 3	0	0
7	С	1	Total Be F 4 1 3	0	0
7	D	1	Total Be F 4 1 3	0	0
7	G	1	Total Be F 4 1 3	0	0
7	Н	1	Total Be F 4 1 3	0	0
7	I	1	Total Be F 4 1 3	0	0



• Molecule 8 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
8	В	1	Total Mg 1 1	0	0
8	С	1	Total Mg 1 1	0	0
8	D	1	Total Mg 1 1	0	0
8	G	1	Total Mg 1 1	0	0
8	Н	1	Total Mg 1 1	0	0
8	I	1	Total Mg 1 1	0	0

• Molecule 9 is ZINC ION (three-letter code: ZN) (formula: Zn).


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
9	В	1	Total Zn 1 1	0	0
9	С	1	Total Zn 1 1	0	0
9	D	1	Total Zn 1 1	0	0
9	E	1	Total Zn 1 1	0	0
9	G	1	Total Zn 1 1	0	0
9	Н	1	Total Zn 1 1	0	0
9	I	1	Total Zn 1 1	0	0
9	J	1	Total Zn 1 1	0	0




3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: DNA polymerase III subunit delta




• Molecule 1: DNA polymerase III subunit delta








• Molecule 2: DNA polymerase III subunit tau




• Molecule 2: DNA polymerase III subunit tau




Chain D: 37%







• Molecule 2: DNA polymerase III subunit tau




• Molecule 2: DNA polymerase III subunit tau









• Molecule 2: DNA polymerase III subunit tau




• Molecule 3: DNA polymerase III subunit delta'



• Molecule 3: DNA polymerase III subunit delta'







4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants	100.26Å 219.95Å 273.15Å	Donositon
a, b, c, α , β , γ	90.00° 90.00° 90.00°	Depositor
Resolution (Å)	49.97 - 3.39	Depositor
Resolution (A)	49.97 - 3.39	EDS
% Data completeness	97.5 (49.97-3.39)	Depositor
(in resolution range)	97.5 (49.97-3.39)	EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	2.52 (at 3.40Å)	Xtriage
Refinement program	PHENIX (phenix.refine)	Depositor
D D	0.221 , 0.260	Depositor
R, R_{free}	0.209 , 0.251	DCC
R_{free} test set	4166 reflections (5.04%)	wwPDB-VP
Wilson B-factor (Å ²)	92.3	Xtriage
Anisotropy	0.197	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.30 , 47.5	EDS
L-test for twinning ²	$ < L >=0.52, < L^2>=0.36$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	28758	wwPDB-VP
Average B, all atoms (Å ²)	107.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 2.15% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of <|L|>, $<L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: BEF, ADP, ZN, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Во	ond lengths	В	Bond angles
IVIOI	Chain	RMSZ	# Z > 5	RMSZ	# Z >5
1	A	0.72	0/2697	0.78	3/3664 (0.1%)
1	F	0.73	0/2697	0.78	3/3664 (0.1%)
2	В	0.72	0/2876	0.81	0/3900
2	С	0.72	$1/2885 \ (0.0\%)$	0.86	$6/3912 \ (0.2\%)$
2	D	0.77	$1/2863 \ (0.0\%)$	0.89	0/3879
2	G	0.75	2/2992~(0.1%)	0.83	1/4057 (0.0%)
2	Н	0.86	$4/2885 \ (0.1\%)$	0.92	4/3912 (0.1%)
2	I	0.92	$4/2863 \; (0.1\%)$	0.96	2/3879 (0.1%)
3	Е	0.89	$2/2666 \ (0.1\%)$	0.91	5/3639 (0.1%)
3	J	0.89	1/2666~(0.0%)	0.91	2/3639 (0.1%)
4	K	1.97	$6/320 \ (1.9\%)$	2.68	$32/492 \ (6.5\%)$
4	M	1.96	5/320 (1.6%)	2.58	27/492 (5.5%)
5	L	1.76	3/223 (1.3%)	2.80	28/342 (8.2%)
5	N	1.76	2/223~(0.9%)	2.75	25/342 (7.3%)
All	All	0.87	31/29176 (0.1%)	1.01	138/39813 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a maintain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	В	0	1
2	С	0	1
2	G	0	1
2	Н	0	1
2	I	0	1
All	All	0	5

The worst 5 of 31 bond length outliers are listed below:



Mol	Chain	Res	Type	Atoms	Z	Observed(A)	$\operatorname{Ideal}(ext{\AA})$
4	M	11	DG	C3'-O3'	-8.40	1.33	1.44
2	Н	127	GLU	CG-CD	8.31	1.64	1.51
4	M	9	DA	C3'-O3'	-8.17	1.33	1.44
5	N	6	DC	C3'-O3'	-7.52	1.34	1.44
2	G	127	GLU	CG-CD	7.35	1.62	1.51

The worst 5 of 138 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$Observed(^o)$	$\operatorname{Ideal}({}^{o})$
4	M	5	DT	O4'-C1'-N1	-11.71	99.80	108.00
4	K	5	DT	O4'-C1'-N1	-11.64	99.85	108.00
5	L	5	DC	O4'-C4'-C3'	-11.36	99.19	106.00
5	N	2	DT	N3-C4-O4	10.52	126.21	119.90
5	N	5	DC	O4'-C4'-C3'	-10.37	99.78	106.00

There are no chirality outliers.

All (5) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	В	246	ASP	Peptide
2	С	245	ASP	Peptide
2	G	246	ASP	Peptide
2	Н	245	ASP	Peptide
2	I	157	THR	Peptide

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2650	0	2703	168	0
1	F	2650	0	2703	158	0
2	В	2829	0	2879	153	1
2	С	2838	0	2886	165	1
2	D	2818	0	2863	173	0
2	G	2941	0	2987	139	0
2	Н	2838	0	2886	160	2
2	I	2818	0	2863	165	1
3	Е	2601	0	2603	108	1



Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	J	2601	0	2603	119	0
4	K	287	0	161	17	0
4	M	287	0	161	14	0
5	L	200	0	115	5	0
5	N	200	0	115	4	0
6	В	27	0	12	3	0
6	С	27	0	12	3	0
6	D	27	0	12	3	0
6	G	27	0	12	2	0
6	Н	27	0	12	4	0
6	I	27	0	12	2	0
7	В	4	0	0	1	0
7	С	4	0	0	0	0
7	D	4	0	0	0	0
7	G	4	0	0	1	0
7	Н	4	0	0	0	0
7	I	4	0	0	1	0
8	В	1	0	0	0	0
8	С	1	0	0	0	0
8	D	1	0	0	0	0
8	G	1	0	0	0	0
8	Н	1	0	0	0	0
8	I	1	0	0	0	0
9	В	1	0	0	0	0
9	С	1	0	0	0	0
9	D	1	0	0	0	0
9	Е	1	0	0	0	0
9	G	1	0	0	0	0
9	Н	1	0	0	0	0
9	I	1	0	0	0	0
9	J	1	0	0	0	0
All	All	28758	0	28600	1426	3

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 25.

The worst 5 of 1426 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	$\begin{array}{c} {\rm Interatomic} \\ {\rm distance} \ ({\rm \AA}) \end{array}$	$egin{aligned} ext{Clash} \ ext{overlap } (ext{Å}) \end{aligned}$
2:H:351:MET:CE	2:I:290:HIS:HA	1.75	1.15
1:F:95:GLU:HA	1:F:98:LEU:HD12	1.33	1.08



Continued from previous page...

Atom-1	Atom-2	$egin{array}{c} ext{Interatomic} \ ext{distance} & (ext{Å}) \end{array}$	$egin{aligned} ext{Clash} \ ext{overlap } (ext{Å}) \end{aligned}$
1:A:95:GLU:HA	1:A:98:LEU:HD12	1.35	1.07
2:H:351:MET:HE1	2:I:290:HIS:HA	1.13	1.06
2:I:216:ASP:OD1	3:J:157:SER:HB2	1.57	1.03

All (3) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	$\begin{array}{c} \text{Interatomic} \\ \text{distance (Å)} \end{array}$	Clash overlap (Å)
2:B:117:ARG:NH2	2:I:117:ARG:NE[2_555]	1.99	0.21
3:E:178:ARG:NH2	2:H:195:HIS:O[4_545]	2.02	0.18
2:C:117:ARG:NE	2:H:117:ARG:NH2[2_555]	2.03	0.17

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	331/343 (96%)	292 (88%)	34 (10%)	5 (2%)	10 39
1	F	331/343 (96%)	298 (90%)	29 (9%)	4 (1%)	13 44
2	В	362/395~(92%)	323 (89%)	36 (10%)	3 (1%)	19 53
2	С	363/395~(92%)	326 (90%)	31 (8%)	6 (2%)	9 36
2	D	360/395~(91%)	317 (88%)	34 (9%)	9 (2%)	5 29
2	G	376/395~(95%)	340 (90%)	33 (9%)	3 (1%)	19 53
2	Н	363/395~(92%)	320 (88%)	37 (10%)	6 (2%)	9 36
2	I	360/395~(91%)	318 (88%)	33 (9%)	9 (2%)	5 29
3	E	332/334~(99%)	300 (90%)	24 (7%)	8 (2%)	6 30
3	J	332/334~(99%)	298 (90%)	28 (8%)	6 (2%)	8 35
All	All	3510/3724 (94%)	3132 (89%)	319 (9%)	59 (2%)	9 36

5 of 59 Ramachandran outliers are listed below:



Mol	Chain	Res	Type
1	A	159	ASN
2	В	307	ALA
2	С	297	LEU
2	D	307	ALA
3	Е	153	ALA

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	283/291 (97%)	256 (90%)	27 (10%)	8 30
1	F	283/291 (97%)	256 (90%)	27 (10%)	8 30
2	В	301/329~(92%)	282 (94%)	19 (6%)	18 49
2	C	302/329~(92%)	267 (88%)	35 (12%)	5 21
2	D	299/329~(91%)	280 (94%)	19 (6%)	17 48
2	G	313/329 (95%)	296 (95%)	17 (5%)	22 53
2	Н	302/329~(92%)	266 (88%)	36 (12%)	5 20
2	I	299/329 (91%)	283 (95%)	16 (5%)	22 53
3	E	270/270 (100%)	254 (94%)	16 (6%)	19 51
3	J	270/270 (100%)	252 (93%)	18 (7%)	16 47
All	All	$2922/3096 \ (94\%)$	2692 (92%)	230 (8%)	12 39

5 of 230 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	Е	282	SER
3	J	188	LEU
2	G	37	ILE
3	J	160	ARG
2	I	52	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 56 such sidechains are listed below:



Mol	Chain	Res	Type
1	F	45	GLN
3	J	267	ASN
2	G	260	ASN
3	J	246	HIS
3	J	51	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

Of 26 ligands modelled in this entry, 14 are monoatomic - leaving 12 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Mol Type Chain Res Lin		Link	Вс	ond leng	ths	Bond angles			
IVIOI	Type	Chain	nes	LIIIK	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
6	ADP	Н	408	7,8	24,29,29	1.24	3 (12%)	29,45,45	1.45	4 (13%)
7	BEF	Н	409	6	0,3,3	-	-	-		
6	ADP	В	400	7,8	24,29,29	1.32	2 (8%)	29,45,45	1.54	6 (20%)
7	BEF	I	411	6	0,3,3	-	=	-		
7	BEF	С	403	6	0,3,3	-	-	-		
6	ADP	I	410	7,8	24,29,29	1.50	4 (16%)	29,45,45	1.57	6 (20%)
7	BEF	G	407	6	0,3,3	-	-	-		
6	ADP	D	404	7,8	24,29,29	1.22	2 (8%)	29,45,45	1.66	9 (31%)
6	ADP	С	402	7,8	24,29,29	1.20	2 (8%)	29,45,45	1.65	5 (17%)



Mol	Type	Chain	hain Res 1		Во	ond leng	ths	В	ond ang	les
MIOI	Туре	Chain	nes	Link	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
6	ADP	G	406	7,8	24,29,29	1.21	3 (12%)	29,45,45	1.49	7 (24%)
7	BEF	В	401	6	0,3,3	-	-	-		
7	BEF	D	405	6	0,3,3	-	-	-		

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	ADP	Н	408	7,8	-	5/12/32/32	0/3/3/3
6	ADP	В	400	7,8	-	2/12/32/32	0/3/3/3
6	ADP	I	410	7,8	-	1/12/32/32	0/3/3/3
6	ADP	D	404	7,8	-	2/12/32/32	0/3/3/3
6	ADP	С	402	7,8	-	6/12/32/32	0/3/3/3
6	ADP	G	406	7,8	-	0/12/32/32	0/3/3/3

The worst 5 of 16 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	Observed(A)	$\operatorname{Ideal}(ext{\AA})$
6	I	410	ADP	C2'-C1'	-3.87	1.47	1.53
6	С	402	ADP	C2'-C1'	-3.34	1.48	1.53
6	В	400	ADP	C2'-C1'	-3.08	1.49	1.53
6	D	404	ADP	C2'-C1'	-2.87	1.49	1.53
6	В	400	ADP	C5-C4	2.85	1.48	1.40

The worst 5 of 37 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$\mathbf{Observed}(^{o})$	$\operatorname{Ideal}(^{o})$
6	С	402	ADP	PA-O3A-PB	-3.99	119.12	132.83
6	D	404	ADP	PA-O3A-PB	-3.59	120.50	132.83
6	С	402	ADP	O2B-PB-O1B	3.55	124.59	110.68
6	I	410	ADP	O2B-PB-O1B	3.52	124.48	110.68
6	Н	408	ADP	PA-O3A-PB	-3.37	121.26	132.83

There are no chirality outliers.

5 of 16 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
6	С	402	ADP	PA-O3A-PB-O3B



Continued from previous page...

Mol	Chain	Res	Type	Atoms
6	С	402	ADP	C5'-O5'-PA-O1A
6	С	402	ADP	C5'-O5'-PA-O2A
6	Н	408	ADP	PA-O3A-PB-O3B
6	Н	408	ADP	C5'-O5'-PA-O2A


There are no ring outliers.


9 monomers are involved in 20 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	Н	408	ADP	4	0
6	В	400	ADP	3	0
7	I	411	BEF	1	0
6	I	410	ADP	2	0
7	G	407	BEF	1	0
6	D	404	ADP	3	0
6	С	402	ADP	3	0
6	G	406	ADP	2	0
7	В	401	BEF	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



















5.7 Other polymers (i)

There are no such residues in this entry.



5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<rsrz></rsrz>	# RSRZ > 2	$OWAB(\AA^2)$	Q<0.9
1	A	333/343~(97%)	0.89	53 (15%) 1 2	73, 127, 209, 244	0
1	F	333/343 (97%)	0.40	25 (7%) 14 17	67, 127, 197, 222	0
2	В	$364/395 \ (92\%)$	0.29	16 (4%) 34 38	76, 119, 164, 189	0
2	С	365/395~(92%)	0.23	12 (3%) 46 50	68, 114, 180, 208	0
2	D	362/395 (91%)	0.21	10 (2%) 53 57	69, 106, 156, 198	0
2	G	378/395~(95%)	0.08	7 (1%) 66 71	76, 101, 146, 182	0
2	Н	365/395~(92%)	-0.05	2 (0%) 91 93	58, 88, 125, 165	0
2	I	362/395~(91%)	-0.02	1 (0%) 94 96	56, 80, 125, 184	0
3	E	$334/334\ (100\%)$	0.06	4 (1%) 79 83	67, 86, 134, 175	0
3	J	$334/334\ (100\%)$	0.02	3 (0%) 84 88	63, 84, 133, 172	0
4	K	14/15~(93%)	0.06	0 100 100	79, 87, 169, 171	0
4	M	14/15~(93%)	-0.16	0 100 100	76, 84, 174, 181	0
5	L	10/10 (100%)	-0.39	0 100 100	85, 88, 95, 105	0
5	N	10/10 (100%)	-0.40	0 100 100	81, 86, 98, 99	0
All	All	3578/3774~(94%)	0.20	133 (3%) 41 45	56, 101, 175, 244	0

The worst 5 of 133 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	53	PHE	10.9
1	A	55	ILE	9.4
1	A	61	TRP	8.7
1	F	61	TRP	8.3
1	A	62	ASN	7.1



6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.


6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\operatorname{B-factors}(\mathring{\mathbf{A}}^2)$	Q<0.9
8	MG	D	414	1/1	0.84	0.39	75,75,75,75	0
9	ZN	J	425	1/1	0.87	0.09	146,146,146,146	0
8	MG	G	415	1/1	0.92	0.35	79,79,79,79	0
8	MG	В	412	1/1	0.92	0.38	79,79,79,79	0
8	MG	Н	416	1/1	0.93	0.26	57,57,57,57	0
6	ADP	D	404	27/27	0.95	0.21	80,89,95,96	0
7	BEF	G	407	4/4	0.95	0.23	79,79,80,80	0
6	ADP	В	400	27/27	0.95	0.22	86,94,99,100	0
8	MG	С	413	1/1	0.95	0.29	77,77,77,77	0
6	ADP	G	406	27/27	0.96	0.23	81,87,92,94	0
6	ADP	С	402	27/27	0.96	0.21	78,86,91,93	0
7	BEF	В	401	4/4	0.97	0.28	79,80,80,81	0
7	BEF	С	403	4/4	0.97	0.26	77,78,78,79	0
7	BEF	D	405	4/4	0.97	0.23	85,85,86,87	0
6	ADP	Н	408	27/27	0.97	0.20	60,62,67,67	0
7	BEF	Н	409	4/4	0.97	0.22	57,57,57,58	0
8	MG	I	417	1/1	0.97	0.39	58,58,58,58	0
9	ZN	D	420	1/1	0.97	0.07	133,133,133,133	0
7	BEF	I	411	4/4	0.97	0.29	62,63,63,64	0
6	ADP	I	410	27/27	0.98	0.23	62,68,72,73	0
9	ZN	Е	421	1/1	0.98	0.06	161,161,161,161	0
9	ZN	Н	423	1/1	0.98	0.10	112,112,112,112	0
9	ZN	В	418	1/1	0.98	0.07	144,144,144,144	0
9	ZN	С	419	1/1	0.99	0.06	130,130,130,130	0
9	ZN	I	424	1/1	0.99	0.10	99,99,99,99	0
9	ZN	G	422	1/1	0.99	0.12	160,160,160,160	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.


























6.5 Other polymers (i)

There are no such residues in this entry.

