



# Full wwPDB X-ray Structure Validation Report ⓘ

Oct 31, 2023 – 05:26 PM JST

PDB ID : 5GMY  
Title : Crystal structure of the Archaeoglobus fulgidus oligosaccharyltransferase (O29867\_ARCFU) tethered with an acceptor peptide containing the NVT sequon via a disulfide bond  
Authors : Matsumoto, S.; Kohda, D.  
Deposited on : 2016-07-18  
Resolution : 3.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.13  
EDS : 2.36  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 3.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1659 (3.60-3.40)
Clashscore	141614	1036 (3.58-3.42)
Ramachandran outliers	138981	1005 (3.58-3.42)
Sidechain outliers	138945	1006 (3.58-3.42)
RSRZ outliers	127900	1559 (3.60-3.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	875	 6% (Poor fit)   81% (0-1 outliers)   16% (2-3 outliers)   . (Not modelled)
2	B	7	 71% (0-1 outliers)   29% (2-3 outliers)

## 2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 6887 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Transmembrane oligosaccharyl transferase, putative.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	848	6830	4540	1084	1189	17	0	0	0

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	617	CYS	GLY	engineered mutation	UNP O29867
A	869	GLU	-	expression tag	UNP O29867
A	870	LEU	-	expression tag	UNP O29867
A	871	ALA	-	expression tag	UNP O29867
A	872	LEU	-	expression tag	UNP O29867
A	873	VAL	-	expression tag	UNP O29867
A	874	PRO	-	expression tag	UNP O29867
A	875	ARG	-	expression tag	UNP O29867

- Molecule 2 is a protein called acceptor peptide, ARG-TYR-ASN-VAL-THR-ALA-CYS.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	7	56	34	11	10	1	0	0	0


- Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Mg	0	0
			1	1		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Transmembrane oligosaccharyl transferase, putative



- Molecule 2: acceptor peptide, ARG-TYR-ASN-VAL-THR-ALA-CYS



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	121.55Å 121.55Å 181.34Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	40.10 – 3.50 46.62 – 3.50	Depositor EDS
% Data completeness (in resolution range)	98.4 (40.10-3.50) 98.9 (46.62-3.50)	Depositor EDS
$R_{merge}$	0.11	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.52 (at 3.48Å)	Xtrriage
Refinement program	PHENIX 1.18.2_3874	Depositor
R, $R_{free}$	0.226 , 0.274 0.226 , 0.274	Depositor DCC
$R_{free}$ test set	894 reflections (5.10%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	142.7	Xtrriage
Anisotropy	0.589	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.28 , 141.1	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.46$ , $\langle L^2 \rangle = 0.29$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	6887	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	202.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.50% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.25	0/7055	0.41	0/9627
2	B	0.25	0/56	0.45	0/75
All	All	0.25	0/7111	0.41	0/9702

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	6830	0	6721	87	0
2	B	56	0	56	2	0
3	A	1	0	0	0	0
All	All	6887	0	6777	87	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (87) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:458:ASN:HB2	1:A:463:ARG:HA	1.56	0.87
1:A:193:LEU:HD23	1:A:195:ALA:H	1.40	0.87
1:A:297:ILE:HD11	1:A:323:LEU:HD13	1.73	0.70
1:A:681:LYS:HA	1:A:685:PHE:HD2	1.57	0.69
1:A:799:ILE:HA	1:A:836:PRO:HA	1.76	0.68
1:A:356:LEU:HD12	1:A:361:VAL:HG21	1.76	0.67
1:A:131:VAL:HG12	1:A:406:PRO:HB2	1.77	0.66
1:A:637:TYR:HA	1:A:670:PRO:HA	1.78	0.65
1:A:74:TYR:HE2	1:A:604:ASN:HB3	1.64	0.62
1:A:183:ARG:O	1:A:197:GLN:NE2	2.32	0.62
1:A:394:TYR:HD1	1:A:441:LEU:HD21	1.65	0.61
1:A:837:ILE:HG13	1:A:838:THR:HG23	1.83	0.61
1:A:75:PRO:HG2	1:A:603:LEU:HD22	1.84	0.60
1:A:126:LEU:HD11	1:A:179:LEU:HD12	1.83	0.60
1:A:566:ILE:HG21	1:A:603:LEU:HD13	1.84	0.59
1:A:551:TRP:HE3	1:A:569:PRO:HA	1.67	0.59
1:A:827:SER:HB2	1:A:858:VAL:HG21	1.82	0.59
1:A:360:GLU:HG3	2:B:3:ASN:H	1.68	0.58
1:A:5:GLU:HA	1:A:9:LYS:HD3	1.85	0.58
1:A:131:VAL:HG21	1:A:410:LEU:HD22	1.86	0.57
1:A:683:HIS:O	1:A:772:LYS:NZ	2.38	0.57
1:A:32:PRO:HG3	1:A:156:ILE:HG23	1.87	0.56
1:A:554:GLY:HA3	1:A:568:ASN:HA	1.88	0.56
1:A:360:GLU:O	1:A:425:ASN:HB3	2.06	0.56
1:A:124:VAL:HG21	1:A:144:ILE:HG12	1.88	0.55
1:A:608:VAL:HG22	1:A:774:PHE:HB2	1.90	0.54
1:A:258:ALA:HB2	1:A:284:LEU:HB3	1.90	0.53
1:A:551:TRP:CE3	1:A:569:PRO:HA	2.44	0.53
1:A:616:THR:OG1	1:A:617:CYS:N	2.38	0.53
1:A:178:LEU:O	1:A:182:ASN:ND2	2.41	0.53
1:A:211:TYR:HH	1:A:430:TYR:HH	1.57	0.52
1:A:125:TYR:HB2	1:A:141:ALA:HB2	1.90	0.52
1:A:124:VAL:O	1:A:128:THR:HG22	2.09	0.51
1:A:109:LEU:HB3	1:A:160:ASN:HD22	1.76	0.51
1:A:545:TYR:HB2	1:A:606:LYS:HB2	1.93	0.51
1:A:71:PHE:HD2	1:A:558:GLU:HG3	1.75	0.51
1:A:786:VAL:HG22	1:A:787:SER:H	1.75	0.51
1:A:698:GLU:OE1	1:A:805:ARG:NH2	2.44	0.51
1:A:74:TYR:O	1:A:540:TYR:OH	2.22	0.51
1:A:458:ASN:OD1	1:A:459:ALA:N	2.42	0.51
1:A:582:PRO:HG3	1:A:602:LYS:HG2	1.93	0.51
1:A:128:THR:HG23	1:A:137:ALA:HA	1.92	0.50

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:842:ILE:HD11	1:A:849:LYS:HB2	1.94	0.50
1:A:234:VAL:HA	1:A:317:PRO:HB3	1.93	0.49
1:A:550:TRP:HE1	2:B:5:THR:HG21	1.77	0.49
1:A:694:ARG:HD3	1:A:825:PRO:O	2.12	0.49
1:A:362:TYR:CE2	1:A:425:ASN:ND2	2.80	0.49
1:A:60:HIS:ND1	1:A:525:LEU:O	2.46	0.48
1:A:226:LEU:HA	1:A:229:MET:HE3	1.96	0.48
1:A:551:TRP:HB3	1:A:567:ALA:HB1	1.95	0.48
1:A:799:ILE:HG12	1:A:836:PRO:HG3	1.96	0.47
1:A:399:PHE:CE2	1:A:408:MET:HB2	2.50	0.47
1:A:714:LEU:HD13	1:A:767:PRO:HG2	1.96	0.47
1:A:320:VAL:O	1:A:324:THR:OG1	2.24	0.46
1:A:484:PRO:HA	1:A:487:ILE:HG22	1.98	0.46
1:A:194:THR:HG22	1:A:194:THR:O	2.16	0.46
1:A:39:TRP:HD1	1:A:40:THR:HG22	1.81	0.46
1:A:233:PHE:HB3	1:A:320:VAL:HG21	1.97	0.46
1:A:548:ILE:HG22	1:A:608:VAL:HG12	1.97	0.45
1:A:109:LEU:HD22	1:A:160:ASN:ND2	2.32	0.45
1:A:765:ILE:O	1:A:765:ILE:HG13	2.15	0.45
1:A:582:PRO:HA	1:A:599:VAL:HG22	1.98	0.44
1:A:547:VAL:O	1:A:565:PRO:HA	2.18	0.44
1:A:576:ASN:HA	1:A:626:ALA:HA	1.99	0.44
1:A:135:ARG:HD3	1:A:453:TYR:CZ	2.53	0.44
1:A:364:PHE:CZ	1:A:376:ASN:HB3	2.53	0.43
1:A:73:TYR:HD2	1:A:77:GLY:H	1.66	0.43
1:A:286:GLY:O	1:A:290:ILE:HG13	2.19	0.43
1:A:246:LEU:HD11	1:A:405:PHE:CZ	2.54	0.43
1:A:231:PHE:HA	1:A:234:VAL:HG22	2.00	0.43
1:A:720:VAL:HG21	1:A:767:PRO:HD3	2.01	0.42
1:A:53:ARG:NH2	1:A:528:TYR:OH	2.50	0.42
1:A:646:THR:HG23	1:A:648:THR:HG23	2.01	0.42
1:A:143:LEU:HD22	1:A:384:LEU:HD21	2.01	0.42
1:A:690:LEU:HD22	1:A:693:TYR:HD2	1.85	0.42
1:A:786:VAL:HG13	1:A:788:ALA:H	1.85	0.41
1:A:189:LEU:C	1:A:191:HIS:H	2.24	0.41
1:A:62:PHE:CG	1:A:63:PRO:HA	2.55	0.41
1:A:674:TYR:O	1:A:677:THR:OG1	2.33	0.41
1:A:191:HIS:HA	1:A:194:THR:OG1	2.21	0.41
1:A:577:LYS:NZ	1:A:636:TYR:OH	2.54	0.41
1:A:510:MET:HG3	1:A:514:THR:HG21	2.03	0.41
1:A:555:HIS:CD2	1:A:569:PRO:HD2	2.56	0.40

*Continued on next page...*



Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:73:TYR:CE2	1:A:76:TYR:HB2	2.55	0.40
1:A:74:TYR:HB3	1:A:566:ILE:HG13	2.04	0.40
1:A:540:TYR:HA	1:A:541:PRO:HD3	1.90	0.40
1:A:685:PHE:HD1	1:A:763:VAL:HG11	1.86	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	844/875 (96%)	793 (94%)	50 (6%)	1 (0%)	51	84
2	B	5/7 (71%)	2 (40%)	3 (60%)	0	100	100
All	All	849/882 (96%)	795 (94%)	53 (6%)	1 (0%)	51	84

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	148	PRO

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	704/728 (97%)	703 (100%)	1 (0%)	93	98

Continued on next page...

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	B	6/6 (100%)	6 (100%)	0	100	100
All	All	710/734 (97%)	709 (100%)	1 (0%)	93	98

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	53	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	160	ASN
1	A	182	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

Of 1 ligands modelled in this entry, 1 is monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	848/875 (96%)	0.24	56 (6%) 18 17	129, 194, 276, 439	0
2	B	7/7 (100%)	3.34	5 (71%) 0 0	198, 213, 234, 273	0
All	All	855/882 (96%)	0.27	61 (7%) 16 15	129, 194, 276, 439	0

All (61) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	792	GLU	8.1
2	B	1	ARG	7.3
2	B	3	ASN	5.8
1	A	643	TYR	5.1
1	A	864	ILE	5.1
1	A	793	VAL	4.6
1	A	842	ILE	4.1
1	A	765	ILE	4.1
1	A	5	GLU	3.5
1	A	868	LEU	3.4
1	A	663	ILE	3.4
1	A	782	VAL	3.3
2	B	4	VAL	3.3
1	A	814	GLU	3.3
1	A	813	VAL	3.2
1	A	865	GLN	3.2
2	B	7	CYS	3.2
1	A	6	SER	3.1
1	A	652	ALA	3.1
1	A	866	LEU	3.1
1	A	757	LYS	3.0
1	A	849	LYS	3.0
1	A	763	VAL	2.9
1	A	851	ILE	2.9

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	160	ASN	2.9
1	A	844	ALA	2.9
1	A	241	ALA	2.9
1	A	641	PHE	2.8
1	A	785	LYS	2.8
1	A	725	LEU	2.7
1	A	334	ILE	2.7
1	A	867	ASP	2.7
1	A	47	ASP	2.7
1	A	642	TYR	2.6
1	A	714	LEU	2.6
1	A	761	ILE	2.6
1	A	786	VAL	2.5
1	A	674	TYR	2.5
1	A	791	THR	2.4
1	A	723	THR	2.4
1	A	664	ILE	2.4
1	A	653	ASN	2.4
2	B	5	THR	2.4
1	A	783	THR	2.3
1	A	821	THR	2.3
1	A	549	SER	2.3
1	A	146	ILE	2.3
1	A	822	VAL	2.2
1	A	613	GLU	2.2
1	A	709	SER	2.2
1	A	820	TYR	2.2
1	A	68	PHE	2.2
1	A	790	VAL	2.2
1	A	333	VAL	2.1
1	A	764	LYS	2.1
1	A	649	PHE	2.1
1	A	426	ARG	2.1
1	A	552	ASP	2.1
1	A	571	GLN	2.1
1	A	46	ASN	2.0
1	A	311	LEU	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	MG	A	901	1/1	0.93	1.11	170,170,170,170	0

### 6.5 Other polymers [i](#)

There are no such residues in this entry.