



# Full wwPDB X-ray Structure Validation Report ⓘ

Apr 21, 2024 – 10:55 am BST

PDB ID : 1GO7  
Title : The metzincin's methionine: PrtC M226C-E189K double mutant  
Authors : Hege, T.  
Deposited on : 2001-10-20  
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.4, CSD as541be (2020)  
Xtrriage (Phenix) : **NOT EXECUTED**  
EDS : **NOT EXECUTED**  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36.2


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	5710 (2.10-2.10)
Ramachandran outliers	138981	5647 (2.10-2.10)
Sidechain outliers	138945	5648 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	P	462	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	PO4	P	1488	-	-	X	-

## 2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 3866 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called PROTEASE C.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	P	462	3518	2186	598	730	4	0	0	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
P	189	LYS	GLU	engineered mutation	UNP P16317
P	226	CYS	MET	engineered mutation	UNP P16317


- Molecule 2 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	P	7	Total	Ca	0	0
			7	7		

- Molecule 3 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	P	1	Total	Zn	0	0
			1	1		

- Molecule 4 is PHOSPHATE ION (three-letter code: PO4) (formula: O<sub>4</sub>P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	P	1	Total	O	P	0	0
			5	4	1		

- Molecule 5 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	P	335	Total	O	0	0
			335	335		

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

- Molecule 1: PROTEASE C



## 4 Data and refinement statistics

Xtrriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	102.70Å 102.70Å 121.27Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	39.22 – 2.10	Depositor
% Data completeness (in resolution range)	100.0 (39.22-2.10)	Depositor
$R_{merge}$	0.10	Depositor
$R_{sym}$	(Not available)	Depositor
Refinement program	REFMAC 5.0.36	Depositor
R, $R_{free}$	0.172 , 0.204	Depositor
Estimated twinning fraction	No twinning to report.	Xtrriage
Total number of atoms	3866	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	27.0	wwPDB-VP

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, PO4, CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	P	0.67	0/3597	1.01	15/4885 (0.3%)

There are no bond length outliers.

All (15) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	P	377	ASP	CB-CG-OD2	7.51	125.06	118.30
1	P	249	ASP	CB-CG-OD2	7.27	124.84	118.30
1	P	280	ASP	CB-CG-OD2	7.06	124.65	118.30
1	P	91	ASP	CB-CG-OD2	7.03	124.62	118.30
1	P	386	ASP	CB-CG-OD2	6.80	124.42	118.30
1	P	415	ASP	CB-CG-OD2	6.76	124.39	118.30
1	P	114	ASP	CB-CG-OD2	6.74	124.36	118.30
1	P	237	ASP	CB-CG-OD2	6.65	124.29	118.30
1	P	35	ASP	CB-CG-OD2	6.58	124.22	118.30
1	P	286	ASP	CB-CG-OD2	6.45	124.11	118.30
1	P	412	ASP	CB-CG-OD1	6.03	123.72	118.30
1	P	213	ASP	CB-CG-OD2	5.69	123.42	118.30
1	P	430	ASP	CB-CG-OD2	5.54	123.28	118.30
1	P	208	ASP	CB-CG-OD2	5.42	123.18	118.30
1	P	465	LEU	CB-CG-CD1	-5.11	102.31	111.00

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	P	3518	0	3209	26	0
2	P	7	0	0	0	0
3	P	1	0	0	0	0
4	P	5	0	0	2	0
5	P	335	0	0	2	1
All	All	3866	0	3209	26	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (26) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:P:413:LYS:HD3	1:P:478:ILE:HD11	1.49	0.95
1:P:203:ASN:HA	1:P:228:TYR:OH	1.81	0.80
1:P:429:GLN:O	5:P:2309:HOH:O	2.10	0.70
1:P:45:LYS:NZ	1:P:214:ALA:O	2.31	0.64
1:P:413:LYS:CD	1:P:478:ILE:HD11	2.28	0.61
1:P:189:LYS:NZ	4:P:1488:PO4:O4	2.36	0.57
1:P:95:VAL:O	1:P:135:ASN:HB2	2.06	0.56
1:P:192:HIS:CD2	4:P:1488:PO4:O2	2.62	0.52
1:P:452:LEU:C	1:P:452:LEU:HD23	2.30	0.52
1:P:421:ASN:HD22	1:P:422:GLU:N	2.09	0.50
1:P:79:LYS:HG3	1:P:123:VAL:O	2.13	0.49
1:P:289:LYS:N	1:P:289:LYS:HD3	2.27	0.49
1:P:328:LEU:HD12	1:P:328:LEU:N	2.28	0.48
1:P:328:LEU:N	1:P:328:LEU:CD1	2.76	0.48
1:P:79:LYS:HD2	1:P:81:LEU:HD23	1.95	0.47
1:P:309:TYR:HB3	1:P:311:ASN:HD22	1.80	0.46
1:P:45:LYS:HE3	1:P:211:TYR:O	2.17	0.44
1:P:443:TRP:CH2	1:P:448:SER:HA	2.52	0.44
1:P:243:GLY:HA3	5:P:2186:HOH:O	2.17	0.44
1:P:134:GLY:O	1:P:165:TRP:HA	2.17	0.44
1:P:202:TYR:CD2	1:P:209:PRO:HB3	2.53	0.44
1:P:19:ASN:O	1:P:420:ARG:NH1	2.44	0.42
1:P:317:LEU:HD21	1:P:343:ALA:HB2	2.02	0.42
1:P:100:GLU:OE1	1:P:177:GLY:N	2.47	0.42
1:P:78:PHE:HA	1:P:131:ILE:O	2.22	0.40
1:P:421:ASN:HD22	1:P:421:ASN:N	2.19	0.40



All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:P:2162:HOH:O	5:P:2298:HOH:O[2_655]	2.09	0.11

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	P	460/462 (100%)	449 (98%)	11 (2%)	0	<a href="#">100</a> <a href="#">100</a>

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	P	361/361 (100%)	359 (99%)	2 (1%)	<a href="#">86</a> <a href="#">90</a>

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	P	421	ASN
1	P	437	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	P	68	ASN
1	P	222	GLN
1	P	232	ASN
1	P	311	ASN
1	P	421	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

Of 9 ligands modelled in this entry, 8 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
4	PO4	P	1488	-	4,4,4	0.98	0	6,6,6	1.65	3 (50%)

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
4	P	1488	PO4	O2-P-O1	-2.17	102.97	110.89
4	P	1488	PO4	O4-P-O2	2.16	114.90	107.97

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	P	1488	PO4	O4-P-O3	-2.01	101.51	107.97

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	P	1488	PO4	2	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

### 6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

### 6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

### 6.4 Ligands

EDS was not executed - this section is therefore empty.

### 6.5 Other polymers

EDS was not executed - this section is therefore empty.