



Full wwPDB X-ray Structure Validation Report ⓘ

Jan 27, 2024 – 10:19 PM EST

PDB ID : 1GRB
Title : SUBSTRATE BINDING AND CATALYSIS BY GLUTATHIONE REDUCTASE AS DERIVED FROM REFINED ENZYME: SUBSTRATE CRYSTAL STRUCTURES AT 2 ANGSTROMS RESOLUTION
Authors : Karplus, P.A.; Schulz, G.E.
Deposited on : 1992-12-15
Resolution : 1.85 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtrriage (Phenix) : 1.13
EDS : 2.36
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.85 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	2625 (1.86-1.86)
Ramachandran outliers	138981	2592 (1.86-1.86)
Sidechain outliers	138945	2592 (1.86-1.86)
RSRZ outliers	127900	2436 (1.86-1.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	478	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	PO4	A	481	-	-	X	-

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 4176 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called GLUTATHIONE REDUCTASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	461	3499	2212	603	660	24	5	0	0

- Molecule 2 is PHOSPHATE ION (three-letter code: PO4) (formula: O₄P).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	O	P		
2	A	1	5	4	1	0	0

- Molecule 3 is FLAVIN-ADENINE DINUCLEOTIDE (three-letter code: FAD) (formula: C₂₇H₃₃N₉O₁₅P₂).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
3	A	1	53	27	9	15	2	0	0

- Molecule 4 is NICOTINAMIDE-ADENINE-DINUCLEOTIDE (three-letter code: NAD) (formula: $C_{21}H_{27}N_7O_{14}P_2$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
4	A	1	44	21	7	14	2	0	0

- Molecule 5 is NADPH DIHYDRO-NICOTINAMIDE-ADENINE-DINUCLEOTIDE PHOSPHATE (three-letter code: NDP) (formula: $C_{21}H_{30}N_7O_{17}P_3$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
5	A	1	48	21	7	17	3	0	0

- Molecule 6 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	527	Total	O	0	0
			527	527		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: GLUTATHIONE REDUCTASE

Chain A:  73% 18%



4 Data and refinement statistics

Property	Value	Source
Space group	B 1 1 2	Depositor
Cell constants a, b, c, α , β , γ	119.80Å 84.50Å 63.20Å 90.00° 90.00° 58.70°	Depositor
Resolution (Å)	10.00 – 1.85 72.20 – 1.85	Depositor EDS
% Data completeness (in resolution range)	(Not available) (10.00-1.85) 82.4 (72.20-1.85)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$	-	Xtrriage
Refinement program	TNT	Depositor
R, R_{free}	0.162 , (Not available) 0.176 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å ²)	16.2	Xtrriage
Anisotropy	0.106	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 91.6	EDS
L-test for twinning ¹	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	4176	wwPDB-VP
Average B, all atoms (Å ²)	23.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.83% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NAD, NDP, PO4, FAD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.89	22/3566 (0.6%)	1.22	35/4824 (0.7%)

All (22) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	77	GLU	CD-OE1	7.47	1.33	1.25
1	A	262	GLU	CD-OE1	6.78	1.33	1.25
1	A	185	GLU	CD-OE1	6.70	1.33	1.25
1	A	124	GLU	CD-OE1	6.46	1.32	1.25
1	A	317	GLU	CD-OE1	6.46	1.32	1.25
1	A	358	GLU	CD-OE2	6.45	1.32	1.25
1	A	394	GLU	CD-OE1	6.31	1.32	1.25
1	A	101	GLU	CD-OE2	6.31	1.32	1.25
1	A	91	GLU	CD-OE2	6.20	1.32	1.25
1	A	473	GLU	CD-OE2	5.98	1.32	1.25
1	A	355	GLU	CD-OE1	5.80	1.32	1.25
1	A	236	GLU	CD-OE1	5.75	1.31	1.25
1	A	427	GLU	CD-OE1	5.72	1.31	1.25
1	A	50	GLU	CD-OE2	5.65	1.31	1.25
1	A	165	GLU	CD-OE2	5.60	1.31	1.25
1	A	244	GLU	CD-OE2	5.51	1.31	1.25
1	A	239	GLU	CD-OE1	5.50	1.31	1.25
1	A	253	GLU	CD-OE2	5.37	1.31	1.25
1	A	428	GLU	CD-OE1	5.26	1.31	1.25
1	A	41	GLU	CD-OE2	5.10	1.31	1.25
1	A	472	GLU	CD-OE2	5.07	1.31	1.25
1	A	386	GLU	CD-OE1	5.05	1.31	1.25

All (35) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	91	GLU	C-N-CA	-9.22	102.93	122.30
1	A	352	ARG	NE-CZ-NH2	8.95	124.78	120.30
1	A	104	ASP	CB-CG-OD1	7.94	125.44	118.30
1	A	104	ASP	CB-CG-OD2	-7.92	111.17	118.30
1	A	461	ASP	CB-CG-OD2	-7.88	111.21	118.30
1	A	309	ASP	CB-CG-OD2	-7.71	111.36	118.30
1	A	352	ARG	NE-CZ-NH1	-7.09	116.76	120.30
1	A	441	ASP	CB-CG-OD2	-6.84	112.14	118.30
1	A	283	ASP	CB-CG-OD1	-6.70	112.27	118.30
1	A	159	MET	CB-CA-C	-6.61	97.19	110.40
1	A	84	ASP	CB-CG-OD1	-6.58	112.38	118.30
1	A	385	ASP	CB-CG-OD1	-6.51	112.44	118.30
1	A	359	ASP	CB-CG-OD2	-6.22	112.70	118.30
1	A	461	ASP	CB-CG-OD1	6.19	123.87	118.30
1	A	308	ASP	CB-CG-OD1	6.08	123.77	118.30
1	A	441	ASP	CB-CG-OD1	6.06	123.76	118.30
1	A	459	ASP	CB-CG-OD2	-6.01	112.89	118.30
1	A	309	ASP	CB-CG-OD1	5.98	123.68	118.30
1	A	316	ASP	CB-CG-OD1	-5.97	112.92	118.30
1	A	297	ASP	CB-CG-OD2	-5.97	112.93	118.30
1	A	359	ASP	CB-CG-OD1	5.94	123.65	118.30
1	A	316	ASP	CB-CG-OD2	5.88	123.59	118.30
1	A	227	ASP	CB-CG-OD1	5.65	123.38	118.30
1	A	97	ARG	NE-CZ-NH2	5.60	123.10	120.30
1	A	22	ASP	CB-CG-OD1	-5.56	113.30	118.30
1	A	84	ASP	CB-CG-OD2	5.45	123.20	118.30
1	A	178	ASP	CB-CG-OD2	-5.45	113.40	118.30
1	A	363	ASP	CB-CG-OD2	-5.37	113.47	118.30
1	A	281	ASP	CB-CG-OD1	-5.37	113.47	118.30
1	A	135	ASP	CB-CG-OD1	-5.35	113.48	118.30
1	A	308	ASP	CB-CG-OD2	-5.26	113.57	118.30
1	A	297	ASP	CB-CG-OD1	5.13	122.92	118.30
1	A	283	ASP	CB-CG-OD2	5.09	122.89	118.30
1	A	91	GLU	CA-C-N	5.08	126.36	116.20
1	A	385	ASP	CB-CG-OD2	5.00	122.80	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3499	0	3539	56	1
2	A	5	0	0	2	0
3	A	53	0	31	2	0
4	A	44	0	18	7	0
5	A	48	0	17	10	0
6	A	527	0	0	12	1
All	All	4176	0	3605	67	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (67) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:479:FAD:O5'	3:A:479:FAD:C5'	1.65	1.45
4:A:480:NAD:O3B	5:A:483:NDP:O2B	1.71	1.00
1:A:296:LYS:HB2	6:A:583:HOH:O	1.85	0.75
1:A:331:ASP:HA	1:A:337:LEU:HD22	1.73	0.70
1:A:93:LYS:O	1:A:93:LYS:HG2	1.90	0.70
1:A:127:ARG:HD2	6:A:721:HOH:O	1.93	0.68
1:A:202:MET:HA	1:A:202:MET:HE2	1.76	0.67
1:A:202:MET:HA	1:A:202:MET:CE	2.28	0.64
4:A:480:NAD:C3B	5:A:483:NDP:HO3A	2.02	0.64
3:A:479:FAD:C5'	3:A:479:FAD:P	2.86	0.64
1:A:292:VAL:HG23	1:A:293:PRO:HD2	1.80	0.63
1:A:78:PHE:HB3	6:A:958:HOH:O	1.98	0.61
1:A:293:PRO:HG3	1:A:312:HIS:CD2	2.36	0.61
1:A:222:VAL:O	1:A:231:SER:HB2	2.02	0.60
4:A:480:NAD:C3B	5:A:483:NDP:O3B	1.81	0.60
2:A:481:PO4:O2	5:A:483:NDP:H1B	2.02	0.60
1:A:185:GLU:HG3	6:A:799:HOH:O	2.01	0.59
1:A:403:PHE:CE1	1:A:473:GLU:HG3	2.39	0.58
1:A:45:ARG:NH2	1:A:124:GLU:OE1	2.38	0.57
1:A:135:ASP:HB3	1:A:136:PRO:HD2	1.86	0.56
1:A:266:VAL:HG22	1:A:276:MET:HG2	1.89	0.54
1:A:388:ILE:HG23	1:A:393:ILE:HG12	1.89	0.54
1:A:167:GLN:HG2	6:A:646:HOH:O	2.08	0.53
1:A:100:LYS:HG2	1:A:100:LYS:O	2.06	0.53
1:A:159:MET:HB3	1:A:160:PRO:HD2	1.91	0.52

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:159:MET:HB3	1:A:160:PRO:CD	2.41	0.51
1:A:125:ILE:HG22	1:A:127:ARG:HD3	1.94	0.49
4:A:480:NAD:C4N	5:A:483:NDP:H41N	0.97	0.49
1:A:334:GLY:HA2	1:A:337:LEU:HD21	1.95	0.49
4:A:480:NAD:H4N	5:A:483:NDP:C4N	0.97	0.49
4:A:480:NAD:C4N	5:A:483:NDP:H42N	0.97	0.48
1:A:164:HIS:HB2	6:A:912:HOH:O	2.13	0.48
1:A:198:ILE:O	1:A:202:MET:HG2	2.13	0.48
1:A:138:PRO:HB3	1:A:326:ILE:HD11	1.95	0.48
1:A:307:THR:HA	1:A:312:HIS:O	2.13	0.48
1:A:227:ASP:OD2	1:A:420:LYS:HE2	2.14	0.48
1:A:421:MET:SD	1:A:457:LYS:HD3	2.54	0.47
1:A:224:ARG:NH1	6:A:970:HOH:O	2.42	0.47
1:A:418:VAL:O	1:A:435:MET:HA	2.16	0.46
1:A:315:VAL:HA	1:A:320:ASN:O	2.16	0.46
1:A:82:HIS:CD2	1:A:87:PHE:CB	2.98	0.46
1:A:352:ARG:HB2	1:A:360:SER:HB3	1.98	0.46
1:A:91:GLU:HG2	1:A:92:GLY:N	2.31	0.46
1:A:427:GLU:OE1	1:A:427:GLU:N	2.46	0.46
1:A:182:GLN:HE21	1:A:182:GLN:HB3	1.60	0.46
1:A:77:GLU:HG2	6:A:1006:HOH:O	2.16	0.45
1:A:135:ASP:CB	1:A:136:PRO:HD2	2.44	0.45
1:A:82:HIS:CD2	1:A:87:PHE:HB3	2.52	0.45
1:A:465:ALA:HB1	1:A:472:GLU:HB2	1.98	0.45
1:A:202:MET:CE	1:A:205:ILE:HD12	2.48	0.43
1:A:324:LYS:HE2	6:A:748:HOH:O	2.17	0.43
1:A:138:PRO:HA	6:A:582:HOH:O	2.19	0.42
2:A:481:PO4:O2	5:A:483:NDP:C1B	2.65	0.42
1:A:135:ASP:HA	1:A:136:PRO:HD3	1.72	0.42
1:A:65:PRO:HB2	1:A:181:PHE:CE1	2.54	0.42
1:A:339:THR:HB	1:A:340:PRO:HD3	2.02	0.42
1:A:337:LEU:O	5:A:483:NDP:H1D	2.19	0.41
1:A:355:GLU:O	1:A:355:GLU:HG2	2.20	0.41
1:A:78:PHE:HD1	1:A:78:PHE:HA	1.69	0.41
1:A:132:PHE:CD1	1:A:303:LEU:HD12	2.55	0.41
1:A:188:GLY:HA3	6:A:927:HOH:O	2.20	0.41
1:A:223:LEU:O	1:A:231:SER:HB3	2.20	0.41
1:A:427:GLU:O	1:A:428:GLU:HB2	2.21	0.41
5:A:483:NDP:H1B	6:A:594:HOH:O	2.20	0.41
1:A:337:LEU:O	4:A:480:NAD:H1D	2.19	0.41
1:A:258:LEU:HA	1:A:258:LEU:HD23	1.86	0.40

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:202:MET:HE2	1:A:205:ILE:HD12	2.03	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
6:A:580:HOH:O	6:A:580:HOH:O[2_665]	0.76	1.44
1:A:90:CYS:SG	1:A:90:CYS:SG[2_665]	1.62	0.58

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	459/478 (96%)	440 (96%)	18 (4%)	1 (0%)	47 33

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	136	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	382/393 (97%)	359 (94%)	23 (6%)	19 5

All (23) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	75	HIS
1	A	89	SER
1	A	90	CYS
1	A	91	GLU
1	A	93	LYS
1	A	127	ARG
1	A	134	SER
1	A	167	GLN
1	A	185	GLU
1	A	212	LYS
1	A	252	LYS
1	A	253	GLU
1	A	256	LYS
1	A	272	ARG
1	A	275	VAL
1	A	276	MET
1	A	296	LYS
1	A	317	GLU
1	A	335	LYS
1	A	394	GLU
1	A	397	LYS
1	A	426	LYS
1	A	478	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	52	HIS
1	A	82	HIS
1	A	117	ASN
1	A	129	HIS
1	A	182	GLN
1	A	301	ASN
1	A	366	ASN
1	A	425	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	PO4	A	481	-	4,4,4	1.43	0	6,6,6	1.59	1 (16%)
4	NAD	A	480	-	42,48,48	1.60	5 (11%)	50,73,73	1.75	13 (26%)
5	NDP	A	483	-	45,52,52	1.78	9 (20%)	53,80,80	1.44	10 (18%)
3	FAD	A	479	-	53,58,58	1.50	6 (11%)	68,89,89	1.33	9 (13%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAD	A	480	-	-	4/26/62/62	0/5/5/5
5	NDP	A	483	-	-	6/30/77/77	0/5/5/5
3	FAD	A	479	-	-	5/30/50/50	0/6/6/6

All (20) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	480	NAD	O4B-C1B	5.50	1.48	1.41
3	A	479	FAD	O4B-C1B	5.44	1.48	1.41

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	479	FAD	O5'-C5'	5.30	1.65	1.44
4	A	480	NAD	O4D-C1D	5.29	1.48	1.41
5	A	483	NDP	O4B-C1B	5.02	1.48	1.41
5	A	483	NDP	C6N-C5N	4.97	1.42	1.33
5	A	483	NDP	C4N-C3N	-4.09	1.41	1.49
3	A	479	FAD	C5'-C4'	3.63	1.56	1.51
5	A	483	NDP	PA-O5B	3.48	1.73	1.59
5	A	483	NDP	P2B-O2B	3.10	1.65	1.59
5	A	483	NDP	O4D-C1D	2.73	1.48	1.42
3	A	479	FAD	P-O2P	-2.48	1.43	1.55
3	A	479	FAD	O5B-C5B	2.26	1.53	1.44
5	A	483	NDP	C7N-C3N	2.24	1.53	1.48
4	A	480	NAD	C2B-C1B	-2.23	1.50	1.53
4	A	480	NAD	C2N-N1N	2.19	1.37	1.35
5	A	483	NDP	C4N-C5N	-2.15	1.43	1.48
5	A	483	NDP	C2A-N1A	2.13	1.37	1.33
4	A	480	NAD	C5N-C4N	2.07	1.43	1.38
3	A	479	FAD	C1'-C2'	2.03	1.55	1.52

All (33) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	480	NAD	C6N-N1N-C2N	-5.59	116.88	121.97
4	A	480	NAD	C3N-C2N-N1N	3.96	124.30	120.43
3	A	479	FAD	C5'-C4'-C3'	-3.95	104.57	112.20
3	A	479	FAD	P-O3P-PA	3.84	146.02	132.83
4	A	480	NAD	C4A-C5A-N7A	3.40	112.95	109.40
3	A	479	FAD	O4B-C1B-C2B	-3.29	102.11	106.93
4	A	480	NAD	O2N-PN-O1N	3.11	127.63	112.24
5	A	483	NDP	O2N-PN-O1N	3.11	127.63	112.24
3	A	479	FAD	O5B-C5B-C4B	-2.95	98.84	108.99
5	A	483	NDP	O2A-PA-O1A	2.83	126.21	112.24
4	A	480	NAD	O2A-PA-O1A	2.83	126.21	112.24
4	A	480	NAD	O4D-C4D-C3D	-2.76	99.66	105.11
5	A	483	NDP	O4D-C4D-C3D	-2.76	99.66	105.11
4	A	480	NAD	O7N-C7N-C3N	-2.74	116.35	119.63
3	A	479	FAD	O2P-P-O1P	2.68	125.51	112.24
5	A	483	NDP	C2A-N1A-C6A	2.68	123.34	118.75
3	A	479	FAD	C4A-C5A-N7A	2.65	112.17	109.40
3	A	479	FAD	O2A-PA-O1A	2.50	124.62	112.24
5	A	483	NDP	C5A-C6A-N1A	-2.49	114.70	120.35
5	A	483	NDP	C4A-C5A-N7A	2.49	111.99	109.40

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	480	NAD	C5N-C4N-C3N	-2.46	117.43	120.34
5	A	483	NDP	C5A-C6A-N6A	2.42	124.03	120.35
5	A	483	NDP	O7N-C7N-C3N	-2.42	116.34	120.90
3	A	479	FAD	C5A-C6A-N6A	2.37	123.95	120.35
4	A	480	NAD	O4B-C1B-C2B	-2.32	103.54	106.93
4	A	480	NAD	O5B-C5B-C4B	-2.30	101.08	108.99
5	A	483	NDP	C1D-N1N-C6N	2.22	125.62	120.83
5	A	483	NDP	C1D-N1N-C2N	-2.19	117.46	121.11
2	A	481	PO4	O4-P-O1	-2.13	103.12	110.89
4	A	480	NAD	C5A-C6A-N6A	2.07	123.50	120.35
4	A	480	NAD	C5A-C6A-N1A	-2.05	115.70	120.35
4	A	480	NAD	O2B-C2B-C3B	-2.02	105.29	111.82
3	A	479	FAD	C2A-N1A-C6A	2.00	122.18	118.75

There are no chirality outliers.

All (15) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	479	FAD	PA-O3P-P-O5'
5	A	483	NDP	O4D-C1D-N1N-C2N
3	A	479	FAD	O4B-C4B-C5B-O5B
5	A	483	NDP	O4B-C4B-C5B-O5B
5	A	483	NDP	C3B-C4B-C5B-O5B
3	A	479	FAD	C3B-C4B-C5B-O5B
4	A	480	NAD	C3D-C4D-C5D-O5D
5	A	483	NDP	C3D-C4D-C5D-O5D
3	A	479	FAD	P-O3P-PA-O2A
4	A	480	NAD	O4B-C4B-C5B-O5B
3	A	479	FAD	P-O3P-PA-O1A
4	A	480	NAD	O4D-C4D-C5D-O5D
5	A	483	NDP	O4D-C4D-C5D-O5D
4	A	480	NAD	PA-O3-PN-O2N
5	A	483	NDP	PA-O3-PN-O2N

There are no ring outliers.

4 monomers are involved in 13 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	481	PO4	2	0
4	A	480	NAD	7	0
5	A	483	NDP	10	0


Continued on next page...


Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	479	FAD	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	461/478 (96%)	-0.64	0 100 100	8, 18, 44, 70	1 (0%)

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	PO4	A	481	5/5	0.94	0.30	21,30,34,38	0
4	NAD	A	480	44/44	0.96	0.32	8,17,27,40	0
5	NDP	A	483	48/48	0.97	0.34	8,18,29,40	0
3	FAD	A	479	53/53	0.98	0.07	5,12,19,21	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.


Electron density around NAD A 480:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around NDP A 483:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.