

wwPDB X-ray Structure Validation Summary Report (i)

Feb 12, 2024 – 04:04 AM EST

PDB ID : 3GRK

Title: Crystal structure of short chain dehydrogenase reductase SDR glucose-ribitol

dehydrogenase from Brucella melitensis

Authors : Seattle Structural Genomics Center for Infectious Disease (SSGCID)

Deposited on : 2009-03-25

Resolution : 2.35 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*A user guide is available at

https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity: 4.02b-467 Xtriage (Phenix): 1.13

EDS: 2.36

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

 $Refmac \quad : \quad 5.8.0158$

CCP4 : 7.0.044 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.36

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X- $RAY\ DIFFRACTION$

The reported resolution of this entry is 2.35 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive	Similar resolution
Metric	$(\# \mathrm{Entries})$	$(\# ext{Entries}, ext{ resolution range}(ext{Å}))$
R_{free}	130704	1164 (2.36-2.36)
Clashscore	141614	1232 (2.36-2.36)
Ramachandran outliers	138981	1211 (2.36-2.36)
Sidechain outliers	138945	1212 (2.36-2.36)
RSRZ outliers	127900	1150 (2.36-2.36)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain		
1	A	293	73%	10% •	15%
1	В	293	76%	9% •	14%
1	С	293	7%	16%	15%
1	D	293	7% 73%	13%	14%
1	Е	293	76%	9%	15%



Mol	Chain	Length	Quality of chain			
1	F	293	72%	13%	•	14%
1	G	293	71%	13%		16%
1	Н	293	5% 74%	12%		14%



2 Entry composition (i)

There are 2 unique types of molecules in this entry. The entry contains 15671 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Enoyl-(acyl-carrier-protein) reductase (NADH).

Mol	Chain	Residues		Ato	oms			ZeroOcc	AltConf	Trace
1	A	248	Total	С	N	О	S	0	0	0
1	A	240	1863	1184	314	359	6	0	U	0
1	В	252	Total	С	N	О	S	0	0	0
1	Б	202	1904	1211	324	363	6	0	U	
1	С	248	Total	С	N	О	S	0	0	0
1		240	1875	1191	320	358	6	0	U	
1	D	251	Total	С	N	О	S	0	0	0
1	D	201	1909	1214	327	362	6	0	U	
1	Е	248	Total	С	N	О	S	0	0	0
1	l Li	240	1883	1196	322	359	6	0		
1	F	252	Total	С	N	О	S	0	0	0
1	I.	202	1907	1211	326	364	6	U	U	U
1	G	247	Total	С	N	О	S	0	0	0
1	G	241	1869	1189	320	354	6	0	0	
1	Н	252	Total	С	N	О	S	0	0	0
1	11	202	1898	1205	324	363	6		U	

There are 168 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-20	MET	-	expression tag	UNP Q8YFK8
A	-19	ALA	-	expression tag	UNP Q8YFK8
A	-18	HIS	-	expression tag	UNP Q8YFK8
A	-17	HIS	-	expression tag	UNP Q8YFK8
A	-16	HIS	-	expression tag	UNP Q8YFK8
A	-15	HIS	-	expression tag	UNP Q8YFK8
A	-14	HIS	-	expression tag	UNP Q8YFK8
A	-13	HIS	-	expression tag	UNP Q8YFK8
A	-12	MET	-	expression tag	UNP Q8YFK8
A	-11	GLY	_	expression tag	UNP Q8YFK8
A	-10	THR	-	expression tag	UNP Q8YFK8
A	-9	LEU		expression tag	UNP Q8YFK8
A	-8	GLU	-	expression tag	UNP Q8YFK8



 $Continued\ from\ previous\ page...$

Chain	Residue	Modelled Modelled	Actual	Comment	Reference
A	-7	ALA	-	expression tag	UNP Q8YFK8
A	-6	GLN	-	expression tag	UNP Q8YFK8
A	-5	THR	-	expression tag	UNP Q8YFK8
A	-4	GLN	-	expression tag	UNP Q8YFK8
A	-3	GLY	-	expression tag	UNP Q8YFK8
A	-2	PRO	_	expression tag	UNP Q8YFK8
A	-1	GLY	-	expression tag	UNP Q8YFK8
A	0	SER	-	expression tag	UNP Q8YFK8
В	-20	MET	-	expression tag	UNP Q8YFK8
В	-19	ALA	-	expression tag	UNP Q8YFK8
В	-18	HIS	-	expression tag	UNP Q8YFK8
В	-17	HIS	-	expression tag	UNP Q8YFK8
В	-16	HIS	-	expression tag	UNP Q8YFK8
В	-15	HIS	-	expression tag	UNP Q8YFK8
В	-14	HIS	-	expression tag	UNP Q8YFK8
В	-13	HIS	-	expression tag	UNP Q8YFK8
В	-12	MET	-	expression tag	UNP Q8YFK8
В	-11	GLY	_	expression tag	UNP Q8YFK8
В	-10	THR	-	expression tag	UNP Q8YFK8
В	-9	LEU	-	expression tag	UNP Q8YFK8
В	-8	GLU	-	expression tag	UNP Q8YFK8
В	-7	ALA	-	expression tag	UNP Q8YFK8
В	-6	GLN	-	expression tag	UNP Q8YFK8
В	-5	THR	-	expression tag	UNP Q8YFK8
В	-4	GLN	-	expression tag	UNP Q8YFK8
В	-3	GLY	-	expression tag	UNP Q8YFK8
В	-2	PRO	-	expression tag	UNP Q8YFK8
В	-1	GLY	-	expression tag	UNP Q8YFK8
В	0	SER	-	expression tag	UNP Q8YFK8
С	-20	MET	_	expression tag	UNP Q8YFK8
С	-19	ALA	_	expression tag	UNP Q8YFK8
С	-18	HIS	-	expression tag	UNP Q8YFK8
С	-17	HIS	-	expression tag	UNP Q8YFK8
С	-16	HIS	-	expression tag	UNP Q8YFK8
С	-15	HIS	-	expression tag	UNP Q8YFK8
С	-14	HIS	-	expression tag	UNP Q8YFK8
С	-13	HIS	-	expression tag	UNP Q8YFK8
С	-12	MET		expression tag	UNP Q8YFK8
С	-11	GLY	-	expression tag	UNP Q8YFK8
С	-10	THR	-	expression tag	UNP Q8YFK8
С	-9	LEU	-	expression tag	UNP Q8YFK8
С	-8	GLU	_	expression tag	UNP Q8YFK8



 $Continued\ from\ previous\ page...$

Chain	Residue	Modelled Modelled	Actual	Comment	Reference
С	-7	ALA	-	expression tag	UNP Q8YFK8
С	-6	GLN	-	expression tag	UNP Q8YFK8
С	-5	THR	-	expression tag	UNP Q8YFK8
С	-4	GLN	-	expression tag	UNP Q8YFK8
С	-3	GLY	-	expression tag	UNP Q8YFK8
С	-2	PRO	-	expression tag	UNP Q8YFK8
С	-1	GLY	-	expression tag	UNP Q8YFK8
С	0	SER	-	expression tag	UNP Q8YFK8
D	-20	MET	-	expression tag	UNP Q8YFK8
D	-19	ALA	-	expression tag	UNP Q8YFK8
D	-18	HIS	-	expression tag	UNP Q8YFK8
D	-17	HIS	-	expression tag	UNP Q8YFK8
D	-16	HIS	-	expression tag	UNP Q8YFK8
D	-15	HIS	-	expression tag	UNP Q8YFK8
D	-14	HIS	-	expression tag	UNP Q8YFK8
D	-13	HIS	-	expression tag	UNP Q8YFK8
D	-12	MET	-	expression tag	UNP Q8YFK8
D	-11	GLY	-	expression tag	UNP Q8YFK8
D	-10	THR	-	expression tag	UNP Q8YFK8
D	-9	LEU	-	expression tag	UNP Q8YFK8
D	-8	GLU	-	expression tag	UNP Q8YFK8
D	-7	ALA	-	expression tag	UNP Q8YFK8
D	-6	GLN	-	expression tag	UNP Q8YFK8
D	-5	THR	-	expression tag	UNP Q8YFK8
D	-4	GLN	-	expression tag	UNP Q8YFK8
D	-3	GLY	-	expression tag	UNP Q8YFK8
D	-2	PRO	-	expression tag	UNP Q8YFK8
D	-1	GLY	-	expression tag	UNP Q8YFK8
D	0	SER	-	expression tag	UNP Q8YFK8
Е	-20	MET	-	expression tag	UNP Q8YFK8
E	-19	ALA	_	expression tag	UNP Q8YFK8
E	-18	HIS	-	expression tag	UNP Q8YFK8
Е	-17	HIS	-	expression tag	UNP Q8YFK8
Е	-16	HIS	-	expression tag	UNP Q8YFK8
Е	-15	HIS	-	expression tag	UNP Q8YFK8
Е	-14	HIS	-	expression tag	UNP Q8YFK8
Е	-13	HIS		expression tag	UNP Q8YFK8
Е	-12	MET		expression tag	UNP Q8YFK8
Е	-11	GLY	-	expression tag	UNP Q8YFK8
Е	-10	THR	-	expression tag	UNP Q8YFK8
Е	-9	LEU	-	expression tag	UNP Q8YFK8
Е	-8	GLU	_	expression tag	UNP Q8YFK8



 $Continued\ from\ previous\ page...$

Chain	Residue	Modelled Modelled	Actual	Comment	Reference
Е	-7	ALA	-	expression tag	UNP Q8YFK8
Е	-6	GLN	-	expression tag	UNP Q8YFK8
Е	-5	THR	-	expression tag	UNP Q8YFK8
Е	-4	GLN	-	expression tag	UNP Q8YFK8
Е	-3	GLY	-	expression tag	UNP Q8YFK8
Е	-2	PRO	_	expression tag	UNP Q8YFK8
Е	-1	GLY	-	expression tag	UNP Q8YFK8
Е	0	SER	-	expression tag	UNP Q8YFK8
F	-20	MET	-	expression tag	UNP Q8YFK8
F	-19	ALA	-	expression tag	UNP Q8YFK8
F	-18	HIS	-	expression tag	UNP Q8YFK8
F	-17	HIS	-	expression tag	UNP Q8YFK8
F	-16	HIS	-	expression tag	UNP Q8YFK8
F	-15	HIS	-	expression tag	UNP Q8YFK8
F	-14	HIS	-	expression tag	UNP Q8YFK8
F	-13	HIS	-	expression tag	UNP Q8YFK8
F	-12	MET	-	expression tag	UNP Q8YFK8
F	-11	GLY	-	expression tag	UNP Q8YFK8
F	-10	THR	-	expression tag	UNP Q8YFK8
F	-9	LEU	-	expression tag	UNP Q8YFK8
F	-8	GLU	-	expression tag	UNP Q8YFK8
F	-7	ALA	-	expression tag	UNP Q8YFK8
F	-6	GLN	-	expression tag	UNP Q8YFK8
F	-5	THR	-	expression tag	UNP Q8YFK8
F	-4	GLN	-	expression tag	UNP Q8YFK8
F	-3	GLY	-	expression tag	UNP Q8YFK8
F	-2	PRO	-	expression tag	UNP Q8YFK8
F	-1	GLY	-	expression tag	UNP Q8YFK8
F	0	SER	-	expression tag	UNP Q8YFK8
G	-20	MET	-	expression tag	UNP Q8YFK8
G	-19	ALA	_	expression tag	UNP Q8YFK8
G	-18	HIS	-	expression tag	UNP Q8YFK8
G	-17	HIS	-	expression tag	UNP Q8YFK8
G	-16	HIS	-	expression tag	UNP Q8YFK8
G	-15	HIS	-	expression tag	UNP Q8YFK8
G	-14	HIS	-	expression tag	UNP Q8YFK8
G	-13	HIS	-	expression tag	UNP Q8YFK8
G	-12	MET	-	expression tag	UNP Q8YFK8
G	-11	GLY		expression tag	UNP Q8YFK8
G	-10	THR	-	expression tag	UNP Q8YFK8
G	-9	LEU	-	expression tag	UNP Q8YFK8
G	-8	GLU	_	expression tag	UNP Q8YFK8



Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
G	-7	ALA	-	expression tag	UNP Q8YFK8
G	-6	GLN	-	expression tag	UNP Q8YFK8
G	-5	THR	-	expression tag	UNP Q8YFK8
G	-4	GLN	-	expression tag	UNP Q8YFK8
G	-3	GLY	-	expression tag	UNP Q8YFK8
G	-2	PRO	-	expression tag	UNP Q8YFK8
G	-1	GLY	-	expression tag	UNP Q8YFK8
G	0	SER	-	expression tag	UNP Q8YFK8
Н	-20	MET	-	expression tag	UNP Q8YFK8
Н	-19	ALA	-	expression tag	UNP Q8YFK8
Н	-18	HIS	-	expression tag	UNP Q8YFK8
Н	-17	HIS	-	expression tag	UNP Q8YFK8
Н	-16	HIS	-	expression tag	UNP Q8YFK8
Н	-15	HIS	-	expression tag	UNP Q8YFK8
Н	-14	HIS	-	expression tag	UNP Q8YFK8
Н	-13	HIS	-	expression tag	UNP Q8YFK8
Н	-12	MET	-	expression tag	UNP Q8YFK8
Н	-11	GLY	-	expression tag	UNP Q8YFK8
Н	-10	THR	-	expression tag	UNP Q8YFK8
Н	-9	LEU	-	expression tag	UNP Q8YFK8
Н	-8	GLU	-	expression tag	UNP Q8YFK8
Н	-7	ALA	-	expression tag	UNP Q8YFK8
Н	-6	GLN	-	expression tag	UNP Q8YFK8
Н	-5	THR	-	expression tag	UNP Q8YFK8
Н	-4	GLN	-	expression tag	UNP Q8YFK8
Н	-3	GLY	-	expression tag	UNP Q8YFK8
Н	-2	PRO	-	expression tag	UNP Q8YFK8
Н	-1	GLY	-	expression tag	UNP Q8YFK8
Н	0	SER	-	expression tag	UNP Q8YFK8

• Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	64	Total O	0	0
		01	64 64		Ŭ
2	В	88	Total O	0	0
	Б	00	88 88		0
2	С	58	Total O	0	0
		90	58 58	0	U
2	D	# # #	Total O	0	0
	D	55	55 55	0	U
2	E	01	Total O	0	0
	E	91	91 91	U	U




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	F	77	Total O 77 77	0	0
2	G	56	Total O 56 56	0	0
2	Н	74	Total O 74 74	0	0




3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Enoyl-(acyl-carrier-protein) reductase (NADH)










H255 M259 L17S L17S ALA ALA ASP ASP ASP ASP ASP ASP ASP ASP ASP

• Molecule 1: Enoyl-(acyl-carrier-protein) reductase (NADH)





4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants	66.10Å 120.10Å 137.70Å	Donositor
a, b, c, α , β , γ	90.00° 99.20° 90.00°	Depositor
Resolution (Å)	19.81 - 2.35	Depositor
Resolution (A)	19.82 - 2.35	EDS
% Data completeness	95.3 (19.81-2.35)	Depositor
(in resolution range)	95.3 (19.82-2.35)	EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	3.81 (at 2.35Å)	Xtriage
Refinement program	REFMAC	Depositor
D D.	0.206 , 0.269	Depositor
R, R_{free}	0.201 , 0.260	DCC
R_{free} test set	4211 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å ²)	29.4	Xtriage
Anisotropy	0.091	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.40 , 44.1	EDS
L-test for twinning ²	$ < L > = 0.50, < L^2> = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	15671	wwPDB-VP
Average B, all atoms (Å ²)	27.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The analyses of the Patterson function reveals a significant off-origin peak that is 38.22 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 3.8119e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.

²Theoretical values of <|L|>, $<L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	Bond lengths		angles
IVIOI	Chain	RMSZ	# Z > 5	RMSZ	# Z > 5
1	A	0.47	0/1896	0.58	0/2573
1	В	0.48	0/1937	0.59	0/2624
1	С	0.47	0/1908	0.60	0/2586
1	D	0.46	0/1942	0.59	0/2628
1	Е	0.50	0/1916	0.60	0/2595
1	F	0.47	0/1940	0.59	0/2628
1	G	0.47	0/1902	0.59	0/2576
1	Н	0.46	0/1931	0.59	0/2619
All	All	0.47	0/15372	0.59	0/20829

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1863	0	1822	20	0
1	В	1904	0	1889	20	0
1	С	1875	0	1849	37	0
1	D	1909	0	1909	24	0
1	Е	1883	0	1866	16	0
1	F	1907	0	1892	28	0
1	G	1869	0	1851	26	0
1	Н	1898	0	1867	22	0



$\alpha \cdots$, r	•	
Continued	trom	mromonie	maaa
-	110116	DICULUUS	Duuc
	J	1	1

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	A	64	0	0	1	0
2	В	88	0	0	2	0
2	С	58	0	0	0	0
2	D	55	0	0	1	0
2	${ m E}$	91	0	0	0	0
2	F	77	0	0	1	0
2	G	56	0	0	2	0
2	Н	74	0	0	0	0
All	All	15671	0	14945	171	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 171 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	$\begin{array}{c} {\rm Interatomic} \\ {\rm distance} \ ({\rm \AA}) \end{array}$	Clash overlap (Å)
1:F:92:LEU:HD11	1:F:130:VAL:CG1	2.10	0.81
1:D:11:LYS:NZ	1:D:238:ASP:OD1	2.15	0.79
1:F:134:ALA:HB1	1:F:138:MET:CE	2.13	0.77
1:F:92:LEU:HD11	1:F:130:VAL:HG12	1.65	0.76
1:C:92:LEU:HD11	1:C:130:VAL:CG1	2.16	0.74

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	244/293~(83%)	237 (97%)	7 (3%)	0	100 100
1	В	248/293~(85%)	238 (96%)	10 (4%)	0	100 100
1	C	244/293~(83%)	235 (96%)	9 (4%)	0	100 100



Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	D	247/293 (84%)	238 (96%)	9 (4%)	0	100	100
1	E	244/293 (83%)	236 (97%)	8 (3%)	0	100	100
1	F	248/293 (85%)	240 (97%)	7 (3%)	1 (0%)	34	38
1	G	243/293 (83%)	234 (96%)	9 (4%)	0	100	100
1	Н	248/293 (85%)	240 (97%)	8 (3%)	0	100	100
All	All	1966/2344 (84%)	1898 (96%)	67 (3%)	1 (0%)	51	63

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	F	69	VAL

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	187/227 (82%)	177 (95%)	10 (5%)	22 26
1	В	192/227 (85%)	186 (97%)	6 (3%)	40 48
1	\mathbf{C}	189/227 (83%)	185 (98%)	4 (2%)	53 65
1	D	194/227 (86%)	189 (97%)	5 (3%)	46 56
1	\mathbf{E}	191/227 (84%)	189 (99%)	2 (1%)	76 85
1	F	193/227 (85%)	188 (97%)	5 (3%)	46 56
1	G	188/227 (83%)	181 (96%)	7 (4%)	34 42
1	Н	190/227 (84%)	187 (98%)	3 (2%)	62 75
All	All	1524/1816 (84%)	1482 (97%)	42 (3%)	43 53

5 of 42 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	F	82	LEU
1	G	176	LEU



Mol	Chain	Res	Type
1	F	101	LYS
1	G	12	ARG
1	G	207	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	G	158	ASN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

There are no ligands in this entry.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<rsrz></rsrz>	#RSI	RZ>2	2	$\mathbf{OWAB}(\mathbf{\mathring{A}}^2)$	Q < 0.9
1	A	248/293~(84%)	0.40	14 (5%)	24	35	15, 29, 49, 59	0
1	В	252/293~(86%)	0.11	13 (5%)	27	39	10, 24, 40, 47	0
1	С	248/293 (84%)	0.34	21 (8%)	10	16	14, 27, 48, 57	0
1	D	251/293~(85%)	0.36	20 (7%)	12	18	13, 27, 51, 61	0
1	Е	248/293 (84%)	-0.00	7 (2%)	53 6	64	11, 21, 34, 49	0
1	F	252/293~(86%)	0.27	15 (5%)	21	32	15, 27, 45, 53	0
1	G	247/293~(84%)	0.23	15 (6%)	21	31	12, 26, 48, 56	0
1	Н	252/293~(86%)	0.27	14 (5%)	24	35	14, 26, 48, 58	0
All	All	1998/2344 (85%)	0.25	119 (5%)	21	32	10, 26, 47, 61	0

The worst 5 of 119 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	F	204	GLY	7.2
1	A	204	GLY	5.7
1	G	215	TYR	5.0
1	A	215	TYR	4.9
1	Н	47	ALA	4.8

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.



6.4 Ligands (i)

There are no ligands in this entry.

6.5 Other polymers (i)

There are no such residues in this entry.

