



# Full wwPDB X-ray Structure Validation Report ⓘ

Sep 18, 2023 – 02:08 PM JST

PDB ID : 8GY8  
Title : The structure of Bax1 from *Pyrococcus furiosus*  
Authors : Wang, W.W.; Liu, X.P.  
Deposited on : 2022-09-21  
Resolution : 1.94 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.35.1  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.35.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.94 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	4310 (1.96-1.92)
Clashscore	141614	1023 (1.94-1.94)
Ramachandran outliers	138981	1007 (1.94-1.94)
Sidechain outliers	138945	1007 (1.94-1.94)
RSRZ outliers	127900	4250 (1.96-1.92)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	375	 6% 79% 19% ..
1	B	375	 8% 82% 16% .

## 2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 6606 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Putative translation initiation factor (IF-2 homolog).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	375	Total 3091	C 1988	N 532	O 560	S 11	0	1	0
1	B	375	Total 3080	C 1982	N 528	O 559	S 11	0	0	0

- Molecule 2 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	O	S		
2	A	1	Total 5	O 4	S 1	0	0

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	201	Total 201	O 201	0	0

*Continued on next page...*


*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Residues</b>	<b>Atoms</b>		<b>ZeroOcc</b>	<b>AltConf</b>
3	B	229	Total 229	O 229	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Putative translation initiation factor (IF-2 homolog)



- Molecule 1: Putative translation initiation factor (IF-2 homolog)



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	82.25Å 110.49Å 126.14Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	45.59 – 1.94 45.59 – 1.94	Depositor EDS
% Data completeness (in resolution range)	100.0 (45.59-1.94) 100.0 (45.59-1.94)	Depositor EDS
$R_{merge}$	0.06	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.24 (at 1.94Å)	Xtrriage
Refinement program	PHENIX 1.20_4459	Depositor
R, $R_{free}$	0.198 , 0.218 0.194 , 0.214	Depositor DCC
$R_{free}$ test set	4299 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	35.6	Xtrriage
Anisotropy	0.037	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 51.8	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	6606	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	47.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.85% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.37	0/3140	0.60	0/4208
1	B	0.37	0/3129	0.61	0/4194
All	All	0.37	0/6269	0.61	0/8402

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	7

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (7) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	11	ARG	Sidechain
1	A	164	ARG	Sidechain
1	A	207	ARG	Sidechain
1	A	237	ARG	Sidechain
1	A	266	ARG	Sidechain
1	A	274	ARG	Sidechain
1	A	357	ARG	Sidechain

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3091	0	3235	53	0
1	B	3080	0	3223	46	0
2	A	5	0	0	0	0
3	A	201	0	0	11	3
3	B	229	0	0	7	3
All	All	6606	0	6458	98	3

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (98) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:357:ARG:HH12	1:B:358:LYS:HE2	1.20	1.04
1:B:37:LYS:O	3:B:401:HOH:O	1.98	0.81
1:B:357:ARG:NH1	1:B:358:LYS:HE2	1.98	0.79
1:B:19:LYS:NZ	3:B:403:HOH:O	2.13	0.79
1:B:303:ASN:HB2	1:B:305:LYS:NZ	2.01	0.76
1:A:94:ARG:NE	3:A:501:HOH:O	2.19	0.73
1:A:53:HIS:ND1	3:A:502:HOH:O	2.20	0.72
1:B:3:PRO:HB2	1:B:72:ARG:NH1	2.05	0.71
1:A:278:VAL:HG23	1:A:302:LYS:HB2	1.73	0.70
1:A:208:LYS:HE2	1:A:208:LYS:H	1.56	0.70
1:B:284:PRO:O	3:B:402:HOH:O	2.09	0.69
1:B:229:LYS:HE2	3:B:446:HOH:O	1.92	0.69
1:B:369:LYS:O	1:B:373:MET:HG3	1.93	0.68
1:B:284:PRO:HG2	1:B:296:PRO:HG2	1.76	0.66
1:B:303:ASN:HB2	1:B:305:LYS:HZ1	1.61	0.66
1:B:276:LEU:HD13	1:B:367:ILE:HG23	1.81	0.63
1:A:273:LYS:NZ	3:A:505:HOH:O	2.29	0.61
1:B:301:ARG:HG3	1:B:306:GLU:HG2	1.82	0.61
1:A:303:ASN:OD1	1:A:304:GLY:N	2.34	0.60
1:B:3:PRO:HD3	3:B:424:HOH:O	2.00	0.60
1:A:2:LEU:O	1:A:6:LEU:HD12	2.01	0.59
1:A:258:VAL:O	1:A:258:VAL:HG12	2.03	0.59

*Continued on next page...*



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:205:MET:HA	1:A:208:LYS:HE3	1.85	0.58
1:A:335:LEU:HD11	1:A:354:ILE:HG13	1.86	0.56
1:B:3:PRO:HB2	1:B:72:ARG:HH11	1.71	0.56
1:B:76:PHE:CG	1:B:133:LYS:HE3	2.41	0.55
1:A:73:GLU:HB2	1:A:139:PRO:HG3	1.89	0.54
1:B:12:ARG:CZ	1:B:13:ARG:HE	2.20	0.54
1:B:303:ASN:HB2	1:B:305:LYS:HZ3	1.73	0.54
1:B:266:ARG:HA	1:B:282:ARG:NH1	2.24	0.53
1:A:105:LYS:O	1:A:109:GLU:HG3	2.09	0.53
1:A:205:MET:HA	1:A:208:LYS:CE	2.38	0.53
1:A:208:LYS:H	1:A:208:LYS:CE	2.22	0.52
1:A:214:ALA:HA	1:A:217:ILE:HD12	1.92	0.52
1:B:266:ARG:HA	1:B:282:ARG:HH12	1.75	0.52
1:A:53:HIS:CE1	3:A:502:HOH:O	2.62	0.51
1:B:141:ILE:HG13	1:B:145:GLU:HB3	1.93	0.51
1:A:23:GLU:HG3	3:A:657:HOH:O	2.11	0.51
1:B:4:LYS:HD2	1:B:5:GLU:N	2.26	0.51
1:A:178:ARG:O	1:A:182:LEU:HG	2.11	0.50
1:B:7:LEU:O	1:B:154:LEU:HD11	2.11	0.50
1:B:262:SER:HA	1:B:265:GLU:HG3	1.93	0.50
1:B:324:LEU:HD11	1:B:353:VAL:HG13	1.94	0.50
1:B:333:LYS:HE2	1:B:373:MET:O	2.12	0.49
1:A:12:ARG:HG3	1:A:13:ARG:HG3	1.95	0.49
1:A:12:ARG:NE	1:A:12:ARG:HA	2.28	0.49
1:A:5:GLU:O	1:A:11:ARG:NH2	2.38	0.48
1:A:171:GLU:OE1	1:A:224:LYS:HG2	2.13	0.48
1:B:164:ARG:NE	1:B:231:GLU:OE1	2.46	0.48
1:B:302:LYS:O	1:B:305:LYS:N	2.44	0.48
1:B:350:ILE:HG13	1:B:353:VAL:HG22	1.96	0.47
1:A:275:ILE:HG13	3:A:508:HOH:O	2.14	0.47
1:A:12:ARG:NH2	3:A:510:HOH:O	2.47	0.47
1:A:58:GLU:HG2	1:A:59:ASN:N	2.30	0.47
1:B:125:MET:HE2	1:B:125:MET:HB2	1.85	0.46
1:A:301:ARG:HE	1:A:306:GLU:CG	2.28	0.46
1:A:96:TYR:CZ	1:A:103:ARG:HD3	2.50	0.46
1:A:6:LEU:HB2	1:A:7:LEU:HD22	1.97	0.46
1:A:222:LYS:NZ	1:A:254:LEU:O	2.35	0.46
1:A:274:ARG:HA	1:A:274:ARG:HD3	1.47	0.46
1:A:4:LYS:HG2	1:A:61:ARG:NH2	2.30	0.45
1:B:68:LYS:HG3	1:B:72:ARG:NH1	2.31	0.45
1:A:269:VAL:HG22	1:A:280:VAL:HG11	1.99	0.44

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:208:LYS:H	1:A:208:LYS:CD	2.29	0.44
1:B:351:HIS:CG	3:B:416:HOH:O	2.70	0.44
1:B:44:TYR:HB2	1:B:133:LYS:HG2	2.00	0.44
1:A:5:GLU:OE2	3:A:503:HOH:O	2.21	0.44
1:B:152:LEU:HD21	1:B:218:PRO:HG3	2.00	0.44
1:B:287:ILE:HG21	1:B:308:TYR:CZ	2.52	0.44
1:A:272:ILE:O	1:A:276:LEU:HB2	2.18	0.43
1:A:363:TYR:O	1:A:367:ILE:HD12	2.18	0.43
1:B:76:PHE:CD1	1:B:133:LYS:HG3	2.54	0.43
1:B:363:TYR:O	1:B:367:ILE:HG13	2.17	0.43
1:A:258:VAL:O	1:A:258:VAL:CG1	2.67	0.43
1:A:351:HIS:O	1:A:352:ASP:HB2	2.19	0.43
1:A:94:ARG:CD	3:A:501:HOH:O	2.65	0.43
1:A:266:ARG:HE	1:A:266:ARG:HB2	1.64	0.43
1:A:354:ILE:HD13	1:A:365:GLU:HG2	2.00	0.43
1:A:141:ILE:HG13	1:A:145:GLU:HB3	1.99	0.42
1:A:273:LYS:NZ	1:A:280:VAL:H	2.18	0.42
1:A:289:ALA:HB1	1:A:330:VAL:HG21	2.00	0.42
1:A:164:ARG:NE	1:A:231:GLU:OE1	2.53	0.42
1:B:28:LEU:HD11	1:B:63:VAL:HG22	2.00	0.42
1:B:317:ARG:HD3	1:B:348:GLY:HA3	2.01	0.42
1:A:169:VAL:HG21	1:A:176:ILE:HD13	2.02	0.42
1:B:276:LEU:CD1	1:B:367:ILE:HG23	2.49	0.42
1:A:129:ARG:NH1	1:B:235:ASP:HA	2.35	0.42
1:A:291:GLN:HG2	3:A:639:HOH:O	2.18	0.41
1:B:302:LYS:O	1:B:305:LYS:HE2	2.19	0.41
1:B:335:LEU:HD12	1:B:336:ILE:N	2.35	0.41
1:B:351:HIS:CE1	3:B:416:HOH:O	2.74	0.41
1:A:303:ASN:CG	1:A:304:GLY:N	2.74	0.41
1:A:354:ILE:HD12	1:A:366:VAL:HG22	2.02	0.41
1:A:13:ARG:NH1	3:A:513:HOH:O	2.54	0.41
1:A:291:GLN:HG3	1:A:292:TYR:CE2	2.56	0.41
1:B:43:LYS:HB2	1:B:43:LYS:HE2	1.69	0.40
1:B:175:ARG:HA	1:B:175:ARG:HD2	1.85	0.40
1:A:179:LEU:HD23	1:A:179:LEU:HA	1.95	0.40

All (3) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:642:HOH:O	3:B:410:HOH:O[1_655]	1.93	0.27

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:657:HOH:O	3:B:534:HOH:O[2_655]	2.00	0.20
3:A:658:HOH:O	3:B:596:HOH:O[2_655]	2.18	0.02

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	374/375 (100%)	362 (97%)	11 (3%)	1 (0%)	41 32
1	B	373/375 (100%)	361 (97%)	9 (2%)	3 (1%)	19 9
All	All	747/750 (100%)	723 (97%)	20 (3%)	4 (0%)	29 17

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	3	PRO
1	B	8	ASP
1	A	273	LYS
1	B	262	SER

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	337/336 (100%)	322 (96%)	15 (4%)	27 12
1	B	336/336 (100%)	323 (96%)	13 (4%)	32 17

Continued on next page...

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
All	All	673/672 (100%)	645 (96%)	28 (4%)	30	14

All (28) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	8	ASP
1	A	164	ARG
1	A	207	ARG
1	A	208	LYS
1	A	244	SER
1	A	255	GLU
1	A	257	SER
1	A	260	TYR
1	A	271	LYS
1	A	274	ARG
1	A	302	LYS
1	A	329	ASN
1	A	358	LYS
1	A	369	LYS
1	A	372	GLU
1	B	4	LYS
1	B	12	ARG
1	B	80	SER
1	B	115	ASN
1	B	125	MET
1	B	137	ARG
1	B	174	LYS
1	B	212	SER
1	B	253	LYS
1	B	259	GLU
1	B	262	SER
1	B	331	ASP
1	B	375	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	SO4	A	401	-	4,4,4	0.11	0	6,6,6	0.31	0

There are no bond length outliers.

There are no bond angle outliers.


There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight  $> 250$  and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



### 5.7 Other polymers [i](#)

There are no such residues in this entry.

### 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	375/375 (100%)	0.63	24 (6%) 19 26	23, 43, 82, 121	0
1	B	375/375 (100%)	0.66	30 (8%) 12 17	22, 44, 82, 109	0
All	All	750/750 (100%)	0.64	54 (7%) 15 21	22, 44, 82, 121	0

All (54) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	258	VAL	10.7
1	A	260	TYR	10.4
1	B	260	TYR	8.6
1	A	3	PRO	6.7
1	A	257	SER	6.3
1	B	258	VAL	6.2
1	B	261	ASP	6.1
1	A	255	GLU	6.1
1	B	256	VAL	5.1
1	B	3	PRO	4.9
1	A	256	VAL	4.8
1	A	2	LEU	4.6
1	A	259	GLU	4.6
1	B	7	LEU	4.4
1	A	349	LYS	4.3
1	B	259	GLU	4.3
1	B	357	ARG	4.2
1	B	2	LEU	4.2
1	A	254	LEU	4.0
1	A	261	ASP	4.0
1	B	257	SER	3.9
1	A	375	ASN	3.9
1	A	303	ASN	3.8
1	A	348	GLY	3.8

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	349	LYS	3.8
1	B	347	LEU	3.7
1	B	350	ILE	3.5
1	A	347	LEU	3.1
1	B	1	MET	3.0
1	B	262	SER	3.0
1	A	274	ARG	2.9
1	B	375	ASN	2.8
1	A	141	ILE	2.8
1	A	1	MET	2.8
1	B	255	GLU	2.8
1	B	264	LEU	2.7
1	B	286	ILE	2.7
1	B	174	LYS	2.7
1	B	351	HIS	2.7
1	B	263	SER	2.6
1	B	232	ILE	2.6
1	B	115	ASN	2.5
1	B	116	THR	2.5
1	A	6	LEU	2.4
1	B	285	GLY	2.3
1	B	4	LYS	2.3
1	A	331	ASP	2.2
1	B	16	ILE	2.2
1	B	233	ILE	2.2
1	A	60	TYR	2.2
1	A	357	ARG	2.1
1	A	266	ARG	2.0
1	B	241	PHE	2.0
1	A	53	HIS	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.




## 6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	SO4	A	401	5/5	0.99	0.10	30,30,35,37	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



## 6.5 Other polymers [i](#)

There are no such residues in this entry.