

Full wwPDB X-ray Structure Validation Report (i)

May 15, 2020 - 02:46 am BST

PDB ID	:	1GZL
Title	:	Crystal structure of C14linkmid/IQN17: a cross-linked inhibitor of HIV-1 en-
		try bound to the gp41 hydrophobic pocket
Authors	:	Sia, S.K.; Carr, P.A.; Cochran, A.G.; Malashkevich, V.M.; Kim, P.S.
Deposited on	:	2002-05-23
Resolution	:	1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (1)) were used in the production of this report:


MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.11
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	$7.0.044 (\mathrm{Gargrove})$
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$egin{array}{c} { m Whole \ archive} \ (\#{ m Entries}) \end{array}$	${f Similar\ resolution}\ (\#{ m Entries},{ m resolution\ range}({ m \AA}))$		
R_{free}	130704	5950 (1.80-1.80)		
Clashscore	141614	6793(1.80-1.80)		
Ramachandran outliers	138981	6697(1.80-1.80)		
Sidechain outliers	138945	6696 (1.80-1.80)		
RSRZ outliers	127900	5850 (1.80-1.80)		

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain					
1	А	46	4%	59%		37%	•	
1	В	46		59%		37%	•	
2	С	12	8%	{	83%		8%	
2	D	12	25%		75%			



2 Entry composition (i)

There are 5 unique types of molecules in this entry. The entry contains 1095 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called FUSION PROTEIN BETWEEN THE HYDROPHOBIC POCKET OF HIV GP41 AND GENERAL CONTROL PROTEIN GCN4-PIQI.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	Δ	46	Total	С	Ν	Ο	S	0	0	0
		40	384	245	69	69	1	0		
1	1 B	46	Total	С	Ν	Ο	S	0	0	0
		B 46		245	69	69	1	0	U	0

Chain	Residue	Modelled	Actual	Comment	Reference
A	5	ILE	LEU	$\operatorname{conflict}$	UNP P03069
A	9	ILE	VAL	conflict	UNP P03069
A	12	ILE	LEU	conflict	UNP P03069
A	13	GLU	LEU	conflict	UNP P03069
A	16	GLN	ASN	conflict	UNP P03069
A	17	LYS	TYR	conflict	UNP P03069
A	18	LYS	HIS	conflict	UNP P03069
A	19	ILE	LEU	conflict	UNP P03069
A	23	ILE	VAL	conflict	UNP P03069
A	26	ILE	LEU	conflict	UNP P03069
В	5	ILE	LEU	$\operatorname{conflict}$	UNP P03069
В	9	ILE	VAL	conflict	UNP P03069
В	12	ILE	LEU	conflict	UNP P03069
В	13	GLU	LEU	conflict	UNP P03069
В	16	GLN	ASN	conflict	UNP P03069
В	17	LYS	TYR	conflict	UNP P03069
В	18	LYS	HIS	conflict	UNP P03069
В	19	ILE	LEU	conflict	UNP P03069
В	23	ILE	VAL	conflict	UNP P03069
В	26	ILE	LEU	$\operatorname{conflict}$	UNP P03069

There are 20 discrepancies between the modelled and reference sequences:

• Molecule 2 is a protein called ENVELOPE GLYCOPROTEIN GP41.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace				
0	2 C	12	Total	С	Ν	0	0	0	0				
		12	117	75	18	24							
0	2 D	Л	Л	П	Л	19	Total	С	Ν	0	0	0	0
		12	117	75	18	24	0	0	0				


There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
С	629	GLU	MET	engineered mutation	UNP P04578
С	636	GLU	ASN	engineered mutation	UNP P04578
D	629	GLU	MET	engineered mutation	UNP P04578
D	636	GLU	ASN	engineered mutation	UNP P04578

• Molecule 3 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	В	1	Total Cl 1 1	0	0
3	А	1	Total Cl 1 1	0	0

• Molecule 4 is PENTANE-1,5-DIAMINE (three-letter code: N2P) (formula: C₅H₁₄N₂).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	С	1	Total C 7 5	N 2	0	0



Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	D	1	TotalCN752	0	0

• Molecule 5 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	А	30	$\begin{array}{cc} {\rm Total} & {\rm O} \\ {\rm 30} & {\rm 30} \end{array}$	0	0
5	В	32	$\begin{array}{cc} \text{Total} & \text{O} \\ 32 & 32 \end{array}$	0	0
5	С	7	Total O 7 7	0	0
5	D	8	Total O 8 8	0	0




3 Residue-property plots (i)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: FUSION PROTEIN BETWEEN THE HYDROPHOBIC POCKET OF HIV GP41 AND GENERAL CONTROL PROTEIN GCN4-PIQI



• Molecule 1: FUSION PROTEIN BETWEEN THE HYDROPHOBIC POCKET OF HIV GP41 AND GENERAL CONTROL PROTEIN GCN4-PIQI





4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 63	Depositor
Cell constants	38.36Å 38.36 Å 169.69 Å	Deperitor
a, b, c, α , β , γ	90.00° 90.00° 120.00°	Depositor
Resolution (Å)	19.58 - 1.80	Depositor
Resolution (A)	19.58 - 1.80	EDS
% Data completeness	81.7(19.58-1.80)	Depositor
(in resolution range)	$81.8\ (19.58\text{-}1.80)$	EDS
R _{merge}	(Not available)	Depositor
R _{sym}	0.04	Depositor
$< I/\sigma(I) > 1$	$3.80 ({\rm at}1.80{ m \AA})$	Xtriage
Refinement program	CNS 1.0	Depositor
R, R_{free}	0.208 , 0.243	Depositor
$\mathbf{n}, \mathbf{n}_{free}$	0.205 , 0.248	DCC
R_{free} test set	685 reflections $(6.42%)$	wwPDB-VP
Wilson B-factor $(Å^2)$	25.1	Xtriage
Anisotropy	0.224	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.42 , 84.9	EDS
L-test for twinning ²	$< L >=0.51, < L^2>=0.34$	Xtriage
Estimated twinning fraction	0.257 for h,-h-k,-l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	1095	wwPDB-VP
Average B, all atoms $(Å^2)$	35.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 12.04% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: N2P, ACE, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
	Chain	RMSZ	# Z > 5	RMSZ	# Z > 5
1	А	0.38	0/383	0.55	0/506
1	В	0.40	0/383	0.50	0/506
2	С	0.53	0/121	0.58	0/162
2	D	0.58	0/121	0.49	0/162
All	All	0.43	0/1008	0.53	0/1336

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	384	0	428	20	0
1	В	384	0	428	23	0
2	С	117	0	93	11	0
2	D	117	0	93	13	0
3	А	1	0	0	0	0
3	В	1	0	0	0	0
4	С	7	0	10	1	0
4	D	7	0	11	5	0
5	А	30	0	0	4	0



001000							
Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes	
5	В	32	0	0	3	0	
5	С	7	0	0	1	0	
5	D	8	0	0	1	0	
All	All	1095	0	1063	60	0	

Continued from previous page...

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 29.

All (60) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic	Clash
		distance (Å)	overlap (Å)
1:A:4:GLN:HA	1:A:4:GLN:HE21	1.02	1.09
1:B:44:ILE:HG22	1:B:45:LEU:HG	1.31	1.07
2:D:632:ASP:OD1	4:D:1640:N2P:H2C1	1.67	0.94
1:A:4:GLN:HA	1:A:4:GLN:NE2	1.84	0.92
1:A:15:LYS:HG3	5:A:2017:HOH:O	1.73	0.87
1:B:28:LYS:HE3	1:B:28:LYS:HA	1.61	0.82
1:B:40:LEU:O	1:B:44:ILE:HG12	1.87	0.74
1:B:43:ARG:HH21	2:D:628:TRP:HE1	1.36	0.72
1:B:28:LYS:HG2	2:D:638:TYR:CZ	2.26	0.71
1:B:44:ILE:CG2	1:B:45:LEU:HG	2.14	0.70
2:C:631:TRP:O	2:C:635:ILE:HG12	1.92	0.69
1:B:25:ARG:HG2	5:B:2023:HOH:O	1.93	0.69
1:B:16:GLN:HG2	1:B:20:GLU:OE2	1.94	0.68
1:B:28:LYS:HG2	2:D:638:TYR:CE2	2.29	0.68
1:A:16:GLN:O	1:A:20:GLU:HG3	1.93	0.68
2:D:633:ARG:HH22	4:D:1640:N2P:H1C1	1.56	0.67
1:B:1:ARG:HG3	5:B:2005:HOH:O	1.97	0.64
1:A:33:THR:O	1:A:37:ILE:HG13	1.98	0.64
1:B:3:LYS:HE2	1:B:7:ASP:OD1	2.00	0.62
1:B:44:ILE:O	1:B:45:LEU:HD23	2.02	0.60
1:A:4:GLN:HE21	1:A:4:GLN:CA	1.92	0.59
1:B:25:ARG:HG3	2:D:638:TYR:HD1	1.67	0.59
2:C:632:ASP:OD2	4:C:1640:N2P:H1C2	2.03	0.59
1:A:41:GLN:HG2	1:A:45:LEU:OXT	2.03	0.58
2:D:633:ARG:HH21	4:D:1640:N2P:H4C2	1.67	0.58
1:B:0:ACE:O	1:B:4:GLN:HG3	2.04	0.57
5:A:2026:HOH:O	2:C:635:ILE:HD12	2.04	0.57
2:D:633:ARG:NH2	4:D:1640:N2P:H1C1	2.19	0.56
1:A:44:ILE:O	1:A:45:LEU:HB2	2.07	0.54
2:C:629:GLU:O	2:C:633:ARG:HG3	2.09	0.53



A 4 1	A 4 5 55 0	Interatomic	Clash
Atom-1	Atom-2	$distance (m \AA)$	overlap (Å)
2:C:639:THR:HG22	2:C:639:THR:OXT	2.08	0.53
1:A:18:LYS:HD2	5:A:2024:HOH:O	2.09	0.52
2:D:639:THR:HG21	5:D:2001:HOH:O	2.10	0.50
1:A:44:ILE:HG22	1:A:45:LEU:N	2.27	0.50
1:B:32:LEU:HD11	2:D:634:GLU:HB3	1.93	0.50
1:A:30:LEU:O	1:A:34:VAL:HG23	2.12	0.49
1:B:43:ARG:HH11	1:B:43:ARG:HG2	1.80	0.47
1:B:44:ILE:HG22	1:B:45:LEU:N	2.28	0.47
2:D:628:TRP:O	2:D:631:TRP:HB3	2.14	0.47
2:D:631:TRP:O	2:D:635:ILE:HG12	2.15	0.47
2:C:628:TRP:N	5:C:2001:HOH:O	2.49	0.46
1:B:44:ILE:HG22	1:B:45:LEU:CG	2.23	0.46
1:B:4:GLN:O	1:B:7:ASP:HB2	2.15	0.46
2:C:636:GLU:HG2	2:C:636:GLU:H	1.42	0.45
1:B:39:GLN:O	1:B:43:ARG:HG3	2.16	0.45
1:A:25:ARG:O	1:A:29:LEU:HB2	2.17	0.44
1:B:35:TRP:CH2	1:B:39:GLN:HG3	2.52	0.44
1:A:29:LEU:O	1:A:33:THR:HG23	2.18	0.44
1:A:2:MET:HB2	5:A:2001:HOH:O	2.16	0.44
2:C:630:GLU:HG2	2:C:633:ARG:HH12	1.83	0.43
1:A:28:LYS:HD3	1:A:28:LYS:O	2.19	0.43
1:A:41:GLN:O	1:A:45:LEU:N	2.52	0.43
1:B:9:ILE:HD12	5:B:2001:HOH:O	2.18	0.43
2:D:633:ARG:NH2	4:D:1640:N2P:C1	2.82	0.42
2:C:630:GLU:O	2:C:634:GLU:HG2	2.20	0.42
1:A:32:LEU:HD11	2:C:634:GLU:HB3	2.01	0.42
1:A:29:LEU:HD12	1:A:29:LEU:HA	1.94	0.42
1:A:29:LEU:HD13	2:C:638:TYR:HB2	2.01	0.41
1:A:38:LYS:NZ	1:A:38:LYS:HB3	2.36	0.41
1:B:35:TRP:CZ2	1:B:39:GLN:HG3	2.56	0.40

Continued from previous page...

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.



Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	А	44/46~(96%)	42 (96%)	1 (2%)	1 (2%)	6	1
1	В	44/46~(96%)	43~(98%)	0	1 (2%)	6	1
2	С	10/12~(83%)	10 (100%)	0	0	100	100
2	D	10/12~(83%)	10 (100%)	0	0	100	100
All	All	108/116~(93%)	105~(97%)	1 (1%)	2(2%)	8	1

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	В	44	ILE
1	А	44	ILE

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percer	ntiles
1	А	42/42~(100%)	40~(95%)	2~(5%)	25	11
1	В	42/42~(100%)	39~(93%)	3~(7%)	14	5
2	С	12/12~(100%)	11 (92%)	1 (8%)	11	3
2	D	12/12~(100%)	11 (92%)	1 (8%)	11	3
All	All	108/108~(100%)	101 (94%)	7 (6%)	17	6

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	А	4	GLN
1	А	14	SER
1	В	28	LYS
1	В	30	LEU
1	В	41	GLN
2	С	636	GLU



Continued from previous page...

Mol	Chain	\mathbf{Res}	Type
2	D	630	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	А	4	GLN
1	В	4	GLN
2	С	637	ASN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no carbohydrates in this entry.

5.6 Ligand geometry (i)

Of 4 ligands modelled in this entry, 2 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol Type Chain		Chain	ain Res	Link	B	Bond lengths			Bond angles		
INIOI	туре	Unam	nes		Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2	
4	N2P	С	1640	2	6,6,6	0.47	0	$5,\!5,\!5$	0.47	0	
4	N2P	D	1640	2	6,6,6	0.71	0	$5,\!5,\!5$	0.27	0	

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the



Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	\mathbf{Res}	Link	Chirals	Torsions	Rings
4	N2P	С	1640	2	-	1/4/4/4	-
4	N2P	D	1640	2	-	1/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	С	1640	N2P	C1-C2-C3-C4
4	D	1640	N2P	C1-C2-C3-C4

There are no ring outliers.

2 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	С	1640	N2P	1	0
4	D	1640	N2P	5	0

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ $>$	# RSRZ > 2	$OWAB(Å^2)$	Q<0.9
1	А	45/46~(97%)	-0.27	2 (4%) 34 28	17, 27, 68, 81	0
1	В	45/46~(97%)	-0.34	0 100 100	18, 25, 41, 71	0
2	С	12/12~(100%)	0.17	0 100 100	45, 50, 59, 62	0
2	D	12/12~(100%)	0.18	0 100 100	45, 49, 55, 56	0
All	All	114/116~(98%)	-0.21	2 (1%) 68 64	17, 28, 62, 81	0

All (2) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	А	44	ILE	2.7
1	А	43	ARG	2.7

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no carbohydrates in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.



Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-factors}(\mathrm{\AA}^2)$	Q<0.9
4	N2P	D	1640	7/7	0.69	0.31	$60,\!65,\!66,\!66$	0
4	N2P	С	1640	7/7	0.76	0.32	57,60,62,62	0
3	CL	В	1046	1/1	0.86	0.18	$36,\!36,\!36,\!36$	1
3	CL	А	1046	1/1	0.99	0.17	$38,\!38,\!38,\!38$	1

6.5 Other polymers (i)

There are no such residues in this entry.

