

# Full wwPDB X-ray Structure Validation Report (i)

#### May 23, 2020 – 05:31 am BST

PDB ID : 4H2X

Title : Crystal structure of engineered Bradyrhizobium japonicum glycine: [carrier

protein ligase complexed with carrier protein from Agrobacterium tumefa-

ciens and an analogue of glycyl adenylate

Authors : Luic, M.; Weygand-Durasevic, I.; Ivic, N.; Mocibob, M.

Deposited on : 2012-09-13

Resolution : 2.15 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity : 4.02b-467

Mogul: 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : 1.13

EDS : 2.11

buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

 $Refmac \quad : \quad 5.8.0158$ 

CCP4 : 7.0.044 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.11

## 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X- $RAY\ DIFFRACTION$ 

The reported resolution of this entry is 2.15 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive	Similar resolution
Metric	$(\# { m Entries})$	$(\#  ext{Entries},  ext{resolution range}( ext{Å}))$
$R_{free}$	130704	1479 (2.16-2.16)
Clashscore	141614	1585 (2.16-2.16)
Ramachandran outliers	138981	1560 (2.16-2.16)
Sidechain outliers	138945	1559 (2.16-2.16)
RSRZ outliers	127900	1456 (2.16-2.16)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain					
1	A	346	81%		• 15%			
2	В	346	81%		• 14%			
3	С	103	39% 65%	10%	25%			
3	D	103	61% 66%	7%	27%			



# 2 Entry composition (i)

There are 8 unique types of molecules in this entry. The entry contains 6243 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Amino acid--[acyl-carrier-protein] ligase 1.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace		
1	Λ	295	Total	С	N	О	S	0	1.4	0
1	Λ.	∠90 	2358	1488	418	434	18	0	14	0

There are 20 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-19	MET	-	EXPRESSION TAG	UNP Q89VT8
A	-18	GLY	_	EXPRESSION TAG	UNP Q89VT8
A	-17	SER	=	EXPRESSION TAG	UNP Q89VT8
A	-16	SER	-	EXPRESSION TAG	UNP Q89VT8
A	-15	HIS	_	EXPRESSION TAG	UNP Q89VT8
A	-14	HIS	-	EXPRESSION TAG	UNP Q89VT8
A	-13	HIS	-	EXPRESSION TAG	UNP Q89VT8
A	-12	HIS	-	EXPRESSION TAG	UNP Q89VT8
A	-11	HIS	-	EXPRESSION TAG	UNP Q89VT8
A	-10	HIS	_	EXPRESSION TAG	UNP Q89VT8
A	-9	SER	-	EXPRESSION TAG	UNP Q89VT8
A	-8	SER	-	EXPRESSION TAG	UNP Q89VT8
A	-7	GLY	-	EXPRESSION TAG	UNP Q89VT8
A	-6	LEU	-	EXPRESSION TAG	UNP Q89VT8
A	-5	VAL	_	EXPRESSION TAG	UNP Q89VT8
A	-4	PRO	=	EXPRESSION TAG	UNP Q89VT8
A	-3	ARG	-	EXPRESSION TAG	UNP Q89VT8
A	-2	GLY	-	EXPRESSION TAG	UNP Q89VT8
A	-1	SER	-	EXPRESSION TAG	UNP Q89VT8
A	0	HIS	-	EXPRESSION TAG	UNP Q89VT8

• Molecule 2 is a protein called Amino acid--[acyl-carrier-protein] ligase 1.

Mol	Chain	Residues		At	oms			ZeroOcc	AltConf	Trace
2	В	296	Total	С	N	0	S	0	14	0
			2373	1496	427	432	18	_		_



There are 20 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
В	-19	MET	-	EXPRESSION TAG	UNP Q89VT8
В	-18	GLY	-	EXPRESSION TAG	UNP Q89VT8
В	-17	SER	=	EXPRESSION TAG	UNP Q89VT8
В	-16	SER	-	EXPRESSION TAG	UNP Q89VT8
В	-15	HIS	-	EXPRESSION TAG	UNP Q89VT8
В	-14	HIS	-	EXPRESSION TAG	UNP Q89VT8
В	-13	HIS	-	EXPRESSION TAG	UNP Q89VT8
В	-12	HIS	-	EXPRESSION TAG	UNP Q89VT8
В	-11	HIS	=	EXPRESSION TAG	UNP Q89VT8
В	-10	HIS	-	EXPRESSION TAG	UNP Q89VT8
В	-9	SER	-	EXPRESSION TAG	UNP Q89VT8
В	-8	SER	_	EXPRESSION TAG	UNP Q89VT8
В	-7	GLY	-	EXPRESSION TAG	UNP Q89VT8
В	-6	LEU	_	EXPRESSION TAG	UNP Q89VT8
В	-5	VAL	-	EXPRESSION TAG	UNP Q89VT8
В	-4	PRO	_	EXPRESSION TAG	UNP Q89VT8
В	-3	ARG	_	EXPRESSION TAG	UNP Q89VT8
В	-2	GLY	-	EXPRESSION TAG	UNP Q89VT8
В	-1	SER	-	EXPRESSION TAG	UNP Q89VT8
В	0	HIS	-	EXPRESSION TAG	UNP Q89VT8

• Molecule 3 is a protein called Aminoacyl carrier protein.

Mol	Chain	Residues	${f Atoms}$			ZeroOcc	AltConf	Trace	
3	С	77	Total 568	C 362		S 2	0	0	0
3	D	75	Total 498	C 319		S 1	0	0	0

There are 40 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference	
С	-19	MET	_	EXPRESSION TAG	UNP A9CHM9	
С	-18	GLY	_	EXPRESSION TAG	UNP A9CHM9	
С	-17	SER	-	EXPRESSION TAG	UNP A9CHM9	
С	-16	SER	_	EXPRESSION TAG	UNP A9CHM9	
С	-15	HIS	-	EXPRESSION TAG	UNP A9CHM9	
С	-14	HIS	-	EXPRESSION TAG	UNP A9CHM9	
С	-13	HIS	-	EXPRESSION TAG	UNP A9CHM9	
С	-12	HIS	_	EXPRESSION TAG	UNP A9CHM9	
С	-11	HIS	-	EXPRESSION TAG	UNP A9CHM9	




 $Continued\ from\ previous\ page...$ 

		Modelled	Actual	Comment	Reference
С	-10	HIS	_	EXPRESSION TAG	UNP A9CHM9
С	-9	SER	_	EXPRESSION TAG	UNP A9CHM9
С	-8	SER	_	EXPRESSION TAG	UNP A9CHM9
С	-7	GLY	_	EXPRESSION TAG	UNP A9CHM9
С	-6	LEU	_	EXPRESSION TAG	UNP A9CHM9
С	-5	VAL	_	EXPRESSION TAG	UNP A9CHM9
С	-4	PRO	_	EXPRESSION TAG	UNP A9CHM9
С	-3	ARG	-	EXPRESSION TAG	UNP A9CHM9
С	-2	GLY	-	EXPRESSION TAG	UNP A9CHM9
С	-1	SER	_	EXPRESSION TAG	UNP A9CHM9
С	0	HIS	-	EXPRESSION TAG	UNP A9CHM9
D	-19	MET	-	EXPRESSION TAG	UNP A9CHM9
D	-18	GLY	_	EXPRESSION TAG	UNP A9CHM9
D	-17	SER	-	EXPRESSION TAG	UNP A9CHM9
D	-16	SER	_	EXPRESSION TAG	UNP A9CHM9
D	-15	HIS	-	EXPRESSION TAG	UNP A9CHM9
D	-14	HIS	_	EXPRESSION TAG	UNP A9CHM9
D	-13	HIS	_	EXPRESSION TAG	UNP A9CHM9
D	-12	HIS	_	EXPRESSION TAG	UNP A9CHM9
D	-11	HIS	_	EXPRESSION TAG	UNP A9CHM9
D	-10	HIS	-	EXPRESSION TAG	UNP A9CHM9
D	-9	SER	_	EXPRESSION TAG	UNP A9CHM9
D	-8	SER	_	EXPRESSION TAG	UNP A9CHM9
D	-7	GLY	_	EXPRESSION TAG	UNP A9CHM9
D	-6	LEU	_	EXPRESSION TAG	UNP A9CHM9
D	-5	VAL	-	EXPRESSION TAG	UNP A9CHM9
D	-4	PRO	-	EXPRESSION TAG	UNP A9CHM9
D	-3	ARG	-	EXPRESSION TAG	UNP A9CHM9
D	-2	GLY	-	EXPRESSION TAG	UNP A9CHM9
D	-1	SER	-	EXPRESSION TAG	UNP A9CHM9
D	0	HIS	-	EXPRESSION TAG	UNP A9CHM9

 $\bullet$  Molecule 4 is 5'-O-(glycylsulfamoyl) adenosine (three-letter code: G5A) (formula:  $\rm C_{12}H_{17}N_7O_7S).$ 





Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
4	٨	1	Total	С	N	О	S	0	0
$\begin{vmatrix} 4 & A \end{vmatrix}$	A	1	27	12	7	7	1	0	0
1	D	1	Total	С	N	О	S	0	0
4	R I	27	12	7	7	1	U		

• Molecule 5 is CHLORIDE ION (three-letter code: CL) (formula: Cl).


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	В	1	Total Cl 1 1	0	0
5	A	1	Total Cl 1 1	0	0

• Molecule 6 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	${f Atoms}$	ZeroOcc	${f AltConf}$
6	В	1	Total Zn 1 1	0	0

 $\bullet \ \ \mathrm{Molecule} \ 7 \ \mathrm{is} \ 4'\mathrm{-PHOSPHOPANTETHEINE} \ (\mathrm{three-letter} \ \mathrm{code} \colon \mathrm{PNS}) \ (\mathrm{formula} \colon \mathrm{C}_{11}\mathrm{H}_{23}\mathrm{N}_2\mathrm{O}_7\mathrm{PS}).$ 





Mol	Chain	Residues		A	tom	ıs			ZeroOcc	AltConf
7	C	1	Total	С	N	О	Р	S	0 0	
'		1	21	11	2	6	1	1	0	
7	D	1	Total	С	N	О	Р	S	0	0
'	ש	1	21	11	2	6	1	1	0	

#### • Molecule 8 is water.


Mol	Chain	Residues	${f Atoms}$	ZeroOcc	${f AltConf}$
8	A	160	Total O 160 160	0	0
8	В	182	Total O 182 182	0	0
8	С	4	Total O 4 4	0	0
8	D	1	Total O 1 1	0	0



# 3 Residue-property plots (i)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Amino acid--[acyl-carrier-protein] ligase 1







# 4 Data and refinement statistics (i)

Property	Value	Source		
Space group	P 21 21 21	Depositor		
Cell constants	99.58Å 101.43Å 103.04Å	Danasitan		
a, b, c, $\alpha$ , $\beta$ , $\gamma$	$90.00^{\circ}$ $90.00^{\circ}$ $90.00^{\circ}$	Depositor		
Resolution (Å)	45.94 - 2.15	Depositor		
Resolution (A)	45.94 - 2.15	EDS		
% Data completeness	99.7 (45.94-2.15)	Depositor		
(in resolution range)	99.7 (45.94-2.15)	EDS		
$R_{merge}$	0.06	Depositor		
$R_{sum}$	(Not available)	Depositor		
$< I/\sigma(I) > 1$	1.96 (at 2.14Å)	Xtriage		
Refinement program	PHENIX dev_1116	Depositor		
D D.	0.167 , 0.198	Depositor		
$R, R_{free}$	0.168 , $0.198$	DCC		
$R_{free}$ test set	2879 reflections $(5.00\%)$	wwPDB-VP		
Wilson B-factor (Å <sup>2</sup> )	34.9	Xtriage		
Anisotropy	0.359	Xtriage		
Bulk solvent $k_{sol}(e/Å^3)$ , $B_{sol}(Å^2)$	$0.32\;,45.2$	EDS		
L-test for twinning <sup>2</sup>	$< L >=0.49, < L^2>=0.32$	Xtriage		
	0.013 for -h,l,k			
	0.012  for -l,-k,-h			
Estimated twinning fraction	0.016 for k,h,-l	Xtriage		
	$0.005  ext{ for }  ext{k,l,h}$			
	0.005 for l,h,k			
$F_o, F_c$ correlation	0.96	EDS		
Total number of atoms	6243	wwPDB-VP		
Average B, all atoms (Å <sup>2</sup> )	41.0	wwPDB-VP		

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 4.03% of the height of the origin peak. No significant pseudotranslation is detected.

<sup>&</sup>lt;sup>2</sup>Theoretical values of <|L|>,  $< L^2>$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



<sup>&</sup>lt;sup>1</sup>Intensities estimated from amplitudes.

# 5 Model quality (i)

## 5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: PNS, CSO, G5A, ZN, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond angles		
IVIOI	Chain	RMSZ	# Z  > 5	RMSZ	# Z  > 5	
1	A	0.41	0/2442	0.54	0/3308	
2	В	0.41	0/2470	0.55	0/3348	
3	С	0.29	0/577	0.45	0/785	
3	D	0.25	0/506	0.42	0/693	
All	All	0.39	0/5995	0.53	0/8134	

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	$\mathbf{H}(\mathbf{added})$	Clashes	Symm-Clashes
1	A	2358	0	2305	9	0
2	В	2373	0	2319	9	0
3	С	568	0	534	8	0
3	D	498	0	429	4	0
4	A	27	0	17	2	0
4	В	27	0	16	1	0
5	A	1	0	0	1	0
5	В	1	0	0	1	0
6	В	1	0	0	0	0



Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
7	С	21	0	21	0	0
7	D	21	0	21	0	0
8	A	160	0	0	0	0
8	В	182	0	0	0	0
8	С	4	0	0	0	0
8	D	1	0	0	0	0
All	All	6243	0	5662	26	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (26) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	$\begin{array}{c} {\rm Interatomic} \\ {\rm distance} \ ({\rm \AA}) \end{array}$	Clash overlap (Å)
1:A:220:ARG:NH2	3:D:53:PHE:O	2.22	0.71
2:B:44[A]:GLU:OE2	2:B:173:ARG:NH1	2.25	0.69
1:A:126[B]:LEU:HD21	2:B:126[B]:LEU:HD21	1.76	0.67
2:B:224:MET:HB2	3:C:58:LEU:HD22	1.81	0.61
1:A:268[A]:GLN:NE2	1:A:272:GLY:O	2.35	0.59
1:A:131:CSO:OD	4:A:401:G5A:N	2.41	0.54
3:D:6:ARG:NH1	3:D:19:VAL:O	2.42	0.53
3:C:42:MET:CE	3:C:58:LEU:HD21	2.39	0.52
3:C:42:MET:HE3	3:C:58:LEU:HD21	1.92	0.51
3:D:24:ASP:HA	3:D:66:ILE:HB	1.94	0.50
1:A:243:ARG:HB3	1:A:243:ARG:CZ	2.42	0.50
1:A:49[B]:ARG:HD3	1:A:310:LEU:O	2.12	0.48
3:C:8:ILE:HA	3:C:11:LYS:HB2	1.95	0.47
3:C:57:LEU:HD13	3:C:72:THR:HG23	1.97	0.47
3:D:41:LEU:O	3:D:45:ILE:HG12	2.16	0.46
2:B:220:ARG:NH1	3:C:46:GLU:OE2	2.46	0.46
4:A:401:G5A:N3	5:A:402:CL:CL	2.86	0.46
1:A:91[B]:CYS:HG	2:B:115:TRP:HH2	1.63	0.46
2:B:232:GLN:OE1	2:B:258[A]:ARG:NH2	2.51	0.43
2:B:93[B]:CYS:SG	2:B:119:LEU:HD13	2.58	0.43
1:A:115:TRP:HH2	2:B:91[B]:CYS:SG	2.42	0.42
3:C:29:TYR:CE2	3:C:64:ALA:HA	2.55	0.42
4:B:402:G5A:N3	5:B:403:CL:CL	2.90	0.42
2:B:18:ASP:HA	2:B:19:PRO:HD3	1.97	0.40
1:A:76:LEU:HD22	1:A:125:VAL:HG21	2.04	0.40
3:C:8:ILE:HD11	3:C:48:ALA:HB3	2.04	0.40



There are no symmetry-related clashes.

### 5.3 Torsion angles (i)

#### 5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Analysed Favoured Allowed Outliers		Outliers	Percentiles		
1	A	305/346~(88%)	303 (99%)	2 (1%)	0	100	100	
2	В	308/346 (89%)	304 (99%)	4 (1%)	0	100	100	
3	С	75/103 (73%)	71 (95%)	4 (5%)	0	100	100	
3	D	73/103 (71%)	70 (96%)	3 (4%)	0	100	100	
All	All	761/898 (85%)	748 (98%)	13 (2%)	0	100	100	

There are no Ramachandran outliers to report.

#### 5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percei	ntiles
1	A	$248/277 \ (90\%)$	245 (99%)	3 (1%)	71	76
2	В	$250/279 \ (90\%)$	247 (99%)	3 (1%)	71	76
3	С	56/84 (67%)	56 (100%)	0	100	100
3	D	38/84 (45%)	38 (100%)	0	100	100
All	All	592/724~(82%)	586 (99%)	6 (1%)	73	81

All (6) residues with a non-rotameric sidechain are listed below:



Mol	Chain	Res	Type
1	A	134	VAL
1	A	173	ARG
1	A	243	ARG
2	В	134	VAL
2	В	156	ASP
2	В	173	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

#### 5.3.3 RNA (i)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains (i)

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Tuno	e Chain	Pog	Tinle	B	ond leng	${ m gths}$	Bond angles		
MIOI	Type		Res	Link	Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
1	CSO	A	131	1	3,6,7	0.22	0	0,6,8	0.00	_
1	CSO	A	279	1	3,6,7	0.51	0	0,6,8	0.00	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

$\mathbf{Mol}$	Type	Chain	${f Res}$	Link	Chirals	Torsions	Rings
1	CSO	A	131	1	-	0/1/5/7	-
1	CSO	A	279	1	ı	0/1/5/7	1

There are no bond length outliers.

There are no bond angle outliers.



There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	Α	131	CSO	1	0

## 5.5 Carbohydrates (i)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry (i)

Of 7 ligands modelled in this entry, 3 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type Chain Res		Link	Bond lengths			Bond angles			
MIOI	туре	Chain	nes	LIIIK	Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
7	PNS	С	101	3	14,20,21	0.18	0	18,26,29	0.57	0
4	G5A	В	402	6	25,29,29	0.93	1 (4%)	26,43,43	1.21	4 (15%)
4	G5A	A	401	-	25,29,29	1.24	2 (8%)	26,43,43	1.23	4 (15%)
7	PNS	D	101	3	14,20,21	0.21	0	18,26,29	0.46	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	${f Res}$	Link	Chirals	Torsions	Rings
7	PNS	С	101	3	-	1/24/26/27	-
4	G5A	В	402	6	-	3/12/33/33	0/3/3/3
4	G5A	A	401	-	-	3/12/33/33	0/3/3/3
7	PNS	D	101	3	-	0/24/26/27	-



All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	$\mathbf{Z}$	${ m Observed}({ m \AA})$	$\mathbf{Ideal}(\mathbf{\AA})$
4	A	401	G5A	S-N3S	4.75	1.68	1.59
4	В	402	G5A	S-N3S	2.88	1.65	1.59
4	A	401	G5A	C-N3S	2.43	1.43	1.38

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	$\mathbf{Z}$	$Observed(^o)$	$\operatorname{Ideal}({}^{o})$
4	В	402	G5A	O-C-N3S	-3.22	117.92	122.08
4	В	402	G5A	C5'-O5'-S	3.07	123.71	117.37
4	A	401	G5A	O-C-N3S	-2.88	118.37	122.08
4	A	401	G5A	C5'-O5'-S	2.57	122.69	117.37
4	В	402	G5A	O2S-S-O1S	2.31	124.36	120.76
4	В	402	G5A	C5-C6-N6	2.21	123.71	120.35
4	A	401	G5A	C5-C6-N6	2.17	123.64	120.35
4	A	401	G5A	C-N3S-S	2.09	127.68	124.30

There are no chirality outliers.

All (7) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	В	402	G5A	C5'-O5'-S-N3S
4	A	401	G5A	C5'-O5'-S-N3S
4	В	402	G5A	C5'-O5'-S-O1S
4	A	401	G5A	C5'-O5'-S-O1S
4	A	401	G5A	C5'-O5'-S-O2S
7	С	101	PNS	O27-C28-C29-C31
4	В	402	G5A	C5'-O5'-S-O2S

There are no ring outliers.


2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	В	402	G5A	1	0
4	A	401	G5A	2	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier.



Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







# 5.7 Other polymers (i)

There are no such residues in this entry.

## 5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



## 6 Fit of model and data (i)

### 6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median,  $95^{th}$  percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ $>$	$\#\mathrm{RSRZ}{>}2$	$OWAB(A^2)$	Q < 0.9
1	A	293/346 (84%)	0.01	2 (0%) 87 91	17, 28, 55, 77	0
2	В	296/346~(85%)	-0.16	1 (0%) 94 95	18, 30, 47, 76	0
3	С	77/103 (74%)	2.35	40 (51%) 0 0	51, 68, 91, 102	0
3	D	75/103 (72%)	3.98	63 (84%) 0 0	62, 99, 114, 122	0
All	All	741/898 (82%)	0.59	106 (14%) 2 3	17, 33, 99, 122	0

All (106) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	D	45	ILE	10.7
3	D	48	ALA	8.4
3	D	17	THR	7.7
2	В	17	ALA	7.7
3	D	73	VAL	7.6
3	D	26	ALA	7.6
3	D	20	ASP	7.3
3	D	51	ILE	7.3
3	D	31	ALA	7.1
3	D	3	ALA	6.9
3	С	48	ALA	6.2
3	D	68	ALA	6.2
3	D	69	ILE	6.1
3	D	49	PHE	6.0
3	D	32	GLY	5.9
3	D	63	PHE	5.8
3	С	17	THR	5.8
3	D	23	ALA	5.7
3	D	74	LYS	5.6
3	D	75	LEU	5.5
3	С	73	VAL	5.5



Continued from previous page...

Continued from previous page								
Mol	Chain	Res	Type	RSRZ				
3	С	16	PRO	5.4				
3	D	27	ASP	5.2				
3	D	66	ILE	5.2				
3	D	53	PHE	5.1				
3	D	64	ALA	5.1				
3	D	71	ASP	5.1				
3	D	54	PRO	5.0				
3	С	75	LEU	4.8				
3	D	61	LYS	4.7				
3	С	57	LEU	4.6				
3	D	50	ASP	4.6				
3	D	39	VAL	4.5				
3	С	21	THR	4.3				
3	С	54	PRO	4.3				
3	D	28	LEU	4.3				
3	D	72	THR	4.3				
3	С	76	ILE	4.2				
3	D	29	TYR	4.2				
3	С	72	THR	4.2				
3	D	65	SER	4.2				
3	D	47	GLU	4.1				
3	С	18	PRO	4.1				
3	D	43	LEU	4.0				
3	D	24	ASP	4.0				
3	D	55	ASP	4.0				
3	D	33	LEU	4.0				
3	С	25	GLU	4.0				
3	D	4	THR	4.0				
3	С	55	ASP	4.0				
3	С	19	VAL	4.0				
3	D	19	VAL	3.9				
3	D	1	MET	3.8				
3	D	70	GLU	3.7				
3	D C	56	ASN	3.7				
3	С	68	ALA	3.7				
3	D	8	ILE	3.6				
3	D	18	PRO	3.5				
3	D	5	ILE	3.5				
3	D	52	GLU	3.5				
3	С	36	PHE	3.4				
3	С	15	LEU	3.4				
3	D	30	ALA	3.4				



Continued from previous page...

Mol	nued from Chain	Res	Type	RSRZ
3	D	25	GLU	3.4
3	С	26	ALA	3.4
3	С	63	PHE	3.3
3	С	74	LYS	3.3
3	С	22	ILE	3.2
3	D	56	ASN	3.2
3	D	57	LEU	3.1
3	С	53	PHE	3.1
3	С	50	ASP	3.1
3	D	16	PRO	3.1
3	D	22	ILE	3.1
3	С	12	PHE	3.1
3	D	12	PHE	3.0
3	D	11	LYS	3.0
3	С	65	SER	3.0
3	С	14	GLN	3.0
3	С	69	ILE	2.9
3	С	20	ASP	2.9
3	D	44	GLY	2.8
3	D	36	PHE	2.7
3	D	46	GLU	2.7
3	С	31	ALA	2.7
3	D	10	ALA	2.7
3	D	13	GLY	2.7
3	D	62	SER	2.7
3	С	44	GLY	2.6
3	D	41	LEU	2.6
1	A	212[A]	TYR	2.6
3	D	58	LEU	2.4
3	С	28	LEU	2.4
3	С	71	ASP	2.4
3	D	21	THR	2.4
3	С	9	LEU	2.3
3	С	47	GLU	2.3
3	С	66	ILE	2.3
3	D	60	ARG	2.3
3	С	49	PHE	2.3
3	D	34	SER	2.3
3	С	58	LEU	2.3
1	A	220	ARG	2.2
3	С	33	LEU	2.2
3	D	2	ASN	2.0



Continued from previous page...

Mol	Chain	$\operatorname{Res}$	Type	RSRZ
3	С	70	GLU	2.0

#### 6.2Non-standard residues in protein, DNA, RNA chains (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median,  $95^{th}$  percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	${f B\text{-factors}}({f \AA}^2)$	Q < 0.9
1	CSO	A	279	7/8	0.97	0.12	22,24,39,61	0
1	CSO	A	131	7/8	0.98	0.11	25,26,50,52	0

#### Carbohydrates (i) 6.3

There are no carbohydrates in this entry.

#### 6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median,  $95^{th}$  percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	${f B-factors}({f A}^2)$	Q<0.9
4	G5A	A	401	27/27	0.89	0.19	19,34,49,52	27
7	PNS	D	101	21/22	0.92	0.12	50,59,68,69	0
7	PNS	С	101	21/22	0.92	0.11	48,57,66,73	0
4	G5A	В	402	27/27	0.93	0.14	23,32,42,51	27
5	CL	A	402	1/1	0.97	0.12	37,37,37,37	0
6	ZN	В	401	1/1	0.97	0.08	38,38,38,38	1
5	CL	В	403	1/1	0.98	0.10	45,45,45,45	0


The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




#### Electron density around G5A A 401:

 $2 \mathrm{mF}_o\text{-DF}_c$  (at 0.7 rmsd) in gray  $\mathrm{mF}_o\text{-DF}_c$  (at 3 rmsd) in purple (negative) and green (positive)










#### Electron density around PNS C 101:


 $2 \text{mF}_o\text{-DF}_c$  (at 0.7 rmsd) in gray  $\text{mF}_o\text{-DF}_c$  (at 3 rmsd) in purple (negative) and green (positive)











## 6.5 Other polymers (i)

There are no such residues in this entry.

