



Full wwPDB X-ray Structure Validation Report ⓘ

Dec 4, 2023 – 06:45 pm GMT

PDB ID : 1H3T
Title : Crystal structure of the human igg1 fc-fragment, glycoform (mn2f)2
Authors : Krapp, S.; Mimura, Y.; Jefferis, R.; Huber, R.; Sondermann, P.
Deposited on : 2002-09-19
Resolution : 2.40 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.4, CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3907 (2.40-2.40)
Clashscore	141614	4398 (2.40-2.40)
Ramachandran outliers	138981	4318 (2.40-2.40)
Sidechain outliers	138945	4319 (2.40-2.40)
RSRZ outliers	127900	3811 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	223	 3% 60% 30% 7%
1	B	223	 13% 61% 24% 9%
2	C	4	 75% 25%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	FUL	C	4	-	-	-	X
3	BMA	B	1447	-	-	-	X

2 Entry composition [i](#)


There are 4 unique types of molecules in this entry. The entry contains 3240 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called IG GAMMA-1 CHAIN C REGION.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	208	1597	1016	270	304	7	0	0	1
1	B	202	1490	954	253	276	7	0	0	5

- Molecule 2 is an oligosaccharide called beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[beta-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	C	4	49	28	2	19	0	0	0

- Molecule 3 is beta-D-mannopyranose (three-letter code: BMA) (formula: C₆H₁₂O₆).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	B	1	Total C O 12 6 6	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	58	Total O 58 58	0	0
4	B	34	Total O 34 34	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: IG GAMMA-1 CHAIN C REGION



- Molecule 1: IG GAMMA-1 CHAIN C REGION



- Molecule 2: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[beta-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	49.57Å 80.78Å 139.65Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.00 – 2.40 19.61 – 2.40	Depositor EDS
% Data completeness (in resolution range)	98.1 (50.00-2.40) 94.1 (19.61-2.40)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.29 (at 2.41Å)	Xtrriage
Refinement program	CNS 1.0	Depositor
R, R_{free}	0.280 , 0.309 0.254 , 0.287	Depositor DCC
R_{free} test set	1041 reflections (4.89%)	wwPDB-VP
Wilson B-factor (Å ²)	43.5	Xtrriage
Anisotropy	0.297	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.27 , 41.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.42$, $\langle L^2 \rangle = 0.25$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.90	EDS
Total number of atoms	3240	wwPDB-VP
Average B, all atoms (Å ²)	41.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.89% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NAG, BMA, FUL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.44	0/1641	0.67	0/2232
1	B	0.41	0/1516	0.73	2/2028 (0.1%)
All	All	0.42	0/3157	0.70	2/4260 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	288	LYS	N-CA-C	-8.35	88.46	111.00
1	B	287	ALA	N-CA-C	7.05	130.04	111.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1597	0	1483	61	0
1	B	1490	0	1361	59	0
2	C	49	0	43	6	0
3	B	12	0	12	0	0
4	A	58	0	0	2	0
4	B	34	0	0	1	0
All	All	3240	0	2899	124	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 21.

All (124) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:350:THR:HB	1:B:441:LEU:HG	1.42	1.00
1:B:323:VAL:HG12	1:B:324:SER:H	1.32	0.94
1:B:287:ALA:HB1	1:B:289:THR:H	1.33	0.91
1:B:298:SER:C	1:B:299:THR:CA	2.44	0.85
1:B:311:GLN:NE2	1:B:311:GLN:H	1.74	0.84
1:A:291:PRO:O	1:A:292:ARG:HG3	1.78	0.82
1:B:242:LEU:HD13	1:B:336:ILE:HG22	1.64	0.78
1:B:350:THR:HG23	1:B:439:LYS:HG3	1.65	0.77
1:B:311:GLN:H	1:B:311:GLN:CD	1.91	0.73
1:B:328:LEU:CG	1:B:328:LEU:CA	2.68	0.71
1:B:276:ASN:N	1:B:276:ASN:C	2.44	0.70
1:B:350:THR:CB	1:B:441:LEU:HG	2.19	0.70
1:A:278:TYR:CE2	1:A:284:VAL:HA	2.28	0.68
1:A:282:VAL:O	1:A:283:GLN:HB3	1.93	0.68
1:A:353:PRO:HG2	1:A:358:MET:HE1	1.77	0.67
1:B:429:HIS:CD2	1:B:431:ALA:H	2.14	0.66
1:A:291:PRO:HA	1:A:303:VAL:O	1.94	0.66
1:A:429:HIS:CD2	1:A:431:ALA:H	2.14	0.65
1:B:262:VAL:HG13	1:B:301:ARG:HE	1.61	0.65
1:B:429:HIS:HD2	1:B:431:ALA:H	1.44	0.65
1:A:295:GLN:HB2	1:A:300:TYR:CE1	2.32	0.64
1:B:249:ASP:OD1	1:B:255:ARG:HD3	1.98	0.63
1:A:429:HIS:HD2	1:A:431:ALA:H	1.48	0.61
1:B:263:VAL:O	1:B:301:ARG:HA	2.00	0.61
1:B:311:GLN:NE2	1:B:311:GLN:N	2.47	0.61
1:A:280:ASP:OD2	1:A:317:LYS:HD3	2.00	0.60
1:A:422:VAL:HG22	1:A:442:SER:OG	2.01	0.60
1:B:249:ASP:HA	1:B:255:ARG:CD	2.31	0.59
1:A:414:LYS:O	1:A:418:GLN:HG3	2.02	0.59
1:B:309:LEU:HB3	1:B:311:GLN:NE2	2.17	0.59
1:A:268:HIS:CE1	1:A:295:GLN:HE21	2.20	0.59
1:A:291:PRO:C	1:A:292:ARG:HG3	2.24	0.58
1:B:287:ALA:HB1	1:B:289:THR:N	2.14	0.57
1:A:353:PRO:HG2	1:A:358:MET:CE	2.35	0.56
1:B:249:ASP:O	1:B:257:PRO:HG3	2.05	0.56
1:A:288:LYS:O	1:A:289:THR:O	2.24	0.56
1:B:350:THR:HB	1:B:441:LEU:CG	2.25	0.56

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:365:LEU:HD12	1:B:410:LEU:HD23	1.88	0.56
1:A:274:LYS:HE3	1:A:276:ASN:HD21	1.71	0.56
4:A:2034:HOH:O	1:B:409:LYS:HE2	2.05	0.55
1:B:350:THR:CG2	1:B:439:LYS:HG3	2.36	0.55
1:B:240:VAL:HG21	1:B:323:VAL:HG21	1.89	0.55
1:A:314:LEU:HD22	1:A:430:GLU:HG3	1.89	0.54
2:C:2:NAG:H82	2:C:4:FUL:C1	2.37	0.54
1:A:438:GLN:O	1:A:439:LYS:HD3	2.08	0.53
1:A:272:GLN:O	1:A:325:ASN:ND2	2.39	0.53
1:A:320:LYS:HB2	1:A:335:THR:HG22	1.90	0.53
1:A:274:LYS:HE3	1:A:276:ASN:ND2	2.24	0.53
1:A:268:HIS:HE1	1:A:295:GLN:HG3	1.74	0.52
1:A:346:PRO:HB3	1:A:372:PHE:HB3	1.92	0.52
1:A:415:SER:O	1:A:419:GLN:HG3	2.10	0.52
4:A:2031:HOH:O	1:B:351:LEU:HD22	2.09	0.52
1:B:336:ILE:HG12	1:B:337:SER:N	2.25	0.51
1:B:323:VAL:HG12	1:B:324:SER:N	2.12	0.51
1:A:288:LYS:HE3	1:A:288:LYS:H	1.76	0.51
1:A:350:THR:O	1:A:351:LEU:HD23	2.11	0.51
1:A:249:ASP:OD1	1:A:255:ARG:HD3	2.11	0.50
1:A:436:TYR:CD1	1:A:437:THR:N	2.79	0.50
1:B:250:THR:HG22	1:B:257:PRO:HB3	1.93	0.50
1:A:283:GLN:CD	1:A:287:ALA:HB2	2.32	0.49
1:B:287:ALA:O	1:B:288:LYS:HG2	2.11	0.49
1:A:351:LEU:HB2	1:A:366:THR:HB	1.94	0.49
1:B:238:PRO:HG2	1:B:328:LEU:CD1	2.41	0.49
1:A:282:VAL:O	1:A:283:GLN:CB	2.61	0.49
2:C:1:NAG:O3	2:C:1:NAG:C7	2.61	0.49
1:A:241:PHE:CE2	2:C:3:BMA:H3	2.48	0.49
1:B:260:THR:HG23	1:B:305:VAL:HG22	1.95	0.49
1:A:283:GLN:O	1:A:283:GLN:HG2	2.13	0.48
1:A:288:LYS:N	1:A:288:LYS:HD3	2.27	0.48
1:A:288:LYS:H	1:A:288:LYS:CE	2.26	0.48
1:B:249:ASP:HA	1:B:255:ARG:HD3	1.95	0.48
2:C:1:NAG:H61	2:C:2:NAG:H82	1.95	0.48
1:B:350:THR:HG23	1:B:439:LYS:CG	2.38	0.48
1:B:351:LEU:HB2	1:B:366:THR:HB	1.96	0.48
1:A:353:PRO:HD3	1:A:365:LEU:HD23	1.95	0.48
1:B:325:ASN:ND2	1:B:326:LYS:H	2.12	0.48
1:B:383:SER:HB2	1:B:388:GLU:OE1	2.14	0.47
1:B:422:VAL:HG22	1:B:442:SER:HB3	1.96	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:309:LEU:HB3	1:B:311:GLN:HE22	1.78	0.47
1:B:406:LEU:HD12	1:B:406:LEU:C	2.34	0.47
1:A:353:PRO:HD3	1:A:365:LEU:CD2	2.43	0.47
1:B:316:GLY:HA2	4:B:2006:HOH:O	2.15	0.47
1:A:246:LYS:HB2	1:A:249:ASP:OD2	2.14	0.47
1:B:278:TYR:O	1:B:282:VAL:O	2.33	0.47
1:B:279:VAL:HA	1:B:318:GLU:O	2.15	0.47
1:B:249:ASP:C	1:B:257:PRO:HG3	2.35	0.47
1:A:239:SER:HB2	1:A:264:VAL:CG2	2.45	0.46
1:A:297:ASN:OD1	1:A:298:SER:N	2.39	0.46
1:A:422:VAL:HA	1:A:442:SER:OG	2.15	0.46
1:B:338:LYS:NZ	1:B:338:LYS:HB3	2.30	0.46
1:A:270:ASP:OD2	1:A:270:ASP:N	2.49	0.46
1:B:344:ARG:CD	1:B:344:ARG:NH1	2.80	0.45
1:A:288:LYS:H	1:A:288:LYS:CD	2.30	0.44
1:A:278:TYR:HA	1:A:282:VAL:O	2.17	0.44
1:A:361:ASN:HD22	1:A:361:ASN:HA	1.61	0.44
1:A:421:ASN:N	1:A:421:ASN:HD22	2.16	0.44
1:B:275:PHE:CE1	1:B:302:VAL:O	2.70	0.44
1:B:278:TYR:CD1	1:B:278:TYR:N	2.86	0.43
1:A:343:PRO:HA	1:A:373:TYR:O	2.18	0.43
1:A:386:GLN:CA	1:A:386:GLN:CD	2.87	0.43
1:A:436:TYR:CD1	1:A:436:TYR:C	2.92	0.43
1:A:239:SER:HB2	1:A:264:VAL:HG22	2.01	0.43
1:B:388:GLU:HB3	1:B:410:LEU:HD11	1.99	0.43
1:A:283:GLN:CD	1:A:287:ALA:CB	2.87	0.43
1:A:320:LYS:CB	1:A:335:THR:HG22	2.49	0.43
1:B:358:MET:CE	1:B:417:TRP:HE1	2.31	0.43
1:A:278:TYR:CD1	1:A:278:TYR:N	2.88	0.42
1:A:265:ASP:OD2	2:C:1:NAG:O7	2.37	0.42
1:B:357:GLU:C	1:B:359:THR:H	2.23	0.42
1:A:406:LEU:C	1:A:406:LEU:HD12	2.40	0.42
1:B:306:LEU:HD12	1:B:307:THR:N	2.35	0.42
1:B:326:LYS:HE3	1:B:326:LYS:HB2	1.93	0.41
1:A:239:SER:O	1:A:263:VAL:HA	2.21	0.41
2:C:1:NAG:H61	2:C:2:NAG:C7	2.51	0.41
1:B:358:MET:HE3	1:B:417:TRP:HE1	1.85	0.41
1:A:269:GLU:C	1:A:271:PRO:HD3	2.41	0.41
1:B:240:VAL:O	1:B:334:LYS:HE3	2.21	0.41
1:A:393:THR:HA	1:A:408:SER:HA	2.03	0.40
1:B:386:GLN:HA	1:B:387:PRO:HD3	1.81	0.40

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:279:VAL:O	1:A:280:ASP:HB2	2.21	0.40
1:A:272:GLN:OE1	1:A:272:GLN:HA	2.21	0.40
1:A:336:ILE:HG13	1:A:337:SER:H	1.86	0.40
1:B:242:LEU:HD23	1:B:242:LEU:HA	1.84	0.40
1:B:256:THR:HA	1:B:257:PRO:HD3	1.95	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	206/223 (92%)	186 (90%)	13 (6%)	7 (3%)	3	3
1	B	185/223 (83%)	165 (89%)	12 (6%)	8 (4%)	2	2
All	All	391/446 (88%)	351 (90%)	25 (6%)	15 (4%)	3	2

All (15) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	288	LYS
1	A	289	THR
1	A	433	HIS
1	A	444	SER
1	B	286	ASN
1	B	287	ALA
1	B	288	LYS
1	A	283	GLN
1	B	279	VAL
1	B	301	ARG
1	B	324	SER
1	B	358	MET
1	A	292	ARG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	293	GLU
1	B	323	VAL

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	171/206 (83%)	162 (95%)	9 (5%)	22	37
1	B	153/206 (74%)	146 (95%)	7 (5%)	27	43
All	All	324/412 (79%)	308 (95%)	16 (5%)	25	40

All (16) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	242	LEU
1	A	254	SER
1	A	270	ASP
1	A	286	ASN
1	A	288	LYS
1	A	294	GLN
1	A	358	MET
1	A	361	ASN
1	A	390	ASN
1	B	242	LEU
1	B	260	THR
1	B	280	ASP
1	B	285	HIS
1	B	311	GLN
1	B	326	LYS
1	B	335	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (17) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	276	ASN
1	A	295	GLN
1	A	311	GLN
1	A	347	GLN
1	A	361	ASN
1	A	384	ASN
1	A	390	ASN
1	A	419	GLN
1	A	421	ASN
1	A	429	HIS
1	A	434	ASN
1	B	311	GLN
1	B	325	ASN
1	B	390	ASN
1	B	418	GLN
1	B	429	HIS
1	B	434	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

4 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	NAG	C	1	2,1	14,14,15	0.51	0	17,19,21	0.82	0
2	NAG	C	2	2	14,14,15	0.51	0	17,19,21	0.80	1 (5%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	BMA	C	3	2	11,11,12	0.47	0	15,15,17	0.38	0
2	FUL	C	4	2	10,10,11	0.68	0	14,14,16	0.51	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	C	1	2,1	-	3/6/23/26	0/1/1/1
2	NAG	C	2	2	-	2/6/23/26	0/1/1/1
2	BMA	C	3	2	-	0/2/19/22	0/1/1/1
2	FUL	C	4	2	-	-	0/1/1/1

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	2	NAG	C2-N2-C7	-2.26	119.69	122.90

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	C	1	NAG	C3-C2-N2-C7
2	C	1	NAG	C8-C7-N2-C2
2	C	1	NAG	O7-C7-N2-C2
2	C	2	NAG	O5-C5-C6-O6
2	C	2	NAG	C4-C5-C6-O6


There are no ring outliers.

4 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	C	2	NAG	3	0
2	C	3	BMA	1	0
2	C	4	FUL	1	0
2	C	1	NAG	4	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths,

bond angles, torsion angles, and ring geometry for oligosaccharide.



5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	BMA	B	1447	-	12,12,12	0.42	0	17,17,17	0.64	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '2' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	BMA	B	1447	-	-	0/2/22/22	0/1/1/1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	208/223 (93%)	0.13	7 (3%) 45 44	15, 33, 57, 68	0
1	B	202/223 (90%)	0.77	30 (14%) 2 2	16, 43, 98, 107	0
All	All	410/446 (91%)	0.45	37 (9%) 9 8	15, 37, 92, 107	0

All (37) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	285	HIS	9.7
1	B	283	GLN	7.9
1	B	324	SER	7.2
1	A	285	HIS	6.8
1	B	267	SER	6.3
1	B	298	SER	5.7
1	B	284	VAL	5.3
1	A	445	PRO	5.1
1	A	444	SER	4.8
1	B	286	ASN	4.8
1	B	297	ASN	4.6
1	B	299	THR	4.5
1	A	420	GLY	4.2
1	B	326	LYS	4.2
1	B	323	VAL	4.1
1	B	263	VAL	3.9
1	B	295	GLN	3.7
1	B	292	ARG	3.6
1	B	277	TRP	3.2
1	B	327	ALA	3.1
1	B	238	PRO	3.0
1	A	433	HIS	3.0
1	B	328	LEU	3.0
1	B	329	PRO	3.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	325	ASN	3.0
1	B	278	TYR	2.8
1	B	300	TYR	2.6
1	B	264	VAL	2.6
1	B	287	ALA	2.5
1	A	284	VAL	2.5
1	B	281	GLY	2.4
1	B	239	SER	2.4
1	B	256	THR	2.3
1	A	385	GLY	2.1
1	B	274	LYS	2.1
1	B	282	VAL	2.0
1	B	276	ASN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)


There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	FUL	C	4	10/11	0.67	0.47	68,70,71,72	0
2	BMA	C	3	11/12	0.76	0.30	59,62,63,64	0
2	NAG	C	2	14/15	0.82	0.27	56,61,62,63	0
2	NAG	C	1	14/15	0.83	0.20	58,60,64,66	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.



6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	BMA	B	1447	12/12	0.72	0.47	95,95,96,96	0

6.5 Other polymers [i](#)

There are no such residues in this entry.