

Full wwPDB X-ray Structure Validation Report (i)

Aug 6, 2020 – 11:32 AM BST

PDB ID : 5H49

Title : Crystal structure of Cbln1

Authors: Zhong, C.; Shen, J.; Zhang, H.; Ding, J.

Deposited on : 2016-10-31

Resolution : 2.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.orgA user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp

with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity : 4.02b-467

Mogul : 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : 1.13 EDS : 2.13.1

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

Refmac: 5.8.0158

CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)

Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.13.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X- $RAY\ DIFFRACTION$

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$\begin{array}{c} \textbf{Whole archive} \\ (\#\text{Entries}) \end{array}$	$\begin{array}{c} {\rm Similar \; resolution} \\ (\#{\rm Entries, \; resolution \; range(\AA)}) \end{array}$
R_{free}	130704	3140 (2.80-2.80)
Clashscore	141614	3569 (2.80-2.80)
Ramachandran outliers	138981	3498 (2.80-2.80)
Sidechain outliers	138945	3500 (2.80-2.80)
RSRZ outliers	127900	3078 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain				
1	A	184	59%	15%	27%		
1	В	184	5%	12% •	30%		
1	С	184	48%	19% •	31%		



2 Entry composition (i)

There are 3 unique types of molecules in this entry. The entry contains 3124 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Cerebellin-1.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace		
1	Λ	135	Total	С	N	О	S	0	0	0
1	Α	133	1073	686	186	196	5	0	U	U
1	D	128	Total	С	N	О	S	0	0	0
1	Б	120	1009	649	171	185	4	0	U	
1	С	127	Total	С	N	О	S	0	0	0
1		121	1006	648	174	180	4	U	U	U

There are 36 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	10	GLY	-	expression tag	UNP P63182
A	11	THR	-	expression tag	UNP P63182
A	12	HIS	-	expression tag	UNP P63182
A	13	HIS	-	expression tag	UNP P63182
A	14	HIS	-	expression tag	UNP P63182
A	15	HIS	_	expression tag	UNP P63182
A	16	HIS	_	expression tag	UNP P63182
A	17	HIS	-	expression tag	UNP P63182
A	18	HIS	-	expression tag	UNP P63182
A	19	HIS	-	expression tag	UNP P63182
A	20	GLY	-	expression tag	UNP P63182
A	21	SER	-	expression tag	UNP P63182
В	10	GLY	-	expression tag	UNP P63182
В	11	THR	-	expression tag	UNP P63182
В	12	HIS	-	expression tag	UNP P63182
В	13	HIS	-	expression tag	UNP P63182
В	14	HIS	-	expression tag	UNP P63182
В	15	HIS	-	expression tag	UNP P63182
В	16	HIS	-	expression tag	UNP P63182
В	17	HIS	-	expression tag	UNP P63182
В	18	HIS	-	expression tag	UNP P63182
В	19	HIS	-	expression tag	UNP P63182
В	20	GLY	-	expression tag	UNP P63182



Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
В	21	SER	=	expression tag	UNP P63182
С	10	GLY	-	expression tag	UNP P63182
С	11	THR	_	expression tag	UNP P63182
С	12	HIS	-	expression tag	UNP P63182
С	13	HIS	-	expression tag	UNP P63182
С	14	HIS	_	expression tag	UNP P63182
С	15	HIS	-	expression tag	UNP P63182
С	16	HIS	_	expression tag	UNP P63182
С	17	HIS	-	expression tag	UNP P63182
С	18	HIS	-	expression tag	UNP P63182
С	19	HIS	_	expression tag	UNP P63182
С	20	GLY	_	expression tag	UNP P63182
С	21	SER	-	expression tag	UNP P63182

• Molecule 2 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: $C_8H_{15}NO_6$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total 14	C 8	N 1	O 5	0	0

• Molecule 3 is water.

Mol	Chain	Residues	${f Atoms}$	ZeroOcc	AltConf
3	A	14	Total O 14 14	0	0




 $Continued\ from\ previous\ page...$

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	В	4	Total O 4 4	0	0
3	С	4	Total O 4 4	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.





4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants	91.64Å 91.64Å 151.05Å	Depositor
a, b, c, α , β , γ	90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.00 - 2.80	Depositor
resolution (A)	44.13 - 2.80	EDS
% Data completeness	99.7 (50.00-2.80)	Depositor
(in resolution range)	99.7 (44.13-2.80)	EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	4.45 (at 2.81Å)	Xtriage
Refinement program	REFMAC 5.8.0155	Depositor
D D.	0.271 , 0.307	Depositor
R, R_{free}	0.271 , 0.310	DCC
R_{free} test set	829 reflections (5.07%)	wwPDB-VP
Wilson B-factor (Å ²)	31.6	Xtriage
Anisotropy	0.077	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.33 , 37.1	EDS
L-test for twinning ²	$ < L >=0.50, < L^2>=0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.85	EDS
Total number of atoms	3124	wwPDB-VP
Average B, all atoms (Å ²)	32.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 3.41% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of <|L|>, $< L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond angles	
IVIOI	Chain	RMSZ	# Z >5	RMSZ	# Z > 5
1	A	0.28	0/1097	0.47	0/1481
1	В	0.27	0/1030	0.49	0/1389
1	С	0.27	0/1027	0.47	0/1384
All	All	0.27	0/3154	0.48	0/4254

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1073	0	1057	19	0
1	В	1009	0	995	16	0
1	С	1006	0	1003	51	0
2	A	14	0	13	1	0
3	A	14	0	0	0	0
3	В	4	0	0	1	0
3	С	4	0	0	0	0
All	All	3124	0	3068	82	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 13.



All (82) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

		Interatomic	Clash
Atom-1	Atom-2	${f distance} \; ({f \mathring{A}})$	overlap (Å)
1:C:102:SER:HB2	1:C:185:PHE:CD2	1.35	1.62
1:C:102:SER:HB2	1:C:185:PHE:CE2	1.40	1.54
1:C:102:SER:CB	1:C:185:PHE:CD2	2.19	1.26
1:C:97:PHE:CZ	1:C:101:ARG:O	1.96	1.18
1:C:102:SER:CB	1:C:185:PHE:CE2	2.33	1.10
1:C:97:PHE:CZ	1:C:102:SER:HA	1.89	1.08
1:C:97:PHE:CD1	1:C:103:THR:O	2.05	1.08
1:C:102:SER:CB	1:C:185:PHE:HD2	1.71	0.94
1:C:97:PHE:HE1	1:C:102:SER:C	1.75	0.88
1:C:102:SER:HB2	1:C:185:PHE:HE2	1.34	0.86
1:C:97:PHE:HZ	1:C:102:SER:HA	1.34	0.86
1:C:97:PHE:CE1	1:C:102:SER:HA	2.12	0.84
1:C:97:PHE:CE1	1:C:102:SER:C	2.53	0.82
1:C:97:PHE:HZ	1:C:101:ARG:O	1.62	0.80
1:C:102:SER:O	1:C:185:PHE:HE2	1.71	0.74
1:C:97:PHE:HD1	1:C:103:THR:O	1.69	0.73
1:C:85:TYR:HB2	1:C:100:GLU:O	1.90	0.71
1:C:97:PHE:CE2	1:C:101:ARG:O	2.42	0.71
1:C:97:PHE:CE1	1:C:102:SER:CA	2.73	0.71
1:B:128:GLN:OE1	1:B:173:ARG:NH1	2.24	0.70
1:A:146:GLN:HE22	1:B:146:GLN:HE21	1.39	0.69
1:C:97:PHE:CZ	1:C:102:SER:CA	2.73	0.69
1:B:126:THR:HG22	1:B:144:GLY:HA2	1.75	0.69
1:C:67:ILE:HD11	1:C:90:LEU:HD11	1.78	0.65
1:A:70:THR:HG22	1:A:72:HIS:H	1.61	0.65
1:B:98:ASP:HB2	1:B:101:ARG:HD3	1.80	0.63
1:C:115:ASN:ND2	1:C:186:SER:OG	2.33	0.61
1:C:149:THR:HG22	1:C:150:ARG:HG3	1.83	0.61
1:C:85:TYR:HA	1:C:101:ARG:HA	1.84	0.60
1:C:102:SER:CB	1:C:185:PHE:HE2	2.00	0.60
1:C:102:SER:HB3	1:C:185:PHE:CD2	2.33	0.60
1:B:71:ASN:HB3	1:B:181:LYS:HE3	1.83	0.60
1:A:126:THR:HA	1:A:144:GLY:HA3	1.84	0.59
1:A:127:ILE:HG22	1:A:174:GLY:HA3	1.85	0.58
1:B:120:LYS:HD3	1:B:127:ILE:HG12	1.85	0.58
1:C:102:SER:HB3	1:C:185:PHE:HD2	1.66	0.58
1:C:126:THR:HG22	1:C:145:ASP:H	1.68	0.57
1:C:102:SER:O	1:C:185:PHE:CE2	2.55	0.57
1:C:99:SER:HB3	1:C:101:ARG:HE	1.70	0.56
1:A:150:ARG:HH21	1:A:181:LYS:HZ3	1.54	0.56



Continued from previous page...

	ious page	Interatomic	Clash
Atom-1	Atom-2	${\rm distance}\;({\rm \AA})$	overlap(Å)
1:C:97:PHE:CE1	1:C:103:THR:O	2.58	0.55
1:C:70:THR:HG23	1:C:181:LYS:HD3	1.88	0.54
1:B:127:ILE:HG22	1:B:174:GLY:HA3	1.90	0.53
1:C:136:TRP:HE3	1:C:137:PRO:HD2	1.74	0.53
1:C:97:PHE:HE1	1:C:102:SER:CA	2.19	0.53
1:C:86:PHE:CE1	1:C:102:SER:HB3	2.44	0.53
1:B:143:ALA:HB1	1:B:151:GLU:HG2	1.92	0.51
1:A:139:ILE:HD13	1:A:157:VAL:HB	1.91	0.51
1:A:65:SER:HB2	1:C:158:LEU:HB2	1.93	0.51
1:A:161:MET:HG3	1:A:165:ASP:HB2	1.92	0.50
1:A:68:ARG:HG2	1:A:180:TRP:O	2.14	0.48
1:C:126:THR:HA	1:C:144:GLY:HA3	1.95	0.48
1:B:98:ASP:CB	1:B:101:ARG:HD3	2.44	0.47
1:A:76:GLU:N	2:A:201:NAG:H82	2.30	0.47
1:A:171:LEU:HD22	1:A:176:LEU:HG	1.96	0.46
1:B:76:GLU:HA	3:B:202:HOH:O	2.16	0.45
1:B:67:ILE:HG13	1:B:184:THR:HG22	1.99	0.45
1:A:158:LEU:HB2	1:B:65:SER:HB2	1.99	0.44
1:C:97:PHE:HE1	1:C:102:SER:O	1.99	0.44
1:C:120:LYS:HE2	1:C:151:GLU:HB3	1.98	0.44
1:C:68:ARG:HD2	1:C:87:ASP:OD1	2.17	0.44
1:A:180:TRP:HE3	1:A:180:TRP:H	1.62	0.44
1:B:176:LEU:HD13	1:B:180:TRP:HB3	1.99	0.44
1:A:97:PHE:CZ	1:A:102:SER:HA	2.53	0.43
1:C:139:ILE:HD13	1:C:157:VAL:HB	1.99	0.43
1:B:84:ILE:O	1:B:102:SER:HB2	2.18	0.43
1:C:176:LEU:HD13	1:C:180:TRP:HB3	2.00	0.43
1:C:85:TYR:CA	1:C:101:ARG:HA	2.49	0.42
1:C:101:ARG:HG3	1:C:103:THR:H	1.84	0.42
1:B:125:GLN:HE21	1:B:174:GLY:HA2	1.85	0.42
1:C:138:VAL:HG11	1:C:159:ILE:HD11	2.02	0.42
1:C:86:PHE:HE1	1:C:102:SER:HB3	1.83	0.42
1:A:68:ARG:HD2	1:A:87:ASP:OD1	2.20	0.42
1:A:74:PRO:HG3	1:A:180:TRP:CZ3	2.54	0.42
1:C:68:ARG:HG2	1:C:180:TRP:CD1	2.55	0.41
1:C:112:TYR:CZ	1:C:189:LEU:HD13	2.55	0.41
1:C:148:VAL:HG23	1:C:151:GLU:HB2	2.02	0.41
1:A:114:PHE:HE2	1:A:161:MET:HE1	1.86	0.41
1:A:144:GLY:O	1:A:146:GLN:HG2	2.20	0.41
1:C:146:GLN:HG2	1:C:147:ASP:H	1.86	0.41
1:B:101:ARG:H	1:B:101:ARG:HD2	1.86	0.41



Continued from previous page...

Atom-1	Atom-2	$egin{array}{c} ext{Interatomic} \ ext{distance} \ (ext{Å}) \end{array}$	$egin{aligned} ext{Clash} \ ext{overlap } (ext{Å}) \end{aligned}$
1:A:149:THR:HG23	1:C:145:ASP:HB2	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	\mathbf{ntiles}
1	A	133/184 (72%)	127 (96%)	6 (4%)	0	100	100
1	В	122/184 (66%)	116 (95%)	6 (5%)	0	100	100
1	С	123/184 (67%)	118 (96%)	5 (4%)	0	100	100
All	All	378/552 (68%)	361 (96%)	17 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	116/158 (73%)	115 (99%)	1 (1%)	78 94
1	В	109/158 (69%)	108 (99%)	1 (1%)	78 94
1	С	107/158 (68%)	104 (97%)	3 (3%)	43 77
All	All	332/474 (70%)	327 (98%)	5 (2%)	65 89



All	(5)	residues	with	a	non-rotame	ric	sidechain	are	listed	below:
	\ \sigma_{/}	LODIGICO	***	~	IIOII IOGGIIIO		DIGCOLLAND	COL C	110004	~ ~ ~ ~ ~ ~ ~ .

Mol	Chain	Res	Type
1	A	98	ASP
1	В	101	ARG
1	С	99	SER
1	С	101	ARG
1	С	145	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	146	GLN
1	С	146	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bo	Bond lengths				les
IVIOI	туре	Chain	nes	LIIIK	Counts	Counts RMSZ #		Counts	RMSZ	$\mid \# Z > 2 \mid$
2	NAG	A	201	1	14,14,15	0.22	0	17,19,21	0.44	0



In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	A	201	1	-	0/6/23/26	0/1/1/1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	201	NAG	1	0

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ $>$	$\#\mathrm{RSRZ}{>}2$	$OWAB(A^2)$	Q < 0.9
1	A	135/184~(73%)	-0.12	5 (3%) 41 31	14, 22, 60, 100	0
1	В	128/184 (69%)	0.29	10 (7%) 13 7	16, 29, 81, 119	0
1	С	127/184 (69%)	0.24	10 (7%) 12 7	16, 25, 63, 115	0
All	All	390/552 (70%)	0.13	25 (6%) 19 12	14, 26, 71, 119	0

All (25) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	С	102	SER	13.3
1	В	99	SER	11.9
1	С	145	ASP	8.9
1	A	100	GLU	5.5
1	С	100	GLU	5.1
1	С	101	ARG	4.7
1	В	147	ASP	4.1
1	A	99	SER	3.8
1	A	76	GLU	3.3
1	В	149	THR	3.2
1	В	81	THR	3.2
1	С	136	TRP	3.1
1	В	98	ASP	2.8
1	A	79	ASN	2.7
1	В	100	GLU	2.7
1	В	75	SER	2.6
1	В	76	GLU	2.6
1	A	136	TRP	2.6
1	С	146	GLN	2.5
1	В	177	MET	2.5
1	С	103	THR	2.4
1	В	101	ARG	2.4
1	С	58	GLY	2.3



Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	С	99	SER	2.1
1	С	80	ARG	2.1

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-}\mathbf{factors}(\mathbf{\mathring{A}}^2)$	Q < 0.9
2	NAG	A	201	14/15	0.86	0.33	48,54,59,60	0

6.5 Other polymers (i)

There are no such residues in this entry.

