



# Full wwPDB X-ray Structure Validation Report ⓘ

Dec 10, 2023 – 12:19 pm GMT

PDB ID : 1H7K  
Title : Formation of a tyrosyl radical intermediate in *Proteus mirabilis* catalase by directed mutagenesis and consequences for nucleotide reactivity  
Authors : Andreoletti, P.; Gambarelli, S.; Gaillard, J.; Sainz, G.; Stojanoff, V.; Jouve, H.M.  
Deposited on : 2001-07-08  
Resolution : 2.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.4, CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.36  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*


The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	3907 (2.40-2.40)
Clashscore	141614	4398 (2.40-2.40)
Ramachandran outliers	138981	4318 (2.40-2.40)
Sidechain outliers	138945	4319 (2.40-2.40)
RSRZ outliers	127900	3811 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	483	

## 2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 4083 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called CATALASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	476	3863	2447	685	716	15	0	0	1

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	215	TYR	PHE	conflict	UNP P42321
A	53	OMT	MET	modified residue	UNP P42321

- Molecule 2 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula:  $C_{34}H_{32}FeN_4O_4$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	Fe	N	O		
2	A	1	43	34	1	4	4	0	0

- Molecule 3 is ACETATE ION (three-letter code: ACT) (formula:  $C_2H_3O_2$ ).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O 4 2 2	0	0

- Molecule 4 is SULFATE ION (three-letter code: SO<sub>4</sub>) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total O S 5 4 1	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	168	Total 168	O 168	0	0

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: CATALASE



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 62 2 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	110.00Å 110.00Å 251.20Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	29.26 – 2.40 29.82 – 2.40	Depositor EDS
% Data completeness (in resolution range)	99.7 (29.26-2.40) 99.6 (29.82-2.40)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.47 (at 2.39Å)	Xtrriage
Refinement program	CNS 1.0	Depositor
R, $R_{free}$	0.237 , 0.240 0.231 , 0.224	Depositor DCC
$R_{free}$ test set	1411 reflections (3.93%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	44.5	Xtrriage
Anisotropy	0.456	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 42.0	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.46$ , $\langle L^2 \rangle = 0.29$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	4083	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	40.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.71% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, HEM, ACT, OMT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.76	14/3963 (0.4%)	0.90	18/5363 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	3

All (14) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	99	ARG	NE-CZ	17.48	1.55	1.33
1	A	99	ARG	CZ-NH1	-14.31	1.14	1.33
1	A	472	GLU	CD-OE2	13.10	1.40	1.25
1	A	357	PRO	N-CA	11.24	1.66	1.47
1	A	364	TYR	CB-CG	-9.98	1.36	1.51
1	A	349	HIS	CG-CD2	-9.31	1.20	1.35
1	A	18	ASN	C-N	-9.06	1.13	1.34
1	A	364	TYR	C-N	8.36	1.53	1.34
1	A	18	ASN	C-O	8.09	1.38	1.23
1	A	356	ALA	C-N	-7.63	1.19	1.34
1	A	153	THR	CA-CB	-7.25	1.34	1.53
1	A	349	HIS	CG-ND1	5.83	1.51	1.38
1	A	103	ASP	CB-CG	5.74	1.63	1.51
1	A	103	ASP	CG-OD2	5.23	1.37	1.25

All (18) bond angle outliers are listed below:



Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	99	ARG	NE-CZ-NH2	-28.45	106.07	120.30
1	A	99	ARG	NH1-CZ-NH2	15.83	136.82	119.40
1	A	364	TYR	CB-CG-CD2	-13.06	113.16	121.00
1	A	364	TYR	CB-CG-CD1	12.58	128.55	121.00
1	A	103	ASP	CB-CG-OD2	-11.85	107.64	118.30
1	A	357	PRO	CA-N-CD	-8.08	100.18	111.50
1	A	413	ARG	NE-CZ-NH2	7.38	123.99	120.30
1	A	18	ASN	CA-C-O	-6.88	105.65	120.10
1	A	99	ARG	NE-CZ-NH1	-6.41	117.09	120.30
1	A	54	HIS	O-C-N	-6.40	112.47	122.70
1	A	317	ASN	O-C-N	6.14	132.53	122.70
1	A	6	LEU	CA-CB-CG	6.13	129.41	115.30
1	A	365	HIS	N-CA-CB	-6.02	99.76	110.60
1	A	153	THR	CA-CB-CG2	5.96	120.75	112.40
1	A	6	LEU	CB-CG-CD1	-5.61	101.46	111.00
1	A	349	HIS	CB-CG-ND1	-5.44	109.60	123.20
1	A	153	THR	CA-CB-OG1	5.37	120.29	109.00
1	A	364	TYR	C-N-CA	-5.25	108.56	121.70

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	103	ASP	Sidechain
1	A	18	ASN	Mainchain
1	A	349	HIS	Sidechain

## 5.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3863	0	3677	25	0
2	A	43	0	30	3	0
3	A	4	0	3	0	0
4	A	5	0	0	0	0
5	A	168	0	0	0	0
All	All	4083	0	3710	28	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (28) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:600:HEM:HBB2	2:A:600:HEM:HMB2	1.59	0.84
1:A:464:HIS:HD2	1:A:466:GLU:H	1.41	0.68
1:A:423:ARG:HD3	1:A:463:VAL:O	1.95	0.66
2:A:600:HEM:HBB2	2:A:600:HEM:CMB	2.27	0.64
1:A:56:LYS:HE3	1:A:101:ALA:O	1.99	0.63
1:A:189:ARG:HD3	1:A:243:ASP:OD1	2.01	0.61
1:A:54:HIS:CE1	1:A:95:VAL:HG22	2.41	0.56
1:A:132:PHE:HB3	1:A:278:LEU:HD13	1.88	0.56
1:A:278:LEU:HD12	1:A:329:MET:HG3	1.88	0.55
1:A:359:CYS:HB2	1:A:360:PRO:CD	2.37	0.55
1:A:310:GLN:HB3	1:A:352:ILE:HD13	1.89	0.54
1:A:54:HIS:HA	1:A:94:THR:O	2.08	0.54
1:A:179:MET:HG3	1:A:441:ILE:HG12	1.91	0.52
1:A:436:ARG:O	1:A:440:ARG:HG3	2.10	0.52
1:A:433:GLU:O	1:A:437:MET:HG2	2.12	0.49
1:A:131:VAL:HG23	1:A:174:GLN:HE21	1.77	0.49
1:A:73:TYR:HE2	1:A:290:MET:HE1	1.79	0.46
1:A:359:CYS:HB2	1:A:360:PRO:HD2	1.98	0.45
2:A:600:HEM:CMC	2:A:600:HEM:HBC2	2.46	0.44
1:A:254:PRO:HG2	1:A:297:LEU:HB2	2.00	0.43
1:A:398:GLU:HB2	1:A:399:PRO:HD2	2.00	0.43
1:A:5:LYS:HE3	1:A:5:LYS:HB2	1.90	0.42
1:A:173:HIS:HA	1:A:422:PRO:HB3	2.00	0.42
1:A:200:SER:OG	1:A:323:SER:HB3	2.21	0.41
1:A:371:ARG:HD2	1:A:371:ARG:HA	1.83	0.41
1:A:437:MET:O	1:A:441:ILE:HG13	2.21	0.40
1:A:253:TYR:HA	1:A:254:PRO:HD2	1.96	0.40
1:A:128:ASN:HA	1:A:192:HIS:O	2.22	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries

of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	473/483 (98%)	454 (96%)	19 (4%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	407/413 (98%)	379 (93%)	28 (7%)	15	25

All (28) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	15	VAL
1	A	30	LEU
1	A	37	LEU
1	A	47	VAL
1	A	52	ARG
1	A	113	LEU
1	A	117	THR
1	A	129	THR
1	A	147	VAL
1	A	172	LEU
1	A	191	VAL
1	A	201	PHE
1	A	237	ARG
1	A	242	ARG
1	A	244	LEU
1	A	270	THR
1	A	271	VAL
1	A	278	LEU
1	A	284	HIS

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	297	LEU
1	A	330	LEU
1	A	334	LEU
1	A	363	ASN
1	A	366	ARG
1	A	414	GLU
1	A	429	LEU
1	A	459	LEU
1	A	467	TYR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	42	HIS
1	A	121	ASN
1	A	127	ASN
1	A	174	GLN
1	A	259	GLN
1	A	300	ASN
1	A	341	HIS
1	A	363	ASN
1	A	464	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
1	OMT	A	53	1	8,9,10	2.09	2 (25%)	6,12,14	2.83	3 (50%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	OMT	A	53	1	-	1/7/8/10	-

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	53	OMT	CG-SD	-4.26	1.72	1.78
1	A	53	OMT	CE-SD	-3.37	1.62	1.75

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	53	OMT	OD2-SD-OD1	-4.68	107.08	117.09
1	A	53	OMT	OD1-SD-CG	-3.10	106.17	108.34
1	A	53	OMT	OD1-SD-CE	3.02	111.94	108.91

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	53	OMT	N-CA-CB-CG

There are no ring outliers.

No monomer is involved in short contacts.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	ACT	A	700	-	3,3,3	1.61	1 (33%)	3,3,3	1.95	1 (33%)
2	HEM	A	600	1	41,50,50	1.67	7 (17%)	45,82,82	1.23	6 (13%)
4	SO4	A	800	-	4,4,4	0.81	0	6,6,6	0.57	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	HEM	A	600	1	-	4/12/54/54	-

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	600	HEM	CBB-CAB	4.78	1.54	1.30
2	A	600	HEM	CBC-CAC	3.88	1.55	1.29
2	A	600	HEM	C3C-C2C	-3.86	1.35	1.40
2	A	600	HEM	C3C-CAC	3.01	1.54	1.47
3	A	700	ACT	O-C	2.36	1.33	1.22
2	A	600	HEM	O2A-CGA	-2.27	1.23	1.30
2	A	600	HEM	CAB-C3B	2.24	1.53	1.47
2	A	600	HEM	C2C-C1C	2.22	1.47	1.42

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	600	HEM	C4C-CHD-C1D	3.44	127.10	122.56
3	A	700	ACT	O-C-CH3	-2.71	111.77	122.33
2	A	600	HEM	CMC-C2C-C3C	2.67	129.68	124.68
2	A	600	HEM	CMA-C3A-C4A	-2.67	124.36	128.46
2	A	600	HEM	C4B-CHC-C1C	2.63	126.03	122.56
2	A	600	HEM	CBB-CAB-C3B	-2.62	114.61	127.62
2	A	600	HEM	O1A-CGA-CBA	-2.54	114.91	123.08

There are no chirality outliers.

All (4) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	600	HEM	CAA-CBA-CGA-O1A
2	A	600	HEM	CAD-CBD-CGD-O2D
2	A	600	HEM	CAA-CBA-CGA-O2A
2	A	600	HEM	CAD-CBD-CGD-O1D

There are no ring outliers.

1 monomer is involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	600	HEM	3	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
1	A	2

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	A	356:ALA	C	357:PRO	N	1.19

*Continued on next page...*



*Continued from previous page...*

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	A	18:ASN	C	19:ASN	N	1.13

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	475/483 (98%)	0.04	14 (2%) 51 50	19, 40, 54, 59	0

All (14) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	360	PRO	4.0
1	A	431	ASP	3.4
1	A	284	HIS	3.2
1	A	81	GLU	2.9
1	A	227	ASP	2.8
1	A	450	LYS	2.7
1	A	362	HIS	2.7
1	A	395	ASP	2.6
1	A	451	GLU	2.6
1	A	414	GLU	2.4
1	A	359	CYS	2.4
1	A	361	PHE	2.3
1	A	479	GLU	2.1
1	A	252	ASP	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
1	OMT	A	53	10/11	0.93	0.14	26,31,37,38	0

### 6.3 Carbohydrates [i](#)


There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	ACT	A	700	4/4	0.83	0.37	74,74,74,74	0
4	SO4	A	800	5/5	0.94	0.12	90,90,90,90	0
2	HEM	A	600	43/43	0.97	0.12	18,23,24,27	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



## 6.5 Other polymers [i](#)

There are no such residues in this entry.