



# Full wwPDB X-ray Structure Validation Report ⓘ

Jan 7, 2024 – 07:14 am GMT

PDB ID : 6H8H  
Title : Molybdenum storage protein - in a recombinant and in vivo-like form  
Authors : Ermler, U.; Bruenle, S.  
Deposited on : 2018-08-02  
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.4, CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.36  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*



The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	6207 (1.90-1.90)
Clashscore	141614	6847 (1.90-1.90)
Ramachandran outliers	138981	6760 (1.90-1.90)
Sidechain outliers	138945	6760 (1.90-1.90)
RSRZ outliers	127900	6082 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	B	269	
2	A	275	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
6	PO4	B	304	-	-	-	X
6	PO4	B	305	-	-	-	X
9	MOO	A	304	-	-	X	-

## 2 Entry composition [i](#)

There are 11 unique types of molecules in this entry. The entry contains 4262 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Molybdenum storage protein subunit beta.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	B	268	Total 2010	C 1277	N 353	O 372	S 8	0	7	0

- Molecule 2 is a protein called Molybdenum storage protein subunit alpha.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	A	244	Total 1824	C 1154	N 339	O 328	S 3	0	1	0

- Molecule 3 is bis(mu4-oxo)-tetrakis(mu3-oxo)-hexakis(mu2-oxo)-hexadeca-oxo-octamolybdenum (VI) (three-letter code: 8M0) (formula: Mo<sub>8</sub>O<sub>28</sub>).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	Mo	O		
3	B	1	Total 36	Mo 8	O 28	28	0
3	A	1	Total 34	Mo 8	O 26	0	0

- Molecule 4 is Mo5 Cluster (three-letter code: FUQ) (formula:  $\text{H}_{20}\text{Mo}_5\text{O}_{25}$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	B	1	Total	Mo	O	24	0
			29	5	24		

- Molecule 5 is Mo14O47 cluster (three-letter code: GX2) (formula:  $\text{H}_{17}\text{Mo}_{14}\text{O}_{47}$ ).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	B	1	Total	Mo	O	18	0
			24	6	18		

- Molecule 6 is PHOSPHATE ION (three-letter code: PO4) (formula:  $\text{O}_4\text{P}$ ).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	B	1	Total O P 5 4 1	0	0
6	B	1	Total O P 5 4 1	0	0

- Molecule 7 is ADENOSINE-5'-TRIPHOSPHATE (three-letter code: ATP) (formula:  $C_{10}H_{16}N_5O_{13}P_3$ ).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	1	Total C N O P 31 10 5 13 3	0	0

- Molecule 8 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	1	Total	Mg	0	0
			1	1		

- Molecule 9 is MOLYBDATE ION (three-letter code: MOO) (formula: MoO<sub>4</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
9	A	1	Total	Mo	O	0	0
			5	1	4		

- Molecule 10 is bis(mu2-oxo)-octa-oxo-dimolybdenum (VI) (three-letter code: M27) (formula: Mo<sub>2</sub>O<sub>10</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
10	A	1	Total	Mo	O	9	0
			11	2	9		

- Molecule 11 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
11	B	111	Total	O	0	0
			111	111		
11	A	136	Total	O	0	0
			136	136		




### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Molybdenum storage protein subunit beta



- Molecule 2: Molybdenum storage protein subunit alpha



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 63 2 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	117.10Å 117.10Å 235.62Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	46.94 – 1.90 46.94 – 1.90	Depositor EDS
% Data completeness (in resolution range)	99.5 (46.94-1.90) 99.3 (46.94-1.90)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.10	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.19 (at 1.90Å)	Xtrriage
Refinement program	PHENIX (1.13_2998: ???)	Depositor
R, $R_{free}$	0.206 , 0.221 0.206 , 0.220	Depositor DCC
$R_{free}$ test set	3786 reflections (5.04%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	32.7	Xtrriage
Anisotropy	0.300	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.42 , 53.4	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	4262	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	37.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.52% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: FUQ, MOO, M27, PO4, 8M0, ATP, MG, GX2

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	B	0.24	0/2069	0.42	0/2809
2	A	0.24	0/1867	0.42	0/2546
All	All	0.24	0/3936	0.42	0/5355

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	2010	0	2083	17	0
2	A	1824	0	1864	9	0
3	A	34	0	0	3	0
3	B	36	0	0	0	0
4	B	29	0	0	0	0
5	B	24	0	0	0	0
6	B	10	0	0	0	0
7	A	31	0	12	1	0
8	A	1	0	0	0	0
9	A	5	0	0	1	1
10	A	11	0	0	0	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
11	A	136	0	0	1	0
11	B	111	0	0	0	0
All	All	4262	0	3959	25	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (25) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
9:A:304:MOO:MO	9:A:304:MOO:O1	1.72	0.60
3:A:303:8M0:MO2	3:A:303:8M0:O6	1.73	0.59
2:A:130:HIS:HB2	3:A:303:8M0:O21	2.04	0.58
1:B:104:SER:HA	1:B:149:MET:HG3	1.85	0.57
1:B:179:GLN:HG2	2:A:157:HIS:HB3	1.87	0.55
1:B:195:TYR:HB3	1:B:205:ALA:HB1	1.89	0.54
1:B:228:PHE:CD2	1:B:229:PRO:HD3	2.44	0.53
1:B:227:GLU:O	1:B:230:VAL:HG12	2.10	0.52
1:B:65:ALA:HA	1:B:68:LYS:HE3	1.92	0.51
2:A:227:PRO:HD3	7:A:301:ATP:C6	2.50	0.47
2:A:190:GLU:OE2	11:A:401:HOH:O	2.20	0.47
2:A:156:HIS:HE1	3:A:303:8M0:O23	1.97	0.47
1:B:130:GLY:HA3	1:B:180:PHE:CZ	2.52	0.45
2:A:193:ASP:HA	2:A:249:ASN:HB2	1.99	0.44
1:B:73:LEU:HD21	1:B:136:LEU:HD11	2.00	0.43
2:A:205:ARG:H	2:A:205:ARG:HG3	1.64	0.43
1:B:222:HIS:CD2	1:B:228:PHE:HB2	2.53	0.43
1:B:149:MET:HE3	1:B:150:PRO:HD2	2.01	0.42
1:B:155:TRP:NE1	2:A:177:LEU:HG	2.34	0.42
1:B:170:ASP:HB2	1:B:227:GLU:CD	2.40	0.42
1:B:151:PRO:O	1:B:153:LYS:HG2	2.20	0.41
1:B:203:LYS:HE3	1:B:204:ASP:OD2	2.20	0.41
1:B:157:ARG:HD2	2:A:235:VAL:HG12	2.03	0.41
1:B:192:ASP:HA	1:B:247:ASN:HB2	2.03	0.41
1:B:149:MET:HA	1:B:150:PRO:HD3	1.98	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
9:A:304:MOO:MO	9:A:304:MOO:O2[3_665]	2.06	0.14

### 5.3 Torsion angles [i](#)

#### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	273/269 (102%)	263 (96%)	9 (3%)	1 (0%)	34	24
2	A	243/275 (88%)	236 (97%)	7 (3%)	0	100	100
All	All	516/544 (95%)	499 (97%)	16 (3%)	1 (0%)	47	38

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	128	GLY

#### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	212/205 (103%)	208 (98%)	4 (2%)	57	53
2	A	188/215 (87%)	188 (100%)	0	100	100
All	All	400/420 (95%)	396 (99%)	4 (1%)	76	76

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	67	ARG
1	B	203	LYS
1	B	228	PHE
1	B	253	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	222	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 10 ligands modelled in this entry, 1 is monoatomic - leaving 9 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	FUQ	B	302	-	0,29,30	-	-	-	-	-
6	PO4	B	305	-	4,4,4	0.92	0	6,6,6	0.43	0
3	8M0	A	303	-	14,46,48	8.46	9 (64%)	-	-	-
10	M27	A	305	-	0,11,12	-	-	-	-	-

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
7	ATP	A	301	8	26,33,33	0.93	1 (3%)	31,52,52	1.40	5 (16%)
3	8M0	B	301	-	22,48,48	8.02	12 (54%)	-		
6	PO4	B	304	-	4,4,4	0.91	0	6,6,6	0.42	0
9	MOO	A	304	-	2,4,4	9.97	1 (50%)	-		
5	GX2	B	303	-	7,25,77	2.67	2 (28%)	-		

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	ATP	A	301	8	-	1/18/38/38	0/3/3/3
4	FUQ	B	302	-	-	-	0/0/1/1
5	GX2	B	303	-	-	-	0/1/1/17
10	M27	A	305	-	-	-	0/1/1/1

All (25) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	301	8M0	O7-MO4	19.09	1.92	1.74
3	A	303	8M0	O20-MO8	18.48	1.92	1.74
3	B	301	8M0	O20-MO8	15.87	1.89	1.74
3	B	301	8M0	O9-MO3	15.65	1.90	1.71
3	A	303	8M0	O7-MO4	15.15	1.89	1.74
9	A	304	MOO	O2-MO	14.08	2.03	1.73
3	A	303	8M0	O23-MO7	11.20	1.88	1.68
3	B	301	8M0	O23-MO7	11.13	1.87	1.68
3	B	301	8M0	O10-MO3	11.07	1.85	1.71
3	A	303	8M0	O9-MO3	10.55	1.90	1.67
3	B	301	8M0	O1-MO1	10.29	1.86	1.68
3	A	303	8M0	O1-MO1	10.23	1.86	1.68
3	B	301	8M0	O28-MO5	8.29	1.81	1.71
3	A	303	8M0	O18-MO6	6.86	1.80	1.68
3	B	301	8M0	O18-MO6	6.42	1.79	1.68
3	B	301	8M0	O26-MO4	5.84	2.18	1.88
5	B	303	GX2	O35-MO2	5.04	1.89	1.73
3	B	301	8M0	O25-MO8	4.88	2.13	1.88
5	B	303	GX2	O43-MO2	-4.53	1.59	1.73
3	A	303	8M0	O13-MO5	4.46	1.77	1.67
3	B	301	8M0	O13-MO5	4.40	1.76	1.71

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	301	8M0	O8-MO2	3.47	1.74	1.68
3	A	303	8M0	O8-MO2	3.45	1.74	1.68
3	A	303	8M0	O3-MO3	2.98	2.00	1.89
7	A	301	ATP	C5-C4	2.61	1.47	1.40

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
7	A	301	ATP	N3-C2-N1	-3.24	123.61	128.68
7	A	301	ATP	C3'-C2'-C1'	3.17	105.76	100.98
7	A	301	ATP	PB-O3B-PG	-2.79	123.24	132.83
7	A	301	ATP	C4-C5-N7	-2.45	106.84	109.40
7	A	301	ATP	PA-O3A-PB	-2.19	125.32	132.83

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
7	A	301	ATP	C3'-C4'-C5'-O5'



There are no ring outliers.


3 monomers are involved in 6 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	303	8M0	3	0
7	A	301	ATP	1	0
9	A	304	MOO	1	1

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	B	268/269 (99%)	0.62	26 (9%) <b>7</b> <b>9</b>	24, 36, 64, 76	0
2	A	244/275 (88%)	0.15	12 (4%) <b>29</b> <b>33</b>	21, 30, 56, 71	0
All	All	512/544 (94%)	0.39	38 (7%) <b>14</b> <b>16</b>	21, 33, 61, 76	0

All (38) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	131	LEU	14.3
1	B	130	GLY	14.0
1	B	128	GLY	11.8
1	B	3	ASN	10.2
1	B	129	ALA	8.6
1	B	222	HIS	7.4
2	A	203	PRO	6.4
1	B	204	ASP	5.9
1	B	203	LYS	5.8
1	B	6	ALA	5.6
1	B	5	THR	5.1
1	B	221	LEU	5.0
1	B	228	PHE	4.9
1	B	223	ASP	4.7
1	B	201	THR	4.7
1	B	7	GLU	4.6
1	B	197	ALA	4.4
1	B	202	SER	4.1
2	A	276	ALA	4.1
2	A	207	GLN	4.0
1	B	4	SER	3.9
1	B	127	GLY	3.6
2	A	204	ASP	3.4
2	A	205	ARG	3.3

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
2	A	208	ALA	3.2
1	B	199	PRO	3.1
2	A	202	GLY	3.0
2	A	206	GLY	3.0
2	A	223	GLU	3.0
1	B	200	LYS	3.0
2	A	274	ARG	2.7
1	B	198	ASN	2.7
2	A	209	ARG	2.5
1	B	132	SER	2.4
2	A	275	PRO	2.4
1	B	205	ALA	2.4
1	B	10	GLU	2.2
1	B	219	LYS	2.1

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
6	PO4	B	305	5/5	0.35	0.63	72,80,86,92	0
6	PO4	B	304	5/5	0.42	0.78	82,85,88,89	0
5	GX2	B	303	24/61	0.42	0.33	61,69,96,103	24
4	FUQ	B	302	29/30	0.81	0.29	55,63,67,70	29
3	8M0	B	301	36/36	0.82	0.20	44,49,54,54	36
8	MG	A	302	1/1	0.86	0.20	33,33,33,33	0
10	M27	A	305	11/12	0.93	0.38	39,41,46,48	11
7	ATP	A	301	31/31	0.96	0.12	25,34,39,42	0
3	8M0	A	303	34/36	0.98	0.17	22,31,41,48	34

*Continued on next page...*

Continued from previous page...




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
9	MOO	A	304	5/5	0.99	0.11	21,25,30,30	1

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.







**Electron density around 8M0 B 301:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around ATP A 301:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





## 6.5 Other polymers [i](#)

There are no such residues in this entry.