

Full wwPDB X-ray Structure Validation Report (i)

May 20, 2024 – 02:45 AM JST

PDB ID : 8HJN

Title : Crystal structure of glycosyltransferase SgUGT94-289-3 in complex with UPG

Authors: Li, M.; Zhang, S.; Cui, S.

Deposited on : 2022-11-23

Resolution : 3.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity : 4.02b-467

Mogul : 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : 1.13

EDS : 2.36.2buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

Refmac : 5.8.0158

CCP4 : 7.0.044 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.36.2

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X- $RAY\ DIFFRACTION$

The reported resolution of this entry is 3.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive	Similar resolution
Metric	$(\# \mathrm{Entries})$	$(\# ext{Entries}, ext{ resolution range}(ext{Å}))$
R_{free}	130704	1133 (3.20-3.20)
Clashscore	141614	1253 (3.20-3.20)
Ramachandran outliers	138981	1234 (3.20-3.20)
Sidechain outliers	138945	1233 (3.20-3.20)
RSRZ outliers	127900	1095 (3.20-3.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain		
1	A	456	79%	15%	6%
1	В	456	72%	21%	7%



2 Entry composition (i)


There are 2 unique types of molecules in this entry. The entry contains 6821 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called glycosyltransferase.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace		
1	A	429	Total 3388	C 2191	N 567	O 614	S 16	0	1	0
1	В	426	Total 3361	C 2176	N 558	O 611	S 16	0	1	0

• Molecule 2 is URIDINE-5'-DIPHOSPHATE-GLUCOSE (three-letter code: UPG) (formula: $C_{15}H_{24}N_2O_{17}P_2$) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf			
2	Λ	1	Total	С	N	О	Р	0	0	
2	A	1	36	15	2	17	2	U	0	
2	D	1	Total	С	N	О	Р	0	0	
	Б	1	36	15	2	17	2	U		



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: glycosyltransferase





4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants	64.22Å 80.09Å 172.51Å	Donositon
a, b, c, α , β , γ	90.00° 90.00° 90.00°	Depositor
Resolution (Å)	51.51 - 3.20	Depositor
Resolution (A)	51.51 - 3.20	EDS
% Data completeness	90.1 (51.51-3.20)	Depositor
(in resolution range)	90.1 (51.51-3.20)	EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	1.63 (at 3.19Å)	Xtriage
Refinement program	PHENIX 1.18.2_3874	Depositor
D D.	0.235 , 0.277	Depositor
R, R_{free}	0.235 , 0.277	DCC
R_{free} test set	718 reflections (5.21%)	wwPDB-VP
Wilson B-factor (Å ²)	63.2	Xtriage
Anisotropy	0.409	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.33 , 57.0	EDS
L-test for twinning ²	$ < L >=0.44, < L^2>=0.27$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	6821	wwPDB-VP
Average B, all atoms (Å ²)	53.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The analyses of the Patterson function reveals a significant off-origin peak that is 55.08 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 3.3320e-05. The detected translational NCS is most likely also responsible for the elevated intensity ratio.

²Theoretical values of <|L|>, $<L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: UPG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond angles		
IVIOI	Chain	RMSZ	# Z > 5	RMSZ	# Z > 5	
1	A	0.28	0/3485	0.53	1/4727 (0.0%)	
1	В	0.29	0/3455	0.55	2/4687 (0.0%)	
All	All	0.28	0/6940	0.54	3/9414 (0.0%)	

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	В	0	1

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$Observed(^o)$	$\operatorname{Ideal}({}^{o})$
1	В	9	THR	CA-C-N	-8.60	98.28	117.20
1	A	18	LEU	C-N-CA	-5.16	111.46	122.30
1	В	113	LEU	CB-CG-CD2	-5.07	102.38	111.00

There are no chirality outliers.

All (1) planarity outliers are listed below:

\mathbf{Mol}	Chain	Res	Type	Group
1	В	9	THR	Mainchain



5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3388	0	3312	42	0
1	В	3361	0	3293	66	0
2	A	36	0	22	0	0
2	В	36	0	22	3	0
All	All	6821	0	6649	107	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (107) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

A 4 1	A 4 0	Interatomic	Clash
Atom-1	Atom-2	${\rm distance}(\mathring{\rm A})$	overlap (Å)
1:A:137:PRO:HG3	1:A:453:GLU:HG3	1.53	0.88
1:B:146:VAL:HG22	1:B:209:ILE:HD13	1.56	0.88
1:A:282:GLU:HG3	1:A:315:PRO:HG3	1.57	0.85
1:B:282:GLU:HG3	1:B:315:PRO:HG3	1.59	0.83
1:B:23:LEU:HD11	1:B:48:ASN:HB3	1.61	0.82
1:A:203:SER:HA	1:A:449:LEU:HD11	1.69	0.74
1:B:203:SER:HA	1:B:449:LEU:HD11	1.71	0.72
1:B:148:VAL:HG11	1:B:374:ASP:HB2	1.72	0.71
1:B:373:VAL:HG21	2:B:501:UPG:H6'2	1.73	0.71
1:A:78:PRO:HG2	1:A:80:HIS:CD2	2.25	0.71
1:B:414:GLU:HG3	1:B:416:THR:HG23	1.74	0.68
1:A:148:VAL:HG21	1:A:374:ASP:HB2	1.75	0.67
1:B:347:PHE:CE2	1:B:349:SER:HB2	2.29	0.67
1:B:72:PRO:HD2	1:B:98:ALA:HB1	1.77	0.67
1:B:27:LEU:HD22	1:B:52:ILE:HG13	1.77	0.65
1:B:392:LYS:O	1:B:400:GLN:NE2	2.30	0.64
1:A:207:ILE:HD11	1:A:231:VAL:HG22	1.80	0.64
1:A:206:VAL:HG12	1:A:230:LYS:HB2	1.81	0.63
1:B:76:GLU:O	1:B:78:PRO:HD3	2.02	0.60
1:B:97:GLN:HG3	1:B:101:MET:HE2	1.84	0.59
1:B:138:ALA:O	1:B:203:SER:HB2	2.03	0.58
1:B:367:ILE:HD13	1:B:408:ILE:HG12	1.84	0.58
1:B:267:PHE:HD2	1:B:347:PHE:HE1	1.50	0.58



Continued from previous page...

Continued from pred		Interatomic	Clash
Atom-1	Atom-2	${\rm distance}(\mathring{\rm A})$	overlap (Å)
1:A:148:VAL:HG22	1:A:377:PHE:HB2	1.84	0.58
1:B:56:LEU:HD11	1:B:64:ILE:HD12	1.86	0.57
1:A:138:ALA:O	1:A:203:SER:HB3	2.05	0.56
1:B:23:LEU:HD11	1:B:48:ASN:CB	2.34	0.56
1:B:137:PRO:HB3	1:B:449:LEU:HD21	1.87	0.56
1:A:436:GLU:HG2	1:B:436:GLU:HG2	1.88	0.56
1:B:296:ILE:HD13	1:B:327:MET:HB3	1.88	0.54
1:A:425:ARG:O	1:A:429:GLU:HG2	2.07	0.54
1:B:148:VAL:HG12	1:B:377:PHE:HB2	1.89	0.54
1:A:240:GLU:OE1	1:A:432:ARG:NH1	2.41	0.54
1:B:267:PHE:CD2	1:B:347:PHE:HE1	2.25	0.54
1:B:359:SER:OG	1:B:364:VAL:O	2.25	0.53
1:B:409:LYS:O	1:B:414:GLU:HG2	2.09	0.53
1:A:248:GLU:HG2	1:A:249:GLY:N	2.24	0.52
1:B:333:ALA:O	2:B:501:UPG:H5	2.09	0.52
1:B:267:PHE:HD2	1:B:347:PHE:CE1	2.27	0.52
1:B:409:LYS:HA	1:B:413:VAL:HB	1.91	0.51
1:A:153:LEU:HA	1:A:156:ILE:HD13	1.92	0.51
1:B:276:PRO:HB2	1:B:281:MET:HG3	1.93	0.51
1:B:16:PRO:HD2	1:B:43:CYS:O	2.12	0.50
1:B:448:LEU:O	1:B:452:ILE:HG13	2.12	0.50
1:B:339:LEU:HD21	1:B:347:PHE:CD1	2.47	0.50
1:A:270:PHE:CZ	1:A:284:ILE:HD13	2.47	0.49
1:A:53:LYS:N	1:A:54:PRO:HD2	2.27	0.49
1:A:236:PRO:HG3	1:A:431:LEU:HD22	1.95	0.49
1:A:53:LYS:HD2	1:A:53:LYS:O	2.12	0.49
1:B:121:ASP:OD1	1:B:122:SER:N	2.43	0.49
1:A:117:LEU:HD11	1:A:139:ILE:HG13	1.94	0.48
1:B:30:ALA:HB1	1:B:40:ILE:HG21	1.94	0.48
1:B:137:PRO:HG3	1:B:453:GLU:OE1	2.14	0.48
1:B:270:PHE:CZ	1:B:284:ILE:HD13	2.49	0.48
1:B:17:TRP:HB3	1:B:121:ASP:OD2	2.14	0.47
1:B:142:ASN:HB3	1:B:209:ILE:HG13	1.97	0.47
1:B:276:PRO:HG2	1:B:281:MET:SD	2.55	0.47
1:B:350:HIS:O	1:B:375:GLN:HB3	2.15	0.46
1:B:347:PHE:CD2	1:B:349:SER:HB2	2.51	0.46
1:A:168:PHE:HE2	1:A:377:PHE:HD1	1.63	0.45
1:A:410:GLU:HB3	1:A:417:ARG:HG3	1.98	0.45
1:B:116:HIS:O	1:B:137:PRO:HD2	2.16	0.45
1:A:146:VAL:CG2	1:A:209:ILE:HG12	2.46	0.45
1:B:319:LEU:O	1:B:322:ALA:O	2.35	0.45



Continued from previous page...

Continued from pred		Interatomic	Clash
Atom-1	Atom-2	${\rm distance} \ (\mathring{\rm A})$	$overlap (\AA)$
1:A:276:PRO:HB2	1:A:281:MET:HG3	1.99	0.45
1:A:142:ASN:HB3	1:A:209:ILE:HD12	1.99	0.45
1:A:294:ASN:HA	1:A:325:ARG:O	2.17	0.45
1:B:256:TRP:CZ3	1:B:296:ILE:HD11	2.51	0.44
1:A:137:PRO:HB3	1:A:449:LEU:HD21	1.98	0.44
1:A:448:LEU:O	1:A:452:ILE:HG13	2.17	0.44
1:B:9:THR:HG22	1:B:9:THR:O	2.18	0.44
1:B:442:MET:O	1:B:446:ILE:HG12	2.17	0.44
1:A:27:LEU:HD22	1:A:52:ILE:HG13	2.00	0.44
1:A:419:ASP:OD1	1:A:419:ASP:N	2.50	0.44
1:A:238:VAL:HG13	1:A:358:GLU:HG2	2.00	0.43
1:B:14:MET:HE3	1:B:29:LEU:HD23	2.00	0.43
1:A:409:LYS:O	1:A:414:GLU:HG2	2.18	0.43
1:A:79:PRO:HA	1:A:82:HIS:CG	2.54	0.43
1:A:283:GLU:OE2	1:A:401:ARG:HG3	2.19	0.43
1:A:426:GLU:O	1:A:430:ILE:HG13	2.18	0.43
1:B:78:PRO:HB2	1:B:80:HIS:CD2	2.54	0.43
1:B:33:LEU:O	1:B:38:PHE:HB2	2.18	0.43
1:A:267:PHE:O	1:A:347:PHE:HD1	2.02	0.43
1:B:106:PHE:HD2	1:B:130:VAL:HG21	1.83	0.43
1:B:49:LEU:HD11	1:B:67:VAL:O	2.19	0.42
1:B:428:SER:O	1:B:432:ARG:HG2	2.19	0.42
1:B:439:PHE:O	1:B:443:VAL:HG23	2.20	0.42
1:A:201:HIS:CD2	1:A:203:SER:HB2	2.54	0.42
1:B:121:ASP:H	1:B:124:GLN:NE2	2.18	0.42
2:B:501:UPG:H6'1	2:B:501:UPG:H1'	1.78	0.42
1:B:88:PRO:HG3	1:B:302:PRO:HB3	2.02	0.42
1:B:386:GLY:O	1:B:420:VAL:HA	2.20	0.42
1:A:35:ARG:O	1:B:36:ARG:HG2	2.20	0.41
1:B:80:HIS:CE1	1:B:81:LEU:HG	2.55	0.41
1:B:272:SER:HA	1:B:300:ARG:HH11	1.84	0.41
1:B:372:HIS:CD2	1:B:373:VAL:HG13	2.55	0.41
1:B:382:VAL:CG1	1:B:388:GLY:HA3	2.49	0.41
1:A:365:PRO:HA	1:A:387:VAL:O	2.21	0.41
1:A:318:PHE:CD2	1:A:319:LEU:HD12	2.56	0.41
1:A:78:PRO:HG2	1:A:80:HIS:NE2	2.34	0.41
1:A:375:GLN:N	1:A:376:PRO:HD2	2.36	0.41
1:B:153:LEU:HD23	1:B:153:LEU:HA	1.94	0.41
1:B:113:LEU:HD23	1:B:113:LEU:HA	1.78	0.41
1:B:207:ILE:HD11	1:B:231:VAL:HG22	2.03	0.41
1:B:448:LEU:HG	1:B:452:ILE:HD11	2.03	0.40



Continued from previous page...

Atom-1	Atom-2	$\begin{array}{c} {\rm Interatomic} \\ {\rm distance} \ ({\rm \AA}) \end{array}$	Clash overlap (Å)
1:B:294:ASN:HA	1:B:325:ARG:O	2.20	0.40
1:A:270:PHE:HZ	1:A:284:ILE:HD13	1.86	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percei	ntiles
1	A	422/456 (92%)	409 (97%)	12 (3%)	1 (0%)	47	79
1	В	419/456 (92%)	411 (98%)	7 (2%)	1 (0%)	47	79
All	All	841/912 (92%)	820 (98%)	19 (2%)	2 (0%)	47	79

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	373	VAL
1	В	373	VAL

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles		
1	A	369/396 (93%)	369 (100%)	0	100	100	
1	В	366/396 (92%)	366 (100%)	0	100	100	



Continued from previous page...

Mol	Chain	Analysed	Rotameric				
All	All	735/792 (93%)	735 (100%)	0	100	100	

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	173	HIS
1	A	201	HIS
1	В	82	HIS
1	В	97	GLN
1	В	201	HIS
1	В	372	HIS

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).



Mol	Type Chain		Chain Res		Bond lengths			Bond angles		
IVIOI	Type	Chain	nes	Link	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	UPG	A	501	-	35,38,38	0.37	0	53,58,58	0.51	0
2	UPG	В	501	-	35,38,38	0.39	0	53,58,58	0.54	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

\mathbf{Mol}	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	UPG	A	501	-	-	1/23/59/59	0/3/3/3
2	UPG	В	501	-	-	9/23/59/59	0/3/3/3

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (10) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	В	501	UPG	O4C-C1C-N1-C2
2	В	501	UPG	O4C-C1C-N1-C6
2	В	501	UPG	C5C-O5C-PA-O2A
2	В	501	UPG	PA-O3A-PB-O3B
2	В	501	UPG	C1'-O3B-PB-O2B
2	В	501	UPG	C1'-O3B-PB-O3A
2	В	501	UPG	C5C-O5C-PA-O3A
2	В	501	UPG	C1'-O3B-PB-O1B
2	В	501	UPG	C5C-O5C-PA-O1A
2	A	501	UPG	O4C-C4C-C5C-O5C

There are no ring outliers.


1 monomer is involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	В	501	UPG	3	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier.



Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ $>$	$\#\mathrm{RSRZ}{>}2$		$\mathbf{OWAB}(\mathbf{\mathring{A}}^2)$	Q < 0.9
1	A	429/456 (94%)	0.01	6 (1%) 75	63	22, 52, 87, 121	0
1	В	$426/456 \ (93\%)$	0.01	9 (2%) 63	49	26, 47, 88, 116	0
All	All	855/912 (93%)	0.01	15 (1%) 68	55	22, 50, 88, 121	0

All (15) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	В	201	HIS	3.8
1	A	456	HIS	3.5
1	В	203	SER	3.3
1	A	193	GLY	2.9
1	В	160	HIS	2.7
1	В	199	CYS	2.6
1	В	202	ALA	2.5
1	В	274	TYR	2.4
1	В	161	SER	2.4
1	В	324	GLU	2.3
1	В	116	HIS	2.1
1	A	176	ALA	2.1
1	A	305	ASP	2.1
1	A	330	LYS	2.0
1	A	324	GLU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.




6.4 Ligands (i)


In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-}\mathbf{factors}(\mathbf{\mathring{A}}^2)$	Q<0.9
2	UPG	В	501	36/36	0.89	0.26	26,54,80,102	0
2	UPG	A	501	36/36	0.91	0.30	35,57,91,105	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.







6.5 Other polymers (i)

There are no such residues in this entry.

