



# Full wwPDB X-ray Structure Validation Report ⓘ

Oct 3, 2021 – 12:08 PM EDT

PDB ID : 3HK0  
Title : Crystal Structure of the RA and PH Domains of Grb10  
Authors : Hubbard, S.R.; Depetris, R.S.; Wu, J.  
Deposited on : 2009-05-22  
Resolution : 2.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.23.2  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.23.2


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	3163 (2.60-2.60)
Clashscore	141614	3518 (2.60-2.60)
Ramachandran outliers	138981	3455 (2.60-2.60)
Sidechain outliers	138945	3455 (2.60-2.60)
RSRZ outliers	127900	3104 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	256	
1	B	256	

## 2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 3542 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Growth factor receptor-bound protein 10.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	230	1818	1162	314	331	11	0	0	0
1	B	216	1707	1096	290	309	12	0	0	0

There are 20 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	102	GLY	-	expression tag	UNP Q13322
A	103	SER	-	expression tag	UNP Q13322
A	104	HIS	-	expression tag	UNP Q13322
A	105	MET	-	expression tag	UNP Q13322
A	145	SER	CYS	engineered mutation	UNP Q13322
A	212	SER	CYS	engineered mutation	UNP Q13322
A	232	SER	CYS	engineered mutation	UNP Q13322
A	270	ALA	LYS	engineered mutation	UNP Q13322
A	271	ALA	GLU	engineered mutation	UNP Q13322
A	331	SER	CYS	engineered mutation	UNP Q13322
B	102	GLY	-	expression tag	UNP Q13322
B	103	SER	-	expression tag	UNP Q13322
B	104	HIS	-	expression tag	UNP Q13322
B	105	MET	-	expression tag	UNP Q13322
B	145	SER	CYS	engineered mutation	UNP Q13322
B	212	SER	CYS	engineered mutation	UNP Q13322
B	232	SER	CYS	engineered mutation	UNP Q13322
B	270	ALA	LYS	engineered mutation	UNP Q13322
B	271	ALA	GLU	engineered mutation	UNP Q13322
B	331	SER	CYS	engineered mutation	UNP Q13322

- Molecule 2 is THIOCYANATE ION (three-letter code: SCN) (formula: CNS).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	S		
2	A	1	3	1	1	1	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
3	A	7	7	7	0	0
3	B	7	7	7	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Growth factor receptor-bound protein 10



- Molecule 1: Growth factor receptor-bound protein 10



## 4 Data and refinement statistics i

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	126.88Å 48.88Å 92.22Å 90.00° 114.14° 90.00°	Depositor
Resolution (Å)	30.00 – 2.60 43.54 – 2.50	Depositor EDS
% Data completeness (in resolution range)	99.6 (30.00-2.60) 98.2 (43.54-2.50)	Depositor EDS
$R_{merge}$	0.06	Depositor
$R_{sym}$	0.06	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.81 (at 2.51Å)	Xtrriage
Refinement program	REFMAC 5.2.0019	Depositor
R, $R_{free}$	0.230 , 0.277 0.227 , 0.271	Depositor DCC
$R_{free}$ test set	907 reflections (5.10%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	59.7	Xtrriage
Anisotropy	0.278	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.31 , 52.3	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	3542	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	48.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.13% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SCN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.37	0/1857	0.53	0/2513
1	B	0.51	3/1743 (0.2%)	0.60	3/2358 (0.1%)
All	All	0.44	3/3600 (0.1%)	0.56	3/4871 (0.1%)

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	253	TYR	CG-CD2	9.04	1.50	1.39
1	B	253	TYR	CE1-CZ	7.98	1.49	1.38
1	B	253	TYR	CE2-CZ	7.02	1.47	1.38

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	253	TYR	CZ-CE2-CD2	-7.29	113.24	119.80
1	B	253	TYR	CB-CG-CD1	-5.49	117.70	121.00
1	B	304	LEU	CA-CB-CG	5.09	127.00	115.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1818	0	1749	17	0
1	B	1707	0	1633	16	0
2	A	3	0	0	0	0
3	A	7	0	0	0	0
3	B	7	0	0	0	0
All	All	3542	0	3382	32	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (32) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:286:ILE:HD11	1:A:304:LEU:HD12	1.72	0.70
1:A:284:SER:HB3	1:A:308:PRO:HA	1.76	0.67
1:B:311:VAL:HG13	1:B:314:GLU:HB2	1.78	0.65
1:B:260:GLY:HA2	1:B:281:LEU:HG	1.80	0.62
1:A:161:GLY:HA3	1:A:291:ALA:HB2	1.82	0.61
1:A:151:TRP:CE2	1:A:189:LYS:HE3	2.38	0.59
1:B:323:ALA:HB1	1:B:328:THR:HG23	1.83	0.59
1:A:156:HIS:CE1	1:A:290:ILE:HD13	2.42	0.55
1:B:236:GLN:HE21	1:B:253:TYR:HE2	1.55	0.55
1:B:118:ASP:N	1:B:118:ASP:OD1	2.39	0.55
1:A:190:ASN:ND2	1:A:192:ALA:H	2.12	0.48
1:B:302:HIS:CD2	1:B:326:GLU:HG2	2.49	0.47
1:B:315:THR:HA	1:B:318:LEU:HD13	1.96	0.47
1:A:240:HIS:HB2	1:A:322:CYS:HB2	1.97	0.47
1:A:260:GLY:HA2	1:A:281:LEU:HG	1.98	0.45
1:B:236:GLN:HB3	1:B:253:TYR:OH	2.16	0.45
1:A:315:THR:O	1:A:318:LEU:HB2	2.16	0.45
1:B:154:VAL:HB	1:B:186:LEU:HB2	1.98	0.44
1:A:194:TYR:HB3	1:A:197:PHE:HD1	1.81	0.44
1:A:343:MET:HG2	1:B:169:HIS:HD2	1.82	0.44
1:B:169:HIS:HB3	1:B:348:ASN:HD21	1.83	0.44
1:B:286:ILE:HD11	1:B:304:LEU:HD23	2.00	0.44
1:B:152:THR:HG22	1:B:188:ARG:O	2.17	0.43
1:A:244:LEU:HD13	1:A:319:ARG:CZ	2.49	0.42
1:A:154:VAL:HB	1:A:186:LEU:HB2	2.01	0.42
1:A:211:TRP:CZ3	1:A:225:ASN:HB3	2.55	0.42
1:A:334:THR:HG22	1:A:338:LEU:HD22	2.02	0.42
1:B:154:VAL:HG21	1:B:188:ARG:CZ	2.49	0.42
1:B:240:HIS:HB2	1:B:322:CYS:HB2	2.02	0.41

*Continued on next page...*



Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:135:LEU:HD12	1:A:172:VAL:HG21	2.03	0.41
1:A:212:SER:OG	1:A:213:GLN:N	2.53	0.41
1:B:326:GLU:OE2	1:B:329:ARG:NH2	2.54	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	224/256 (88%)	216 (96%)	7 (3%)	1 (0%)	34 57
1	B	208/256 (81%)	200 (96%)	7 (3%)	1 (0%)	29 52
All	All	432/512 (84%)	416 (96%)	14 (3%)	2 (0%)	29 52

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	292	GLY
1	A	213	GLN

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	191/230 (83%)	177 (93%)	14 (7%)	14 28

Continued on next page...

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	179/230 (78%)	166 (93%)	13 (7%)	14	28
All	All	370/460 (80%)	343 (93%)	27 (7%)	14	28

All (27) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	112	VAL
1	A	118	ASP
1	A	122	LYS
1	A	135	LEU
1	A	190	ASN
1	A	244	LEU
1	A	256	LEU
1	A	273	ARG
1	A	284	SER
1	A	290	ILE
1	A	304	LEU
1	A	318	LEU
1	A	319	ARG
1	A	338	LEU
1	B	118	ASP
1	B	122	LYS
1	B	133	ARG
1	B	135	LEU
1	B	171	LEU
1	B	244	LEU
1	B	256	LEU
1	B	297	ASN
1	B	301	ASP
1	B	304	LEU
1	B	319	ARG
1	B	328	THR
1	B	338	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	190	ASN
1	A	224	GLN
1	A	347	GLN
1	A	348	ASN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	174	GLN
1	B	302	HIS
1	B	347	GLN
1	B	348	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	SCN	A	15	-	1,2,2	1.84	0	0,1,1	-	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	230/256 (89%)	-0.17	2 (0%) 84 82	41, 48, 55, 57	0
1	B	216/256 (84%)	-0.04	3 (1%) 75 71	41, 49, 57, 59	0
All	All	446/512 (87%)	-0.11	5 (1%) 80 78	41, 49, 56, 59	0

All (5) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	126	ILE	2.6
1	A	141	TYR	2.6
1	B	277	LEU	2.2
1	B	275	LEU	2.1
1	B	296	TYR	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	SCN	A	15	3/3	0.97	0.17	55,55,55,55	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.