



Full wwPDB X-ray Structure Validation Report ⓘ

May 16, 2020 – 02:50 am BST

PDB ID : 1HL8
Title : CRYSTAL STRUCTURE OF THERMOTOGA MARITIMA ALPHA-FUCOSIDASE
Authors : Sulzenbacher, G.; Bignon, C.; Bourne, Y.; Henrissat, B.
Deposited on : 2003-03-14
Resolution : 2.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3907 (2.40-2.40)
Clashscore	141614	4398 (2.40-2.40)
Ramachandran outliers	138981	4318 (2.40-2.40)
Sidechain outliers	138945	4319 (2.40-2.40)
RSRZ outliers	127900	3811 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	449	 3% 83% 11% • 5%
1	B	449	 4% 80% 13% • 6%

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 7234 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called PUTATIVE ALPHA-L-FUCOSIDASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	426	Total 3552	C 2315	N 587	O 643	S 7	0	4	0
1	B	422	Total 3520	C 2296	N 582	O 635	S 7	0	3	0


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	87	Total 87	O 87	0	0
2	B	75	Total 75	O 75	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: PUTATIVE ALPHA-L-FUCOSIDASE



- Molecule 1: PUTATIVE ALPHA-L-FUCOSIDASE



4 Data and refinement statistics

Property	Value	Source
Space group	H 3 2	Depositor
Cell constants a, b, c, α , β , γ	172.46Å 172.46Å 167.41Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	37.27 – 2.40 37.15 – 2.40	Depositor EDS
% Data completeness (in resolution range)	99.1 (37.27-2.40) 99.0 (37.15-2.40)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.91 (at 2.39Å)	Xtrriage
Refinement program	REFMAC 5.1.24	Depositor
R, R_{free}	0.185 , 0.228 0.214 , 0.245	Depositor DCC
R_{free} test set	2642 reflections (7.13%)	wwPDB-VP
Wilson B-factor (Å ²)	41.6	Xtrriage
Anisotropy	0.018	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 48.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	7234	wwPDB-VP
Average B, all atoms (Å ²)	60.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.80% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: KCX

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.42	0/3669	0.73	11/4978 (0.2%)
1	B	0.40	0/3631	0.72	12/4924 (0.2%)
All	All	0.41	0/7300	0.72	23/9902 (0.2%)

There are no bond length outliers.

All (23) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	219	ASP	CB-CG-OD2	6.94	124.55	118.30
1	A	56	ASP	CB-CG-OD2	6.58	124.22	118.30
1	A	98	ASP	CB-CG-OD2	6.02	123.72	118.30
1	B	177	ASP	CB-CG-OD2	5.93	123.64	118.30
1	A	152	ASP	CB-CG-OD2	5.87	123.59	118.30
1	A	177	ASP	CB-CG-OD2	5.81	123.53	118.30
1	A	112	ASP	CB-CG-OD2	5.50	123.25	118.30
1	B	112	ASP	CB-CG-OD2	5.45	123.21	118.30
1	B	313	ASP	CB-CG-OD2	5.43	123.19	118.30
1	B	11	ASP	CB-CG-OD2	5.41	123.17	118.30
1	A	400	ASP	CB-CG-OD2	5.35	123.12	118.30
1	A	11	ASP	CB-CG-OD2	5.33	123.10	118.30
1	B	162	ARG	NE-CZ-NH1	5.26	122.93	120.30
1	B	330	ASP	CB-CG-OD2	5.25	123.02	118.30
1	A	274	ASP	CB-CG-OD2	5.24	123.02	118.30
1	B	25	ASP	CB-CG-OD2	5.18	122.96	118.30
1	B	98	ASP	CB-CG-OD2	5.14	122.93	118.30
1	B	253	ASP	CB-CG-OD2	5.14	122.92	118.30
1	A	253	ASP	CB-CG-OD2	5.12	122.91	118.30
1	B	219	ASP	CB-CG-OD2	5.12	122.91	118.30
1	A	190	ASP	CB-CG-OD2	5.11	122.89	118.30
1	B	106	ASP	CB-CG-OD2	5.01	122.81	118.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	224	ASP	CB-CG-OD2	5.01	122.81	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3552	0	3422	8	0
1	B	3520	0	3392	14	0
2	A	87	0	0	0	0
2	B	75	0	0	0	0
All	All	7234	0	6814	22	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (22) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:18:HIS:CD2	1:B:243:TYR:CZ	2.85	0.65
1:B:18:HIS:HE1	1:B:247:PRO:O	1.80	0.63
1:A:32:PHE:CD2	1:A:322:LEU:HD13	2.39	0.58
1:B:18:HIS:CD2	1:B:243:TYR:CE1	2.93	0.56
1:A:432:LYS:HA	1:A:435:LEU:HD12	1.91	0.53
1:B:296:GLU:HA	1:B:300:HIS:CE1	2.44	0.52
1:A:396:ILE:HB	1:A:428:ILE:HG23	1.92	0.51
1:A:299:GLU:OE1	1:A:300:HIS:CE1	2.64	0.51
1:B:58:TRP:HZ2	1:B:64:TYR:CE2	2.29	0.50
1:B:58:TRP:HZ2	1:B:64:TYR:HE2	1.56	0.50
1:B:142:PHE:CE2	1:B:179:ARG:HD2	2.47	0.49
1:B:124:PRO:HD2	1:B:169:GLY:O	2.16	0.46
1:B:134:LEU:HD13	1:B:153:LEU:HD12	2.00	0.44
1:B:408:VAL:HG22	1:B:446:ALA:HB2	1.99	0.44
1:B:292:TYR:CE2	1:B:328:LYS:HG2	2.54	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:396:ILE:HD12	1:A:430:VAL:HG23	2.01	0.41
1:B:191:LEU:O	1:B:195:ARG:NE	2.52	0.41
1:B:363:ARG:NH2	1:B:400:ASP:O	2.53	0.41
1:B:300:HIS:CD2	1:B:300:HIS:N	2.87	0.41
1:A:18:HIS:CE1	1:A:243:TYR:CE1	3.09	0.40
1:A:276:PRO:HB3	1:A:278:TYR:CE1	2.57	0.40
1:A:40:VAL:HB	1:A:41:PRO:CD	2.52	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	423/449 (94%)	403 (95%)	18 (4%)	2 (0%)	29 41
1	B	416/449 (93%)	395 (95%)	20 (5%)	1 (0%)	47 62
All	All	839/898 (93%)	798 (95%)	38 (4%)	3 (0%)	34 48

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	405	ALA
1	B	405	ALA
1	A	440	ILE

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	372/388 (96%)	340 (91%)	32 (9%)	10	16
1	B	368/388 (95%)	335 (91%)	33 (9%)	9	14
All	All	740/776 (95%)	675 (91%)	65 (9%)	10	15

All (65) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	14[A]	SER
1	A	14[B]	SER
1	A	28	LYS
1	A	32	PHE
1	A	45	THR
1	A	75	LYS
1	A	85	LYS
1	A	94	GLU
1	A	99	LEU
1	A	112	ASP
1	A	113	LEU
1	A	134	LEU
1	A	138[A]	LYS
1	A	138[B]	LYS
1	A	176	LEU
1	A	179	ARG
1	A	215	LEU
1	A	263	LYS
1	A	266	GLU
1	A	275	LEU
1	A	299	GLU
1	A	311	LEU
1	A	322	LEU
1	A	362	GLU
1	A	370	ASP
1	A	379	LYS
1	A	395	LYS
1	A	404	SER
1	A	409	ARG
1	A	416	ARG
1	A	424	LYS
1	A	430	VAL
1	B	7	ARG
1	B	16[A]	ARG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	16[B]	ARG
1	B	32	PHE
1	B	45	THR
1	B	66	GLU
1	B	76	GLU
1	B	85	LYS
1	B	113	LEU
1	B	134	LEU
1	B	215	LEU
1	B	221	LEU
1	B	222	TRP
1	B	251	VAL
1	B	294	ARG
1	B	300	HIS
1	B	301	MET
1	B	311	LEU
1	B	322	LEU
1	B	339	GLU
1	B	344	LEU
1	B	349	ARG
1	B	353	ASP
1	B	363	ARG
1	B	367	LYS
1	B	378	ARG
1	B	379	LYS
1	B	383	ILE
1	B	388	LEU
1	B	420	LYS
1	B	428	ILE
1	B	430	VAL
1	B	432	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	208	GLN
1	A	300	HIS
1	A	402	ASN
1	B	18	HIS
1	B	208	GLN
1	B	295	ASN
1	B	402	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	KCX	B	338	1	7,11,12	1.38	1 (14%)	4,12,14	0.74	0
1	KCX	A	338	1	7,11,12	1.21	1 (14%)	4,12,14	0.49	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	KCX	B	338	1	-	3/7/10/12	-
1	KCX	A	338	1	-	1/7/10/12	-

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	338	KCX	CE-NZ	3.08	1.52	1.45
1	A	338	KCX	CE-NZ	2.47	1.51	1.45

There are no bond angle outliers.

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	B	338	KCX	CG-CD-CE-NZ

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
1	B	338	KCX	CE-CD-CG-CB
1	B	338	KCX	C-CA-CB-CG
1	A	338	KCX	C-CA-CB-CG

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	425/449 (94%)	0.20	15 (3%) 44 43	30, 60, 85, 101	0
1	B	421/449 (93%)	0.27	16 (3%) 40 39	30, 61, 100, 121	0
All	All	846/898 (94%)	0.24	31 (3%) 41 41	30, 61, 93, 121	0

All (31) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	267	TYR	7.7
1	B	405	ALA	6.2
1	B	267	TYR	5.8
1	B	58	TRP	5.2
1	A	297	GLY	4.8
1	A	58	TRP	4.7
1	B	268	HIS	3.9
1	A	404	SER	3.7
1	B	413	THR	3.5
1	A	274	ASP	3.3
1	B	56	ASP	3.3
1	A	76	GLU	3.3
1	B	432	LYS	2.9
1	A	85	LYS	2.9
1	A	305	GLU	2.9
1	B	380	CYS	2.8
1	A	381	ASN	2.7
1	A	411	PHE	2.5
1	A	84	VAL	2.5
1	B	403	LEU	2.5
1	A	299	GLU	2.5
1	B	85	LYS	2.4
1	A	372	THR	2.4
1	A	56	ASP	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	351	TYR	2.3
1	B	430	VAL	2.2
1	B	372	THR	2.2
1	B	406	GLY	2.1
1	B	383	ILE	2.1
1	B	45	THR	2.1
1	B	418	SER	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	KCX	B	338	12/13	0.82	0.20	80,84,86,87	0
1	KCX	A	338	12/13	0.85	0.25	81,83,86,86	0

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.