



Full wwPDB X-ray Structure Validation Report

Sep 20, 2023 – 07:59 PM EDT

PDB ID : 5HNM
Title : Crystal structure of vancomycin resistance D,D-pentapeptidase VanY E175A mutant from VanB-type resistance cassette in complex with Zn(II)
Authors : Stogios, P.J.; Chun, J.; Wawrzak, Z.; Evdokimova, E.; Di Leo, R.; Yim, V.; Courvalin, P.; Savchenko, A.; Anderson, W.F.; Center for Structural Genomics of Infectious Diseases (CSGID)
Deposited on : 2016-01-18
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the  symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references](#) ) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.35.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION





The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.





Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5042 (2.30-2.30)
Clashscore	141614	5643 (2.30-2.30)
Ramachandran outliers	138981	5575 (2.30-2.30)
Sidechain outliers	138945	5575 (2.30-2.30)
RSRZ outliers	127900	4938 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	197	 5% 85% 6% 9%
1	B	197	 3% 87% 5% 8%
1	C	197	 6% 83% 7% 10%
1	D	197	 8% 82% 9% 9%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	E	197	 4% 85% 6% 9%
1	F	197	 6% 85% 6% 9%

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 9261 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called D-alanyl-D-alanine carboxypeptidase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	179	1440	911	245	282	2	0	0	0
1	B	181	1462	926	248	286	2	0	1	0
1	C	178	1436	909	244	281	2	0	1	0
1	D	179	1440	911	245	282	2	0	0	0
1	E	180	1458	922	248	286	2	0	2	0
1	F	180	1460	925	248	285	2	0	2	0

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	175	ALA	GLU	engineered mutation	UNP Q47746
B	175	ALA	GLU	engineered mutation	UNP Q47746
C	175	ALA	GLU	engineered mutation	UNP Q47746
D	175	ALA	GLU	engineered mutation	UNP Q47746
E	175	ALA	GLU	engineered mutation	UNP Q47746
F	175	ALA	GLU	engineered mutation	UNP Q47746

- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Zn	0	0
			1	1		
2	B	1	Total	Zn	0	0
			1	1		
2	C	1	Total	Zn	0	0
			1	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	D	1	Total	Zn	0	0
			1	1		
2	E	1	Total	Zn	0	0
			1	1		
2	F	1	Total	Zn	0	0
			1	1		

- Molecule 3 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		
3	E	1	Total	O	S	0	0
			5	4	1		
3	F	1	Total	O	S	0	0
			5	4	1		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	88	Total	O	0	5
			93	93		
4	B	112	Total	O	0	1
			113	113		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	C	70	Total 70	O 70	0	0
4	D	83	Total 85	O 85	0	2
4	E	89	Total 89	O 89	0	0
4	F	89	Total 89	O 89	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: D-alanyl-D-alanine carboxypeptidase




- Molecule 1: D-alanyl-D-alanine carboxypeptidase




- Molecule 1: D-alanyl-D-alanine carboxypeptidase




- Molecule 1: D-alanyl-D-alanine carboxypeptidase




- Molecule 1: D-alanyl-D-alanine carboxypeptidase





- Molecule 1: D-alanyl-D-alanine carboxypeptidase



4 Data and refinement statistics i

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	55.71Å 77.32Å 85.86Å 101.34° 108.13° 90.06°	Depositor
Resolution (Å)	18.88 – 2.30 18.88 – 2.30	Depositor EDS
% Data completeness (in resolution range)	95.8 (18.88-2.30) 91.8 (18.88-2.30)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.70 (at 2.30Å)	Xtrriage
Refinement program	PHENIX 1.9_1692	Depositor
R, R_{free}	0.219 , 0.275 0.220 , 0.275	Depositor DCC
R_{free} test set	2894 reflections (5.04%)	wwPDB-VP
Wilson B-factor (Å ²)	30.7	Xtrriage
Anisotropy	0.033	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 42.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.29$	Xtrriage
Estimated twinning fraction	0.039 for h,-k,-h-l	Xtrriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	9261	wwPDB-VP
Average B, all atoms (Å ²)	49.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 55.08 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 3.3303e-05. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.24	0/1475	0.38	0/2008
1	B	0.21	0/1500	0.38	0/2041
1	C	0.24	0/1474	0.44	0/2006
1	D	0.23	0/1475	0.39	0/2008
1	E	0.27	0/1499	0.40	0/2040
1	F	0.21	0/1501	0.38	0/2043
All	All	0.24	0/8924	0.39	0/12146

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	C	0	1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	C	96	ARG	Mainchain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within

the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1440	0	1374	6	0
1	B	1462	0	1402	6	0
1	C	1436	0	1372	10	0
1	D	1440	0	1374	9	0
1	E	1458	0	1395	7	0
1	F	1460	0	1401	7	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
2	D	1	0	0	0	0
2	E	1	0	0	0	0
2	F	1	0	0	0	0
3	A	5	0	0	0	0
3	B	5	0	0	0	0
3	E	5	0	0	1	0
3	F	5	0	0	0	0
4	A	93	0	0	0	0
4	B	113	0	0	0	0
4	C	70	0	0	0	0
4	D	85	0	0	0	0
4	E	89	0	0	2	0
4	F	89	0	0	0	0
All	All	9261	0	8318	44	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (44) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:95:ASN:OD1	1:C:97:GLN:HG2	1.76	0.86
1:F:109:GLU:HB3	1:F:119:ILE:HG22	1.79	0.65
1:E:95:ASN:HD22	1:E:97:GLN:H	1.45	0.64
1:D:110:GLN:HG2	1:D:116:ARG:HG2	1.82	0.60
1:C:109:GLU:HB3	1:C:119:ILE:HG22	1.83	0.60
1:B:141:VAL:HG13	1:B:194[B]:VAL:HG13	1.84	0.59
1:E:105:ASP:N	1:E:105:ASP:OD1	2.35	0.58
1:B:184:GLY:HA2	1:B:189:GLN:HE22	1.69	0.57
1:D:93:LEU:HD11	1:D:261:GLU:HG2	1.85	0.57
1:F:226:PRO:HB2	1:F:229:LYS:HB2	1.86	0.57

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:146:ARG:NH1	1:B:151:GLN:OE1	2.39	0.54
1:C:97:GLN:HG3	1:C:98:ASN:N	2.22	0.54
1:D:223:ARG:NH2	1:D:237:ASN:OD1	2.41	0.54
1:C:97:GLN:CG	1:C:98:ASN:N	2.72	0.53
1:D:109:GLU:HB3	1:D:119:ILE:HG22	1.90	0.52
1:F:118:ASP:HB3	1:F:121[A]:ILE:HG12	1.91	0.52
1:D:150:LYS:HE3	1:D:154:ILE:HD11	1.91	0.51
1:D:223:ARG:HG3	1:D:242:TYR:CZ	2.44	0.51
1:D:258:LEU:HD21	1:D:263:TYR:HB2	1.93	0.51
1:E:173:LYS:NZ	4:E:405:HOH:O	2.41	0.51
1:E:109:GLU:HB3	1:E:119:ILE:HG22	1.92	0.50
3:E:302:SO4:O4	4:E:401:HOH:O	2.18	0.49
1:F:94:VAL:HG21	1:F:190:LEU:HD21	1.95	0.49
1:D:229:LYS:HG3	1:D:259:CYS:HB3	1.95	0.48
1:F:117:ILE:HB	1:F:121[B]:ILE:HD11	1.95	0.48
1:C:100:ILE:HG22	1:C:101:PRO:O	2.14	0.48
1:E:106:VAL:HG23	1:E:118:ASP:OD2	2.14	0.48
1:A:167:TYR:OH	1:F:107:GLU:OE1	2.21	0.47
1:A:187:GLU:OE2	1:A:241:HIS:NE2	2.42	0.46
1:A:109:GLU:HB3	1:A:119:ILE:HG22	1.98	0.46
1:C:117:ILE:HB	1:C:121:ILE:HD11	1.99	0.45
1:E:184:GLY:C	1:E:189[B]:GLN:HE22	2.20	0.45
1:F:121[A]:ILE:HD11	1:F:192:LEU:HD22	1.99	0.44
1:B:124:TYR:OH	1:C:127:ASP:OD2	2.30	0.44
1:E:186:SER:H	1:E:189[B]:GLN:NE2	2.16	0.44
1:C:93:LEU:HD21	1:C:97:GLN:HE21	1.83	0.44
1:A:146:ARG:NH2	1:A:195:ASP:OD2	2.50	0.43
1:A:100:ILE:HG22	1:A:184:GLY:O	2.19	0.43
1:B:186:SER:HB3	1:B:189:GLN:HG3	2.01	0.42
1:D:96:ARG:NH2	1:D:181:ALA:HA	2.34	0.42
1:C:93:LEU:HD21	1:C:97:GLN:NE2	2.34	0.42
1:A:132:ALA:HB1	1:A:137:VAL:HB	2.00	0.42
1:C:149:GLU:O	1:C:153:GLU:HG3	2.19	0.42
1:B:91:LEU:HB3	1:B:264:LEU:HD21	2.02	0.42

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	177/197 (90%)	171 (97%)	6 (3%)	0	100	100
1	B	180/197 (91%)	175 (97%)	5 (3%)	0	100	100
1	C	177/197 (90%)	165 (93%)	12 (7%)	0	100	100
1	D	177/197 (90%)	171 (97%)	6 (3%)	0	100	100
1	E	180/197 (91%)	174 (97%)	6 (3%)	0	100	100
1	F	180/197 (91%)	171 (95%)	9 (5%)	0	100	100
All	All	1071/1182 (91%)	1027 (96%)	44 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	150/165 (91%)	150 (100%)	0	100	100
1	B	153/165 (93%)	153 (100%)	0	100	100
1	C	150/165 (91%)	148 (99%)	2 (1%)	69	82
1	D	150/165 (91%)	150 (100%)	0	100	100
1	E	153/165 (93%)	150 (98%)	3 (2%)	55	72
1	F	153/165 (93%)	152 (99%)	1 (1%)	84	92
All	All	909/990 (92%)	903 (99%)	6 (1%)	86	92

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	98	ASN
1	C	107	GLU
1	E	104	TYR
1	E	189[A]	GLN
1	E	189[B]	GLN
1	F	87	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	C	97	GLN
1	E	95	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 10 ligands modelled in this entry, 6 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	SO4	B	302	2	4,4,4	0.14	0	6,6,6	0.05	0
3	SO4	F	302	-	4,4,4	0.14	0	6,6,6	0.05	0
3	SO4	A	302	2	4,4,4	0.14	0	6,6,6	0.05	0
3	SO4	E	302	2	4,4,4	0.14	0	6,6,6	0.07	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	E	302	SO4	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	179/197 (90%)	0.12	10 (5%) 24 30	25, 42, 72, 107	0
1	B	181/197 (91%)	0.08	6 (3%) 46 53	24, 37, 63, 130	0
1	C	178/197 (90%)	0.29	11 (6%) 20 26	32, 51, 88, 114	0
1	D	179/197 (90%)	0.66	15 (8%) 11 15	42, 60, 98, 141	0
1	E	180/197 (91%)	0.17	7 (3%) 39 46	28, 41, 76, 123	0
1	F	180/197 (91%)	0.20	11 (6%) 21 27	26, 43, 72, 121	0
All	All	1077/1182 (91%)	0.25	60 (5%) 24 30	24, 46, 83, 141	0

All (60) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	103	GLN	7.0
1	D	266	THR	7.0
1	F	266	THR	6.7
1	F	102	ALA	6.3
1	D	105	ASP	6.3
1	E	103	GLN	6.3
1	D	102	ALA	6.1
1	B	266	THR	5.9
1	E	87	GLN	5.4
1	E	102	ALA	5.3
1	B	102	ALA	5.2
1	C	257	GLY	5.2
1	C	103	GLN	5.1
1	A	257	GLY	5.1
1	C	97	GLN	5.0
1	D	200	GLY	4.8
1	A	103	GLN	4.4
1	C	102	ALA	4.1
1	D	208	VAL	4.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	D	204	THR	3.9
1	B	268	LYS	3.9
1	F	87	GLN	3.8
1	E	266	THR	3.6
1	D	265	ASN	3.5
1	D	257	GLY	3.3
1	D	103	GLN	3.0
1	C	200	GLY	3.0
1	A	210	ARG	3.0
1	C	201	ILE	3.0
1	E	228	ASP	3.0
1	D	201	ILE	2.9
1	D	135	ASP	2.7
1	D	206	ASN	2.6
1	F	103	GLN	2.6
1	A	102	ALA	2.6
1	A	266	THR	2.6
1	F	134	ALA	2.6
1	A	228	ASP	2.5
1	E	95	ASN	2.5
1	E	206	ASN	2.5
1	A	106	VAL	2.4
1	F	228	ASP	2.4
1	D	205	GLY	2.4
1	A	105	ASP	2.4
1	C	105	ASP	2.4
1	F	230	THR	2.3
1	F	201	ILE	2.3
1	C	104	TYR	2.2
1	A	255	HIS	2.2
1	B	228	ASP	2.2
1	C	202	HIS	2.2
1	C	258	LEU	2.2
1	F	255	HIS	2.1
1	A	230	THR	2.1
1	F	104	TYR	2.1
1	D	96	ARG	2.1
1	D	149	GLU	2.1
1	F	210	ARG	2.1
1	C	88	GLU	2.0
1	B	104	TYR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	SO4	F	302	5/5	0.80	0.18	109,110,111,114	0
3	SO4	B	302	5/5	0.86	0.25	97,97,99,99	0
3	SO4	A	302	5/5	0.87	0.20	92,99,102,103	0
3	SO4	E	302	5/5	0.88	0.20	98,100,102,106	0
2	ZN	D	500	1/1	0.96	0.04	48,48,48,48	0
2	ZN	F	301	1/1	0.96	0.08	37,37,37,37	0
2	ZN	B	301	1/1	0.96	0.15	48,48,48,48	0
2	ZN	E	301	1/1	0.97	0.06	30,30,30,30	1
2	ZN	C	500	1/1	0.97	0.10	64,64,64,64	0
2	ZN	A	301	1/1	1.00	0.08	32,32,32,32	1

6.5 Other polymers [i](#)

There are no such residues in this entry.