

wwPDB NMR Structure Validation Summary Report (i)

Aug 20, 2022 – 09:32 AM EDT

PDB ID : 2HSX

Title: NMR Structure of the nonstructural protein 1 (nsp1) from the SARS coron-

avirus

Authors: Almeida, M.S.; Herrmann, T.; Geralt, M.; Johnson, M.A.; Saikatendu, K.;

Joseph, J.; Subramanian, R.C.; Neuman, B.W.; Buchmeier, M.J.; Stevens, R.C.; Kuhn, P.; Wilson, I.A.; Wuthrich, K.; Joint Center for Structural Ge-

nomics (JCSG)

Deposited on : 2006-07-24

This is a wwPDB NMR Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/NMRValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity: 4.02b-467

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

RCI : v 1n 11 5 13 A (Berjanski et al., 2005)

PANAV : Wang et al. (2010)

ShiftChecker : 2.29

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)

Validation Pipeline (wwPDB-VP) : 2.29

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $SOLUTION\ NMR$

The overall completeness of chemical shifts assignment was not calculated.

There are no overall percentile quality scores available for this entry.

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Mol	Chain	Length	Quality of chain
1	A	116	100%



2 Ensemble composition and analysis (i)

This entry contains 1 models. Identification of well-defined residues and clustering analysis are not possible.



3 Entry composition (i)

There is only 1 type of molecule in this entry. The entry contains 1802 atoms, of which 911 are hydrogens and 0 are deuteriums.

• Molecule 1 is a protein called Leader protein; p65 homolog; NSP1 (EC 3.4.22.-).

\mathbf{M}	ol	Chain	Residues			Aton	ns			Trace
1		Λ	116	Total	С	Н	N	О	S	0
1	-	A	110	1802	562	911	161	166	2	U

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference			
Α	1	GLY	THR	engineered mutation	UNP P59641			



4 Residue-property plots (i)

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

• Molecule 1: Leader protein; p65 homolog; NSP1 (EC 3.4.22.-)

Ch	ai	n	A	۲:																						10	0%	ó																				l			
G1 H2	V3	1.5	36	L7	P8	7 1	110	V12	R13	D14	V15	L16	V17	R18	G19	F20	621	770	V24	E25	E26	A27	L28	S29	E30	A31	R32	E33	H34	K36	N37	388	T39	C40	G41	L42 V43	F.44	1.45	E46	K47	G48	V49	L50	P51	1.53	E54	990	P56	Y57	F59	160
K61 R62	863	D64 A65	799T	298	T68	N69	271	471 H72	K73	V74	V75	E76	177	N78	A 79	E80	MOI	787	20 7	082	786 V	C87	R88	888	065	191	T92	L93	G94 Vor	96.1	767	P98	99	9	9	2 5	9	9	9	10	9	9:	i :	: :	1 =	K114	큐	H			



Refinement protocol and experimental data overview (i) 5



The models were refined using the following method: ATNOS/CANDID/CYANA.

Of the 120 calculated structures, 1 were deposited, based on the following criterion: Closest to the mean structure.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version				
CYANA	structure solution	1.0.3				
CYANA	refinement	1.0.3				

No chemical shift data was provided.



6 Model quality (i)

6.1 Standard geometry (i)

There are no covalent bond-length or bond-angle outliers.

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

There are no planarity outliers.

6.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
All	All	0	0	0	-

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is -.

There are no clashes.

6.3 Torsion angles (i)

6.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	0	-	_	_	-
All	All	0	-	-	-	-

There are no Ramachandran outliers.



6.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	0	-	-	-
All	All	0	-	-	-

There are no protein residues with a non-rotameric sidechain to report.

6.3.3 RNA (i)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates (i)

There are no monosaccharides in this entry.

6.6 Ligand geometry (i)

There are no ligands in this entry.

6.7 Other polymers (i)

There are no such molecules in this entry.

6.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



7 Chemical shift validation (i)

No chemical shift data were provided

