



# Full wwPDB X-ray Structure Validation Report ⓘ

May 25, 2020 – 03:43 am BST

PDB ID : 6HUH  
Title : CRYSTAL STRUCTURE OF OXA-427 class D BETA-LACTAMASE  
Authors : Zavala, A.; Retailleau, P.; Bogaerts, P.; Glupczynski, Y.; Naas, T.; Iorga, B.  
Deposited on : 2018-10-08  
Resolution : 2.78 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.78 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	4107 (2.80-2.76)
Clashscore	141614	4575 (2.80-2.76)
Ramachandran outliers	138981	4487 (2.80-2.76)
Sidechain outliers	138945	4489 (2.80-2.76)
RSRZ outliers	127900	4027 (2.80-2.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	274	 3% 71% 16% • 12%
1	B	274	 11% 73% 14% • 12%

## 2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 3971 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Beta-lactamase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	242	1912	1228	328	351	5	0	1	0
1	B	242	1912	1228	328	351	5	0	1	0

There are 20 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	265	VAL	-	expression tag	UNP A0A1Y6M6D7
A	266	GLU	-	expression tag	UNP A0A1Y6M6D7
A	267	HIS	-	expression tag	UNP A0A1Y6M6D7
A	268	HIS	-	expression tag	UNP A0A1Y6M6D7
A	269	HIS	-	expression tag	UNP A0A1Y6M6D7
A	270	HIS	-	expression tag	UNP A0A1Y6M6D7
A	271	HIS	-	expression tag	UNP A0A1Y6M6D7
A	272	HIS	-	expression tag	UNP A0A1Y6M6D7
A	273	HIS	-	expression tag	UNP A0A1Y6M6D7
A	274	HIS	-	expression tag	UNP A0A1Y6M6D7
B	265	VAL	-	expression tag	UNP A0A1Y6M6D7
B	266	GLU	-	expression tag	UNP A0A1Y6M6D7
B	267	HIS	-	expression tag	UNP A0A1Y6M6D7
B	268	HIS	-	expression tag	UNP A0A1Y6M6D7
B	269	HIS	-	expression tag	UNP A0A1Y6M6D7
B	270	HIS	-	expression tag	UNP A0A1Y6M6D7
B	271	HIS	-	expression tag	UNP A0A1Y6M6D7
B	272	HIS	-	expression tag	UNP A0A1Y6M6D7
B	273	HIS	-	expression tag	UNP A0A1Y6M6D7
B	274	HIS	-	expression tag	UNP A0A1Y6M6D7

- Molecule 2 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			3	2	1		
2	B	1	Total	O	S	0	0
			5	4	1		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	69	Total	O	0	0
			69	69		
3	B	65	Total	O	0	0
			65	65		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Beta-lactamase



- Molecule 1: Beta-lactamase



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	143.79Å 42.58Å 99.71Å 90.00° 114.37° 90.00°	Depositor
Resolution (Å)	68.10 – 2.78 68.10 – 2.78	Depositor EDS
% Data completeness (in resolution range)	94.2 (68.10-2.78) 94.2 (68.10-2.78)	Depositor EDS
$R_{merge}$	0.16	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.26 (at 2.77Å)	Xtrriage
Refinement program	BUSTER 2.10.3	Depositor
R, $R_{free}$	0.205 , 0.269 0.213 , 0.279	Depositor DCC
$R_{free}$ test set	681 reflections (5.09%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	41.4	Xtrriage
Anisotropy	0.433	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 73.3	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.46$ , $\langle L^2 \rangle = 0.29$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	3971	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	50.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 19.88 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 9.8380e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, KCX

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.51	0/1948	0.72	0/2644
1	B	0.50	0/1948	0.73	0/2644
All	All	0.50	0/3896	0.72	0/5288

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1912	0	1881	16	0
1	B	1912	0	1881	17	0
2	A	8	0	0	0	0
2	B	5	0	0	0	0
3	A	69	0	0	0	0
3	B	65	0	0	0	0
All	All	3971	0	3762	32	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (32) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:185:ARG:HH22	1:B:193:GLN:HG2	1.43	0.81
1:B:205:LEU:HB3	1:B:210:LEU:HB3	1.77	0.65
1:B:220:VAL:HG12	1:B:233:VAL:HG13	1.82	0.61
1:A:124:ARG:HD3	1:A:171:PRO:HG3	1.84	0.59
1:A:220:VAL:HG12	1:A:233:VAL:HG13	1.84	0.59
1:A:198:THR:HA	1:A:218:TRP:O	2.04	0.58
1:B:198:THR:HB	1:B:246:SER:HB2	1.86	0.58
1:B:124:ARG:HD3	1:B:171:PRO:HG3	1.85	0.57
1:A:58:ALA:HB2	1:A:98:MET:HE2	1.87	0.56
1:A:59:LEU:HD11	1:A:165:LEU:HD22	1.87	0.56
1:B:198:THR:HA	1:B:218:TRP:O	2.06	0.55
1:B:59:LEU:HD11	1:B:165:LEU:HD22	1.89	0.53
1:A:203:PRO:HG3	1:A:216:ILE:HD11	1.94	0.48
1:A:253:VAL:O	1:A:257:LEU:HB2	2.16	0.46
1:B:207:ASP:HB3	1:B:210:LEU:HG	1.97	0.45
1:A:198:THR:HB	1:A:246:SER:HB2	1.96	0.45
1:A:70:GLU:HG3	1:A:183:ILE:HG23	1.98	0.45
1:A:217:GLY:O	1:A:235:THR:HA	2.16	0.44
1:B:205:LEU:HD13	1:B:212:ARG:HA	2.00	0.43
1:B:70:GLU:HG3	1:B:183:ILE:HG23	2.00	0.43
1:A:59:LEU:HD22	1:A:172:VAL:HG21	2.01	0.43
1:A:62:TYR:HB2	1:A:176:THR:HG23	2.01	0.43
1:B:62:TYR:HB2	1:B:176:THR:HG23	2.01	0.42
1:B:27:MET:HG2	1:B:234:HIS:CE1	2.54	0.42
1:B:210:LEU:HD13	1:B:211:ASP:H	1.85	0.41
1:B:30:ALA:HB3	1:B:231:ILE:HB	2.02	0.41
1:B:192:TRP:CH2	1:B:264:LEU:HD23	2.56	0.41
1:A:243:GLN:HB2	1:A:248:ARG:NH2	2.35	0.41
1:A:187:PRO:HA	1:B:71:GLN:OE1	2.20	0.41
1:A:54[B]:KCX:HZ	1:A:103:ILE:HA	1.86	0.41
1:B:192:TRP:HH2	1:B:264:LEU:HD23	1.86	0.41
1:A:58:ALA:O	1:A:62:TYR:HD2	2.05	0.40

There are no symmetry-related clashes.



## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	239/274 (87%)	219 (92%)	17 (7%)	3 (1%)	12	33
1	B	239/274 (87%)	221 (92%)	17 (7%)	1 (0%)	34	64
All	All	478/548 (87%)	440 (92%)	34 (7%)	4 (1%)	19	47

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	204	LYS
1	B	208	GLY
1	A	205	LEU
1	A	208	GLY

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	200/226 (88%)	181 (90%)	19 (10%)	8	23
1	B	200/226 (88%)	183 (92%)	17 (8%)	10	28
All	All	400/452 (88%)	364 (91%)	36 (9%)	9	26

All (36) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	26	CYS
1	A	87	GLU

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	90	GLN
1	A	117	ARG
1	A	144	GLN
1	A	147	LEU
1	A	154	SER
1	A	162	LEU
1	A	165	LEU
1	A	166	VAL
1	A	205	LEU
1	A	206	LEU
1	A	210	LEU
1	A	215	GLN
1	A	227	ASP
1	A	230	LEU
1	A	238	GLN
1	A	239	LYS
1	A	262	LYS
1	B	26	CYS
1	B	27	MET
1	B	90	GLN
1	B	117	ARG
1	B	144	GLN
1	B	147	LEU
1	B	154	SER
1	B	162	LEU
1	B	165	LEU
1	B	166	VAL
1	B	204	LYS
1	B	206	LEU
1	B	210	LEU
1	B	212	ARG
1	B	227	ASP
1	B	230	LEU
1	B	248	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
1	KCX	B	54[B]	-	7,8,12	0.75	0	4,8,14	7.71	1 (25%)
1	KCX	A	54[B]	-	7,8,12	0.78	0	4,8,14	1.62	1 (25%)
1	KCX	A	54[A]	-	7,11,12	1.03	1 (14%)	4,12,14	0.96	0
1	KCX	B	54[A]	-	7,11,12	1.05	1 (14%)	4,12,14	1.09	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	KCX	B	54[B]	-	-	1/7/7/12	-
1	KCX	A	54[B]	-	-	5/7/7/12	-
1	KCX	A	54[A]	-	-	3/7/10/12	-
1	KCX	B	54[A]	-	-	1/7/10/12	-

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	54[A]	KCX	CE-NZ	2.18	1.50	1.45
1	A	54[A]	KCX	CE-NZ	2.05	1.50	1.45

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	54[B]	KCX	CE-NZ-CX	15.32	148.89	122.95
1	A	54[B]	KCX	CE-NZ-CX	2.98	127.99	122.95

There are no chirality outliers.

All (10) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	B	54[B]	KCX	CD-CE-NZ-CX
1	A	54[B]	KCX	C-CA-CB-CG
1	A	54[B]	KCX	CD-CE-NZ-CX
1	A	54[A]	KCX	C-CA-CB-CG
1	A	54[B]	KCX	CG-CD-CE-NZ
1	A	54[B]	KCX	CA-CB-CG-CD
1	A	54[B]	KCX	N-CA-CB-CG
1	A	54[A]	KCX	CE-CD-CG-CB
1	B	54[A]	KCX	C-CA-CB-CG
1	A	54[A]	KCX	N-CA-CB-CG

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	54[B]	KCX	1	0

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	SO4	B	301	-	4,4,4	0.23	0	6,6,6	0.07	0
2	SO4	A	302	-	0,2,4	0.00	-	0,1,6	0.00	-
2	SO4	A	301	-	4,4,4	0.20	0	6,6,6	0.13	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	241/274 (87%)	0.32	9 (3%) 41 36	23, 42, 84, 127	0
1	B	241/274 (87%)	0.71	29 (12%) 4 3	27, 49, 92, 130	0
All	All	482/548 (87%)	0.51	38 (7%) 12 9	23, 45, 92, 130	0

All (38) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	206	LEU	11.6
1	A	210	LEU	10.0
1	B	208	GLY	9.5
1	A	208	GLY	8.4
1	B	207	ASP	6.1
1	A	211	ASP	5.4
1	B	209	SER	5.0
1	B	82	PRO	4.7
1	B	212	ARG	4.6
1	B	241	GLY	4.4
1	B	210	LEU	4.4
1	B	213	ASP	4.1
1	B	205	LEU	3.6
1	B	256	ALA	3.6
1	B	214	GLN	3.5
1	B	237	ILE	3.5
1	B	264	LEU	3.4
1	B	227	ASP	3.0
1	B	211	ASP	2.9
1	B	47	LEU	2.9
1	B	240	PRO	2.8
1	B	84	PHE	2.8
1	A	209	SER	2.7
1	A	213	ASP	2.7

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	148	SER	2.5
1	B	137	GLY	2.5
1	B	192	TRP	2.5
1	B	184	LEU	2.4
1	B	126	ASP	2.4
1	B	206	LEU	2.3
1	A	200	THR	2.3
1	B	145	ALA	2.3
1	B	189	ILE	2.3
1	B	136	PRO	2.2
1	B	238	GLN	2.2
1	B	239	LYS	2.1
1	A	227	ASP	2.1
1	B	147	LEU	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
1	KCX	B	54[B]	9/13	0.96	0.21	28,28,31,34	5
1	KCX	B	54[A]	12/13	0.96	0.21	21,26,31,34	8
1	KCX	A	54[A]	12/13	0.98	0.17	15,23,28,29	8
1	KCX	A	54[B]	9/13	0.98	0.17	26,26,28,29	5

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	SO4	B	301	5/5	0.97	0.15	87,88,90,90	0
2	SO4	A	302	3/5	0.98	0.44	45,45,46,46	3
2	SO4	A	301	5/5	0.98	0.16	88,88,88,90	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.