



# Full wwPDB X-ray Structure Validation Report ⓘ

Nov 1, 2023 – 01:13 AM EDT

PDB ID : 3HW1  
Title : Structure of Bace (beta secretase) in complex with ligand EV2  
Authors : Godemann, R.; Madden, J.; Kramer, J.; Smith, M.A.; Barker, J.; Ebnet, A.  
Deposited on : 2009-06-17  
Resolution : 2.48 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.36  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.48 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	5857 (2.50-2.46)
Clashscore	141614	6594 (2.50-2.46)
Ramachandran outliers	138981	6469 (2.50-2.46)
Sidechain outliers	138945	6471 (2.50-2.46)
RSRZ outliers	127900	5738 (2.50-2.46)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	411	<div style="display: flex; align-items: center;"> <div style="width: 5%; height: 10px; background-color: red; margin-right: 5px;"></div> <div style="width: 79%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 10%; height: 10px; background-color: yellow; margin-right: 5px;"></div> <div style="width: 10%; height: 10px; background-color: grey;"></div> </div> <p style="margin-left: 20px;">5%      79%      10%      •      10%</p>
1	B	411	<div style="display: flex; align-items: center;"> <div style="width: 8%; height: 10px; background-color: red; margin-right: 5px;"></div> <div style="width: 81%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 9%; height: 10px; background-color: yellow; margin-right: 5px;"></div> <div style="width: 10%; height: 10px; background-color: grey;"></div> </div> <p style="margin-left: 20px;">8%      81%      9%      10%</p>
1	C	411	<div style="display: flex; align-items: center;"> <div style="width: 7%; height: 10px; background-color: red; margin-right: 5px;"></div> <div style="width: 84%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 7%; height: 10px; background-color: yellow; margin-right: 5px;"></div> <div style="width: 9%; height: 10px; background-color: grey;"></div> </div> <p style="margin-left: 20px;">7%      84%      7%      9%</p>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	EV2	B	393[A]	-	-	-	X
2	EV2	B	393[B]	-	-	-	X
3	GOL	C	394	-	-	-	X

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 8985 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Beta-secretase 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	370	2921	1871	486	550	14	10	1	0
1	B	371	2920	1869	486	551	14	21	0	0
1	C	375	2957	1893	494	556	14	4	1	0

There are 15 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-18	LEU	-	expression tag	UNP P56817
A	-17	PRO	-	expression tag	UNP P56817
A	-16	ARG	-	expression tag	UNP P56817
A	-5	LYS	ARG	engineered mutation	UNP P56817
A	-4	THR	ARG	engineered mutation	UNP P56817
B	-18	LEU	-	expression tag	UNP P56817
B	-17	PRO	-	expression tag	UNP P56817
B	-16	ARG	-	expression tag	UNP P56817
B	-5	LYS	ARG	engineered mutation	UNP P56817
B	-4	THR	ARG	engineered mutation	UNP P56817
C	-18	LEU	-	expression tag	UNP P56817
C	-17	PRO	-	expression tag	UNP P56817
C	-16	ARG	-	expression tag	UNP P56817
C	-5	LYS	ARG	engineered mutation	UNP P56817
C	-4	THR	ARG	engineered mutation	UNP P56817

- Molecule 2 is 3-pyrrolidin-1-ylquinoxalin-2-amine (three-letter code: EV2) (formula:  $C_{12}H_{14}N_4$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	N	0	0
			16	12	4		
2	B	1	Total	C	N	0	1
			32	24	8		
2	C	1	Total	C	N	0	0
			16	12	4		

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	C	1	Total	C	O	0	0
			6	3	3		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	40	Total O 40 40	0	0
4	B	31	Total O 31 31	0	0
4	C	46	Total O 46 46	0	0


### 3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Beta-secretase 1




- Molecule 1: Beta-secretase 1



- Molecule 1: Beta-secretase 1







## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	230.62Å 99.88Å 62.90Å 90.00° 103.12° 90.00°	Depositor
Resolution (Å)	48.80 – 2.48 48.82 – 2.48	Depositor EDS
% Data completeness (in resolution range)	100.0 (48.80-2.48) 98.4 (48.82-2.48)	Depositor EDS
$R_{merge}$	0.05	Depositor
$R_{sym}$	0.05	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.64 (at 2.48Å)	Xtrriage
Refinement program	REFMAC 5.2.0005	Depositor
R, $R_{free}$	0.226 , 0.281 0.227 , 0.279	Depositor DCC
$R_{free}$ test set	986 reflections (2.03%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	61.1	Xtrriage
Anisotropy	0.713	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.30 , 55.1	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.031 for -h-2*1,-k,l	Xtrriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	8985	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	21.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.32% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, EV2

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.54	0/2995	0.65	1/4067 (0.0%)
1	B	0.46	0/2992	0.59	0/4062
1	C	0.53	0/3034	0.63	0/4120
All	All	0.51	0/9021	0.63	1/12249 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	0	VAL	O-C-N	-5.50	113.90	122.70

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2921	0	2832	22	0
1	B	2920	0	2833	13	0
1	C	2957	0	2875	14	0
2	A	16	0	14	0	0
2	B	32	0	28	0	0
2	C	16	0	14	0	0
3	C	6	0	8	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	A	40	0	0	0	0
4	B	31	0	0	0	0
4	C	46	0	0	0	0
All	All	8985	0	8604	47	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (47) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:194:ARG:HH22	1:A:385:ASN:HD21	1.35	0.72
1:A:181:HIS:HA	1:A:184:TYR:CE1	2.34	0.62
1:A:282:VAL:HG13	1:A:299:THR:HG23	1.85	0.59
1:A:194:ARG:NH2	1:A:385:ASN:HD21	2.01	0.58
1:C:278:ASN:H	1:C:278:ASN:HD22	1.55	0.55
1:B:199:TYR:HB3	1:B:352:ILE:HD11	1.90	0.54
1:B:261:PHE:CE1	1:B:322:PHE:HB2	2.43	0.53
1:A:261:PHE:CD1	1:A:268:VAL:HG23	2.43	0.53
1:C:42:ALA:CB	1:C:101:ALA:HB1	2.40	0.51
1:C:48:LEU:HD21	1:C:109:PHE:CD2	2.46	0.51
1:C:69:VAL:HG21	1:C:76:TRP:CZ2	2.47	0.50
1:A:199:TYR:HB3	1:A:352:ILE:HD11	1.94	0.49
1:A:181:HIS:HA	1:A:184:TYR:HE1	1.78	0.48
1:A:143:GLN:HG3	1:C:1:GLU:OE1	2.13	0.48
1:A:276:PRO:O	1:A:279:ILE:HG12	2.14	0.48
1:A:357:SER:HB3	1:A:360:HIS:HB3	1.95	0.47
1:B:65:LYS:HB3	1:B:80:LEU:HD12	1.95	0.47
1:A:362:HIS:HB2	1:A:366:ARG:O	2.15	0.46
1:C:123:TYR:CD2	1:C:196:GLU:HB3	2.51	0.46
1:A:222:TYR:HA	1:A:223:ASP:HA	1.75	0.46
1:A:140:LEU:HD21	1:A:176:ILE:HG21	1.97	0.45
1:A:248:ILE:HD11	1:A:283:ILE:HD11	1.98	0.45
1:C:222:TYR:CD1	1:C:223:ASP:HA	2.51	0.45
1:A:121:LEU:O	1:A:199:TYR:OH	2.27	0.45
1:B:14:TYR:CG	1:B:154:LEU:HD22	2.52	0.45
1:B:25:GLN:HE22	1:B:53:GLN:HB2	1.82	0.43
1:B:357:SER:HB3	1:B:360:HIS:HB3	1.99	0.43
1:B:222:TYR:HA	1:B:223:ASP:HA	1.75	0.43
1:B:261:PHE:CD1	1:B:268:VAL:HG23	2.54	0.42
1:C:22:SER:O	1:C:57:SER:HA	2.19	0.42

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:217:CYS:HA	1:A:220:TYR:CD1	2.55	0.42
1:B:142:LYS:NZ	1:B:142:LYS:HB3	2.35	0.42
1:B:235:ARG:HB3	1:B:327:SER:HB2	2.01	0.42
1:C:199:TYR:HB3	1:C:352:ILE:HD11	2.02	0.42
1:C:235:ARG:HB3	1:C:327:SER:HB2	2.01	0.42
1:B:238:LYS:NZ	1:C:385:ASN:HB2	2.36	0.41
1:B:298:ILE:HB	1:B:341:PHE:CZ	2.55	0.41
1:A:20:VAL:HG12	1:A:85:VAL:HG22	2.03	0.41
1:C:234:LEU:HB2	1:C:336:VAL:CG2	2.50	0.41
1:A:130:ASP:OD2	1:A:132:SER:OG	2.22	0.41
1:C:385:ASN:HD22	1:C:385:ASN:H	1.68	0.41
1:A:69:VAL:HG21	1:A:76:TRP:CZ2	2.56	0.41
1:A:362:HIS:HD2	1:A:363:ASP:O	2.03	0.41
1:B:69:VAL:HB	1:B:128:ARG:HG3	2.02	0.41
1:A:237:PRO:O	1:A:238:LYS:C	2.59	0.41
1:A:188:LEU:HD11	1:A:344:VAL:HG21	2.03	0.41
1:C:384:TYR:O	1:C:385:ASN:C	2.60	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	365/411 (89%)	345 (94%)	18 (5%)	2 (0%)	29	46
1	B	363/411 (88%)	344 (95%)	18 (5%)	1 (0%)	41	59
1	C	370/411 (90%)	355 (96%)	14 (4%)	1 (0%)	41	59
All	All	1098/1233 (89%)	1044 (95%)	50 (5%)	4 (0%)	34	52

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	238	LYS
1	B	378	ASP
1	A	156	GLY
1	C	156	GLY

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	316/352 (90%)	308 (98%)	8 (2%)	47	71
1	B	317/352 (90%)	304 (96%)	13 (4%)	30	53
1	C	320/352 (91%)	311 (97%)	9 (3%)	43	67
All	All	953/1056 (90%)	923 (97%)	30 (3%)	39	64

All (30) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	125	GLU
1	A	197	TRP
1	A	213	LEU
1	A	239	LYS
1	A	282	VAL
1	A	336	VAL
1	A	359	CYS
1	A	363	ASP
1	B	35	SER
1	B	54	ARG
1	B	63	LEU
1	B	86	SER
1	B	131	ASP
1	B	138	ASP
1	B	145	HIS
1	B	149	LEU
1	B	169	SER
1	B	197	TRP
1	B	292	THR

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	311	ASP
1	B	317	ASP
1	C	0	VAL
1	C	138	ASP
1	C	173	SER
1	C	197	TRP
1	C	278	ASN
1	C	282	VAL
1	C	317	ASP
1	C	367	THR
1	C	376	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	28	ASN
1	A	266	GLN
1	A	278	ASN
1	A	362	HIS
1	A	385	ASN
1	C	28	ASN
1	C	181	HIS
1	C	278	ASN
1	C	362	HIS
1	C	385	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry

5 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	EV2	A	393	-	18,18,18	2.18	4 (22%)	22,25,25	1.54	4 (18%)
2	EV2	B	393[B]	-	18,18,18	2.30	4 (22%)	22,25,25	1.54	4 (18%)
2	EV2	B	393[A]	-	18,18,18	2.21	3 (16%)	22,25,25	1.51	5 (22%)
3	GOL	C	394	-	5,5,5	0.45	0	5,5,5	0.60	0
2	EV2	C	393	-	18,18,18	2.18	4 (22%)	22,25,25	1.35	3 (13%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	EV2	A	393	-	-	4/4/11/11	0/3/3/3
2	EV2	B	393[B]	-	-	3/4/11/11	0/3/3/3
2	EV2	B	393[A]	-	-	4/4/11/11	0/3/3/3
3	GOL	C	394	-	-	4/4/4/4	-
2	EV2	C	393	-	-	4/4/11/11	0/3/3/3

All (15) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	393[B]	EV2	C9-N7	7.75	1.40	1.31
2	B	393[A]	EV2	C9-N7	7.49	1.40	1.31
2	C	393	EV2	C9-N7	7.37	1.40	1.31
2	A	393	EV2	C9-N7	7.36	1.39	1.31
2	B	393[B]	EV2	C10-N12	3.01	1.41	1.34
2	B	393[A]	EV2	C10-N12	2.96	1.41	1.34
2	C	393	EV2	C10-N12	2.78	1.41	1.34
2	A	393	EV2	C10-C9	-2.53	1.40	1.44

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	393[B]	EV2	C9-N11	2.46	1.43	1.37
2	A	393	EV2	C10-N12	2.40	1.40	1.34
2	B	393[A]	EV2	C9-N11	2.21	1.43	1.37
2	C	393	EV2	C10-C9	-2.15	1.40	1.44
2	A	393	EV2	C9-N11	2.12	1.43	1.37
2	C	393	EV2	C9-N11	2.08	1.42	1.37
2	B	393[B]	EV2	C10-C9	-2.03	1.41	1.44

All (16) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	393	EV2	C14-C16-N11	3.78	108.25	103.45
2	A	393	EV2	C15-N11-C16	-3.58	104.95	111.46
2	B	393[B]	EV2	C14-C16-N11	3.57	107.98	103.45
2	C	393	EV2	C15-N11-C16	-3.35	105.37	111.46
2	B	393[A]	EV2	C14-C16-N11	3.23	107.55	103.45
2	C	393	EV2	C13-C15-N11	3.14	107.44	103.45
2	B	393[B]	EV2	C15-N11-C16	-3.10	105.82	111.46
2	B	393[A]	EV2	C15-N11-C16	-3.10	105.83	111.46
2	B	393[B]	EV2	C13-C15-N11	2.93	107.17	103.45
2	B	393[A]	EV2	C10-C9-N11	2.67	124.83	121.69
2	A	393	EV2	N7-C9-N11	2.63	119.50	117.62
2	B	393[B]	EV2	C9-N7-C5	2.55	121.50	116.11
2	B	393[A]	EV2	C13-C15-N11	2.46	106.58	103.45
2	B	393[A]	EV2	C9-N7-C5	2.40	121.18	116.11
2	C	393	EV2	C9-N7-C5	2.27	120.91	116.11
2	A	393	EV2	C9-N7-C5	2.06	120.45	116.11

There are no chirality outliers.

All (19) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	393	EV2	C10-C9-N11-C16
2	A	393	EV2	N7-C9-N11-C16
2	B	393[A]	EV2	C10-C9-N11-C16
2	B	393[A]	EV2	C10-C9-N11-C15
2	B	393[A]	EV2	N7-C9-N11-C16
2	B	393[A]	EV2	N7-C9-N11-C15
2	B	393[B]	EV2	C10-C9-N11-C15
2	B	393[B]	EV2	N7-C9-N11-C15
2	C	393	EV2	C10-C9-N11-C16
2	C	393	EV2	C10-C9-N11-C15

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	Atoms
2	C	393	EV2	N7-C9-N11-C16
2	C	393	EV2	N7-C9-N11-C15
3	C	394	GOL	O1-C1-C2-O2
3	C	394	GOL	O1-C1-C2-C3
3	C	394	GOL	O2-C2-C3-O3
3	C	394	GOL	C1-C2-C3-O3
2	A	393	EV2	C10-C9-N11-C15
2	A	393	EV2	N7-C9-N11-C15
2	B	393[B]	EV2	C10-C9-N11-C16

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	370/411 (90%)	0.56	21 (5%) 23 24	4, 21, 27, 36	7 (1%)
1	B	371/411 (90%)	0.70	32 (8%) 10 10	9, 21, 30, 39	10 (2%)
1	C	375/411 (91%)	0.74	30 (8%) 12 11	11, 23, 30, 38	6 (1%)
All	All	1116/1233 (90%)	0.67	83 (7%) 14 14	4, 21, 30, 39	23 (2%)

All (83) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	72	THR	4.4
1	A	71	TYR	4.3
1	C	376	THR	4.0
1	C	379	MET	3.9
1	C	244	ALA	3.7
1	A	380	GLU	3.7
1	A	72	THR	3.6
1	A	84	LEU	3.5
1	B	-4	THR	3.4
1	B	145	HIS	3.4
1	A	68	TYR	3.4
1	B	385	ASN	3.3
1	A	145	HIS	3.2
1	A	92	ASN	3.2
1	A	86[A]	SER	3.2
1	C	168	ALA	3.1
1	B	377	LEU	3.1
1	A	110	ILE	3.1
1	B	47	PHE	3.0
1	C	257	PHE	3.0
1	C	384	TYR	3.0
1	C	74	GLY	3.0
1	B	157	ALA	3.0

*Continued on next page...*

*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Res</b>	<b>Type</b>	<b>RSRZ</b>
1	A	52	TYR	2.9
1	B	379	MET	2.9
1	B	324	ILE	2.9
1	C	71	TYR	2.9
1	B	263	LEU	2.8
1	C	262	TRP	2.8
1	C	68	TYR	2.8
1	B	108	PHE	2.8
1	B	333	MET	2.8
1	C	292	THR	2.7
1	B	265	GLU	2.7
1	B	41	GLY	2.7
1	B	76	TRP	2.7
1	A	117	GLY	2.6
1	B	70	PRO	2.6
1	C	76	TRP	2.6
1	C	220	TYR	2.6
1	B	261	PHE	2.6
1	B	262	TRP	2.6
1	C	256	LYS	2.6
1	C	380	GLU	2.6
1	C	286	TYR	2.5
1	C	67	VAL	2.5
1	B	305	TYR	2.5
1	A	24	PRO	2.5
1	B	110	ILE	2.5
1	A	262	TRP	2.5
1	A	50	ARG	2.5
1	C	377	LEU	2.4
1	B	227	VAL	2.4
1	B	337	ILE	2.4
1	B	234	LEU	2.4
1	C	242	GLU	2.4
1	B	378	ASP	2.3
1	B	69	VAL	2.3
1	B	214	LYS	2.3
1	B	376	THR	2.3
1	C	31	VAL	2.2
1	B	104	GLU	2.2
1	C	75	LYS	2.2
1	B	79	GLU	2.2
1	C	215	MET	2.1

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	C	227	VAL	2.1
1	B	66	GLY	2.1
1	C	254	THR	2.1
1	A	91	PRO	2.1
1	B	119	LEU	2.1
1	C	240	VAL	2.1
1	C	118	ILE	2.1
1	C	271	GLN	2.1
1	A	108	PHE	2.1
1	B	113	SER	2.1
1	B	128	ARG	2.1
1	A	171	GLY	2.1
1	A	49	HIS	2.1
1	A	118	ILE	2.1
1	C	78	GLY	2.1
1	A	79	GLU	2.0
1	A	48	LEU	2.0
1	C	334	GLY	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	EV2	B	393[A]	16/16	0.61	0.42	54,54,55,56	16
2	EV2	B	393[B]	16/16	0.61	0.42	41,42,42,42	16
3	GOL	C	394	6/6	0.66	0.62	56,59,60,61	0
2	EV2	A	393	16/16	0.79	0.25	36,37,37,37	16
2	EV2	C	393	16/16	0.89	0.32	38,39,40,40	16

## 6.5 Other polymers

There are no such residues in this entry.