

wwPDB X-ray Structure Validation Summary Report (i)

May 15, 2020 – 01:47 pm BST

PDB ID : 6HWN

Title: Structure of Thermus thermophilus ClpP in complex with a tripeptide.

Authors : Felix, J.; Schanda, P.; Fraga, H.; Morlot, C.

Deposited on : 2018-10-12

Resolution : 1.95 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity : 4.02b-467

Mogul : 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : 1.13 EDS : 2.11

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

Refmac: 5.8.0158

 $\begin{array}{cccc} & CCP4 & : & 7.0.044 \; (Gargrove) \\ Ideal \; geometry \; (proteins) & : & Engh \; \& \; Huber \; (2001) \end{array}$

Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.11

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X- $RAY\ DIFFRACTION$

The reported resolution of this entry is 1.95 Å.


Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$\begin{array}{c} \text{Whole archive} \\ (\#\text{Entries}) \end{array}$	$\begin{array}{c} {\rm Similar \; resolution} \\ (\#{\rm Entries, \; resolution \; range(\AA)}) \end{array}$
R_{free}	130704	2580 (1.96-1.96)
Clashscore	141614	2705 (1.96-1.96)
Ramachandran outliers	138981	2678 (1.96-1.96)
Sidechain outliers	138945	2678 (1.96-1.96)
RSRZ outliers	127900	2539 (1.96-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain		
1	A	204	79% 139	06	8%
			5%	70	070
1	В	204	82% 89 4%	%	9%
1	С	204		9%	8%
1	D	204	82%	0%	8%
1	Е	204	5% 84%	7%	9%
1	F	204	83%	3%	8%





Continued from previous page...

Mol	Chain	Length	Quality of chain	
1	G	204	83%	7% 10%
2	Н	3	67%	33%
2	I	3	67%	33%
2	J	3	67%	33%
2	K	3	100%	
2	L	3	67%	33%
2	M	3	67%	33%
2	N	3	67%	33%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	PEG	D	302	-	-	X	-
3	PEG	E	302	-	-	X	-



2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 11036 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called ATP-dependent Clp protease proteolytic subunit.

Mol	Chain	Residues		At	oms			ZeroOcc	AltConf	Trace
1	A	187	Total	С	N	О	S	0	3	0
1	A	107	1448 928 246 269 5	J 3						
1	В	186	Total	С	N	О	S	0	2	0
1	Б	100	1437	921	242	270	4	0	2	0
1	С	188	Total	С	N	О	S	0	2	0
1		100	1438	923	242	268	5	0	Δ	U
1	D	187	Total	С	N O S	0	5	0		
1	D	107	1473	947	249	272	5	U	J J	U
1	Е	186	Total	С	N	О	S	0	3	0
1	E	100	1451	931	247	268	5	0	J 3	
1	F	187	Total	С	N	О	S	0	5	0
1	I'	107	1483	950	251	277	5	0	9	
1	G	184	Total	С	N	О	S	0	3	0
1	G	104	1447	926	245	271	5	U	ა	U

There are 70 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	195	LEU	-	expression tag	UNP Q72L15
A	196	GLU	-	expression tag	UNP Q72L15
A	197	HIS	_	expression tag	UNP Q72L15
A	198	HIS	_	expression tag	UNP Q72L15
A	199	HIS	_	expression tag	UNP Q72L15
A	200	HIS	_	expression tag	UNP Q72L15
A	201	HIS	_	expression tag	UNP Q72L15
A	202	HIS	_	expression tag	UNP Q72L15
A	203	HIS	_	expression tag	UNP Q72L15
A	204	HIS	_	expression tag	UNP Q72L15
В	195	LEU	_	expression tag	UNP Q72L15
В	196	GLU	_	expression tag	UNP Q72L15
В	197	HIS	_	expression tag	UNP Q72L15
В	198	HIS	-	expression tag	UNP Q72L15
В	199	HIS	-	expression tag	UNP Q72L15



Continued from previous page...

Chain	Residue	Modelled Modelled	Actual	Comment	Reference
В	200	HIS	-	expression tag	UNP Q72L15
В	201	HIS	-	expression tag	UNP Q72L15
В	202	HIS	-	expression tag	UNP Q72L15
В	203	HIS	_	expression tag	UNP Q72L15
В	204	HIS	-	expression tag	UNP Q72L15
С	195	LEU	_	expression tag	UNP Q72L15
С	196	GLU	-	expression tag	UNP Q72L15
С	197	HIS	-	expression tag	UNP Q72L15
С	198	HIS	-	expression tag	UNP Q72L15
С	199	HIS	-	expression tag	UNP Q72L15
С	200	HIS	-	expression tag	UNP Q72L15
С	201	HIS	-	expression tag	UNP Q72L15
С	202	HIS	_	expression tag	UNP Q72L15
С	203	HIS	-	expression tag	UNP Q72L15
С	204	HIS	-	expression tag	UNP Q72L15
D	195	LEU	-	expression tag	UNP Q72L15
D	196	GLU	-	expression tag	UNP Q72L15
D	197	HIS	_	expression tag	UNP Q72L15
D	198	HIS	-	expression tag	UNP Q72L15
D	199	HIS	-	expression tag	UNP Q72L15
D	200	HIS	-	expression tag	UNP Q72L15
D	201	HIS	-	expression tag	UNP Q72L15
D	202	HIS	_	expression tag	UNP Q72L15
D	203	HIS	-	expression tag	UNP Q72L15
D	204	HIS	_	expression tag	UNP Q72L15
E	195	LEU	-	expression tag	UNP Q72L15
E	196	GLU	-	expression tag	UNP Q72L15
E	197	HIS	-	expression tag	UNP Q72L15
Е	198	HIS	_	expression tag	UNP Q72L15
Ε	199	HIS	_	expression tag	UNP Q72L15
Ε	200	HIS	_	expression tag	UNP Q72L15
Ε	201	HIS	_	expression tag	UNP Q72L15
Ε	202	HIS	_	expression tag	UNP Q72L15
E	203	HIS	-	expression tag	UNP Q72L15
E	204	HIS	-	expression tag	UNP Q72L15
F	195	LEU	-	expression tag	UNP Q72L15
F	196	GLU	-	expression tag	UNP Q72L15
F	197	HIS	-	expression tag	UNP Q72L15
F	198	HIS	-	expression tag	UNP Q72L15
F	199	HIS	-	expression tag	UNP Q72L15
F	200	HIS	-	expression tag	UNP Q72L15
F	201	HIS	-	expression tag	UNP Q72L15



 $Continued\ from\ previous\ page...$


Chain	Residue	Modelled	Actual	Comment	Reference
F	202	HIS	-	expression tag	UNP Q72L15
F	203	HIS	-	expression tag	UNP Q72L15
F	204	HIS	_	expression tag	UNP Q72L15
G	195	LEU	-	expression tag	UNP Q72L15
G	196	GLU	-	expression tag	UNP Q72L15
G	197	HIS	-	expression tag	UNP Q72L15
G	198	HIS	-	expression tag	UNP Q72L15
G	199	HIS	-	expression tag	UNP Q72L15
G	200	HIS	-	expression tag	UNP Q72L15
G	201	HIS	-	expression tag	UNP Q72L15
G	202	HIS	=	expression tag	UNP Q72L15
G	203	HIS	_	expression tag	UNP Q72L15
G	204	HIS	-	expression tag	UNP Q72L15

• Molecule 2 is a protein called Unknown tripeptide.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf	Trace
2	Н	3	Total C N O 16 9 3 4	0	0	0
2	I	3	Total C N O 16 9 3 4	0	0	0
2	J	3	Total C N O 16 9 3 4	0	0	0
2	K	3	Total C N O 16 9 3 4	0	0	0
2	L	3	Total C N O 16 9 3 4	0	0	0
2	М	3	Total C N O 16 9 3 4	0	0	0
2	N	3	Total C N O 16 9 3 4	0	0	0

• Molecule 3 is DI(HYDROXYETHYL)ETHER (three-letter code: PEG) (formula: $C_4H_{10}O_3$).





Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O 7 4 3	0	0
3	A	1	Total C O 7 4 3	0	0
3	A	1	Total C O 7 4 3	0	0
3	В	1	Total C O 7 4 3	0	0
3	В	1	Total C O 7 4 3	0	0
3	В	1	Total C O 7 4 3	0	0
3	С	1	Total C O 7 4 3	0	0
3	С	1	Total C O 7 4 3	0	0
3	С	1	Total C O 7 4 3	0	0
3	D	1	Total C O 7 4 3	0	0
3	D	1	Total C O 7 4 3	0	0
3	D	1	Total C O 7 4 3	0	0
3	D	1	Total C O 7 4 3	0	0
3	E	1	Total C O 7 4 3	0	0



Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	E	1	Total C O 7 4 3	0	0
3	F	1	Total C O 7 4 3	0	0
3	G	1	Total C O 7 4 3	0	0

• Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	70	Total O 70 70	0	0
4	Н	1	Total O 1 1	0	0
4	В	70	Total O 70 70	0	0
4	I	2	Total O 2 2	0	0
4	С	76	Total O 76 76	0	0
4	J	2	Total O 2 2	0	0
4	D	91	Total O 91 91	0	0
4	K	2	Total O 2 2	0	0
4	E	99	Total O 99 99	0	0
4	L	2	Total O 2 2	0	0
4	F	108	Total O 108 108	0	0
4	М	2	Total O 2 2	0	0
4	G	100	Total O 100 100	0	0
4	N	3	Total O 3 3	0	0





Chain E:

3 Residue-property plots (i)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: ATP-dependent Clp protease proteolytic subunit







Chain M: 67% 33%

REP

• Molecule 2: Unknown tripeptide

Chain N: 67% 33%



4 Data and refinement statistics (i)

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants	105.98Å 162.79Å 107.95Å	Depositor
a, b, c, α , β , γ	90.00° 116.34° 90.00°	Depositor
Resolution (Å)	46.59 - 1.95	Depositor
Resolution (A)	47.11 - 1.95	EDS
% Data completeness	99.3 (46.59-1.95)	Depositor
(in resolution range)	99.3 (47.11-1.95)	EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	1.37 (at 1.95Å)	Xtriage
Refinement program	PHENIX (1.13_2998: ???)	Depositor
D D.	0.186 , 0.211	Depositor
R, R_{free}	0.186 , 0.211	DCC
R_{free} test set	11807 reflections (10.00%)	wwPDB-VP
Wilson B-factor (Å ²)	35.9	Xtriage
Anisotropy	0.497	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.37, 53.2	EDS
L-test for twinning ²	$ < L > = 0.50, < L^2 > = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	11036	wwPDB-VP
Average B, all atoms (Å ²)	40.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 3.56% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of <|L|>, $< L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: PEG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bo	nd lengths	Bond	angles
MIOI	Chain	RMSZ $ $ # $ Z > 5$		RMSZ	# Z > 5
1	A	0.53	1/1471 (0.1%)	0.59	0/1999
1	В	0.48	0/1460	0.58	0/1985
1	С	0.40	0/1461	0.56	0/1987
1	D	0.45	0/1499	0.57	0/2039
1	Е	0.51	0/1476	0.59	0/2005
1	F	0.46	0/1508	0.56	0/2049
1	G	0.48	0/1470	0.60	0/1994
All	All	0.48	1/10345~(0.0%)	0.58	0/14058

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$\operatorname{Observed}(ext{\AA})$	$\operatorname{Ideal}(\operatorname{\AA})$
1	A	141	GLU	CD-OE2	-5.01	1.20	1.25

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1448	0	1463	22	0
1	В	1437	0	1449	16	0
1	С	1438	0	1444	16	0



Continued from previous page...

Mol	Chain	Non-H		H(added)	Clashes	Symm-Clashes
1	D	1473	0	1482	19	0
1	Ε	1451	0	1464	16	0
1	F	1483	0	1490	13	0
1	G	1447	0	1472	11	0
2	Н	16	0	5	2	0
2	I	16	0	5	3	0
2	J	16	0	5	3	0
2	K	16	0	5	0	0
2	L	16	0	5	2	0
2	M	16	0	5	2	0
2	N	16	0	5	2	0
3	A	21	0	30	4	0
3	В	21	0	30	1	0
3	С	21	0	30	0	0
3	D	28	0	40	9	0
3	E	14	0	20	5	0
3	F	7	0	10	1	0
3	G	7	0	10	0	0
4	A	70	0	0	1	0
4	В	70	0	0	0	0
4	С	76	0	0	1	0
4	D	91	0	0	1	0
4	E	99	0	0	1	0
4	F	108	0	0	2	0
4	G	100	0	0	1	0
4	Н	1	0	0	0	0
4	I	2	0	0	0	0
4	J	2	0	0	0	0
4	K	2	0	0	0	0
4	L	2	0	0	0	0
4	M	2	0	0	0	0
4	N	3	0	0	0	0
All	All	11036	0	10469	106	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 106 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1			Clash overlap (Å)
1:E:155:LYS:HZ2	E:155:LYS:HZ2 3:E:302:PEG:H42		1.06



Continued from previous page...

Atom-1	Atom-2	$egin{aligned} ext{Interatomic} \ ext{distance} \ (ext{Å}) \end{aligned}$	$egin{aligned} ext{Clash} \ ext{overlap } (ext{Å}) \end{aligned}$
1:E:155:LYS:NZ	3:E:302:PEG:H42	1.70	1.06
1:B:101:VAL:HG22	1:B:153:LEU:HD21	1.38	1.01
1:C:139:ALA:O	1:C:143:LEU:HD23	1.67	0.95
1:B:97:SER:OG	2:I:3:UNK:C	2.21	0.88

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	A	186/204~(91%)	183 (98%)	3 (2%)	0	100	100
1	В	184/204 (90%)	182 (99%)	2 (1%)	0	100	100
1	С	186/204 (91%)	183 (98%)	3 (2%)	0	100	100
1	D	188/204 (92%)	185 (98%)	3 (2%)	0	100	100
1	Ε	185/204 (91%)	183 (99%)	2 (1%)	0	100	100
1	F	188/204 (92%)	186 (99%)	2 (1%)	0	100	100
1	G	183/204 (90%)	181 (99%)	2 (1%)	0	100	100
All	All	$1300/1428 \; (91\%)$	1283 (99%)	17 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.



Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	149/167 (89%)	146 (98%)	3 (2%)	55 48
1	В	149/167 (89%)	147 (99%)	2 (1%)	69 65
1	$^{\mathrm{C}}$	147/167 (88%)	143 (97%)	4 (3%)	44 34
1	D	152/167 (91%)	150 (99%)	2 (1%)	69 65
1	E	149/167 (89%)	146 (98%)	3 (2%)	55 48
1	F	154/167 (92%)	150 (97%)	4 (3%)	46 36
1	G	152/167 (91%)	150 (99%)	2 (1%)	69 65
All	All	$1052/1169 \ (90\%)$	1032 (98%)	20 (2%)	65 50

5 of 20 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	125[A]	TRP
1	D	125[B]	TRP
1	F	94[B]	MET
1	С	122	HIS
1	С	125	TRP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	38	GLN
1	F	187	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no carbohydrates in this entry.



5.6 Ligand geometry (i)

17 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	T	Chain	Dog	T : 1-	В	ond leng	$_{ m gths}$	В	ond ang	gles
Mol	Type	Chain	Res	Link	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	PEG	С	301	-	6,6,6	0.48	0	5,5,5	0.27	0
3	PEG	D	304	-	6,6,6	0.48	0	5,5,5	0.32	0
3	PEG	D	303	-	6,6,6	0.46	0	5,5,5	0.52	0
3	PEG	E	301	-	6,6,6	0.49	0	5,5,5	0.44	0
3	PEG	A	301	_	6,6,6	0.47	0	5,5,5	0.35	0
3	PEG	В	303	-	6,6,6	0.48	0	5,5,5	0.30	0
3	PEG	G	301	-	6,6,6	0.49	0	5,5,5	0.24	0
3	PEG	E	302	-	6,6,6	0.47	0	5,5,5	0.23	0
3	PEG	A	303	-	6,6,6	0.48	0	5,5,5	0.36	0
3	PEG	A	302	-	6,6,6	0.47	0	5,5,5	0.18	0
3	PEG	В	302	-	6,6,6	0.50	0	5,5,5	0.24	0
3	PEG	С	302	-	6,6,6	0.47	0	5,5,5	0.29	0
3	PEG	D	302	-	6,6,6	0.48	0	5,5,5	0.43	0
3	PEG	F	301	-	6,6,6	0.47	0	5,5,5	0.23	0
3	PEG	D	301	_	6,6,6	0.48	0	5,5,5	0.23	0
3	PEG	В	301	-	6,6,6	0.45	0	5,5,5	0.35	0
3	PEG	С	303	_	6,6,6	0.46	0	5,5,5	0.27	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	PEG	С	301	_	-	1/4/4/4	1
3	PEG	D	304	-	-	3/4/4/4	-
3	PEG	D	303	-	-	2/4/4/4	-
3	PEG	Ε	301	_	-	3/4/4/4	1
3	PEG	A	301	_	-	2/4/4/4	-
3	PEG	В	303	_	-	2/4/4/4	-



 $Continued\ from\ previous\ page...$

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	PEG	G	301	_	-	2/4/4/4	-
3	PEG	Е	302	-	-	3/4/4/4	-
3	PEG	A	303	-	-	2/4/4/4	-
3	PEG	A	302	-	-	2/4/4/4	-
3	PEG	В	302	_	-	1/4/4/4	-
3	PEG	С	302	_	-	2/4/4/4	-
3	PEG	D	302	-	-	3/4/4/4	-
3	PEG	F	301	-	-	2/4/4/4	-
3	PEG	D	301	_	-	3/4/4/4	-
3	PEG	В	301	-	-	2/4/4/4	-
3	PEG	С	303	_	-	1/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

5 of 36 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	С	301	PEG	O2-C3-C4-O4
3	D	301	PEG	O1-C1-C2-O2
3	E	301	PEG	O2-C3-C4-O4
3	A	302	PEG	O1-C1-C2-O2
3	С	302	PEG	O2-C3-C4-O4

There are no ring outliers.

9 monomers are involved in 20 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	D	304	PEG	1	0
3	D	303	PEG	1	0
3	В	303	PEG	1	0
3	E	302	PEG	5	0
3	A	303	PEG	2	0
3	A	302	PEG	2	0
3	D	302	PEG	6	0
3	F	301	PEG	1	0
3	D	301	PEG	1	0



5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<rsrz></rsrz>	$\#\mathrm{RSRZ}{>}2$	$OWAB(\AA^2)$	Q < 0.9
1	A	187/204 (91%)	0.19	12 (6%) 19 28	29, 37, 72, 94	1 (0%)
1	В	186/204 (91%)	0.12	10 (5%) 25 34	30, 39, 65, 109	1 (0%)
1	С	188/204 (92%)	-0.07	9 (4%) 30 40	29, 39, 69, 93	1 (0%)
1	D	187/204 (91%)	0.07	9 (4%) 30 40	26, 34, 61, 101	0
1	Е	186/204 (91%)	-0.01	10 (5%) 25 34	24, 32, 66, 104	0
1	F	187/204 (91%)	0.04	9 (4%) 30 40	24, 33, 55, 101	0
1	G	184/204 (90%)	-0.06	8 (4%) 35 45	25, 35, 60, 85	1 (0%)
2	Н	0/3	-	-	-	-
2	I	0/3	-	-	-	-
2	J	0/3	-	-	-	-
2	K	0/3	-	-	-	-
2	L	0/3	-	-	-	_
2	M	0/3	-	-	-	
2	N	0/3	_	-	-	_
All	All	1305/1449~(90%)	0.04	67 (5%) 28 37	24, 36, 67, 109	4 (0%)

The worst 5 of 67 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	E	7	ILE	9.2
1	В	7	ILE	7.4
1	E	15	ARG	6.9
1	В	2	VAL	6.4
1	В	16	VAL	6.1

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.



6.3 Carbohydrates (i)

There are no carbohydrates in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\operatorname{B-factors}({ ext{\AA}}^2)$	Q<0.9
3	PEG	Е	301	7/7	0.57	0.16	55,60,69,69	0
3	PEG	В	303	7/7	0.57	0.22	56,63,73,75	0
3	PEG	F	301	7/7	0.59	0.22	67,73,75,75	0
3	PEG	A	302	7/7	0.72	0.32	61,70,79,80	0
3	PEG	A	303	7/7	0.72	0.26	70,71,77,80	0
3	PEG	В	302	7/7	0.77	0.15	65,67,72,73	0
3	PEG	G	301	7/7	0.78	0.17	69,70,72,73	0
3	PEG	D	304	7/7	0.80	0.20	64,67,75,76	0
3	PEG	D	302	7/7	0.81	0.27	38,46,57,60	0
3	PEG	Е	302	7/7	0.82	0.14	57,58,60,60	0
3	PEG	С	302	7/7	0.82	0.14	61,67,71,72	0
3	PEG	С	303	7/7	0.85	0.17	62,62,65,66	0
3	PEG	D	301	7/7	0.86	0.20	52,53,61,63	0
3	PEG	D	303	7/7	0.87	0.24	51,58,72,73	0
3	PEG	A	301	7/7	0.88	0.12	63,64,67,68	0
3	PEG	В	301	7/7	0.90	0.10	59,60,61,63	0
3	PEG	С	301	7/7	0.91	0.15	52,54,57,58	0

6.5 Other polymers (i)

There are no such residues in this entry.

