



# Full wwPDB X-ray Structure Validation Report ⓘ

Jan 15, 2024 – 06:29 pm GMT

PDB ID : 6I58  
Title : Allosteric activation of human prekallikrein by apple domain disc rotation  
Authors : Li, C.; Pathak, M.; McCrae, K.; Dreveny, I.; Emsley, J.  
Deposited on : 2018-11-13  
Resolution : 2.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.4, CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.36  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	3163 (2.60-2.60)
Clashscore	141614	3518 (2.60-2.60)
Ramachandran outliers	138981	3455 (2.60-2.60)
Sidechain outliers	138945	3455 (2.60-2.60)
RSRZ outliers	127900	3104 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	607	 6% 80% 14% . .
2	B	2	 100%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	PEG	A	706	-	-	X	-
5	NAG	A	708	-	-	-	X

## 2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 4753 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Coagulation factor XI.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	585	4602	2903	796	864	39	36	1	0

- Molecule 2 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	B	2	28	16	2	10	0	0	0

- Molecule 3 is CHLORIDE ION (three-letter code: CL) (formula: Cl).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	5	Total	Cl	0	0
			5	5		

- Molecule 4 is DI(HYDROXYETHYL)ETHER (three-letter code: PEG) (formula: C<sub>4</sub>H<sub>10</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			7	4	3		

- Molecule 5 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula:  $C_8H_{15}NO_6$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	1	Total	C	N	O	0	0
			14	8	1	5		
5	A	1	Total	C	N	O	0	0
			14	8	1	5		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	83	Total 83	O 83	0	0


### 3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Coagulation factor XI



- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	80.76Å 80.76Å 251.06Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	37.71 – 2.60 33.16 – 2.60	Depositor EDS
% Data completeness (in resolution range)	98.9 (37.71-2.60) 99.0 (33.16-2.60)	Depositor EDS
$R_{merge}$	0.09	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.42 (at 2.61Å)	Xtrriage
Refinement program	REFMAC 5.8.0158	Depositor
R, $R_{free}$	0.208 , 0.280 0.217 , 0.281	Depositor DCC
$R_{free}$ test set	2641 reflections (10.06%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	70.1	Xtrriage
Anisotropy	0.021	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.32 , 49.2	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	4753	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	75.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.60% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NAG, PEG, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.62	0/4712	0.81	0/6380

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	391	SER	Peptide

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4602	0	4479	57	0
2	B	28	0	25	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	5	0	0	0	0
4	A	7	0	10	4	0
5	A	28	0	26	1	0
6	A	83	0	0	0	0
All	All	4753	0	4540	58	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (58) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:367:LYS:HD2	1:A:368:PRO:HD2	1.19	1.13
1:A:406:GLN:HG3	1:A:471:THR:HG22	1.31	1.06
1:A:390:THR:HG23	1:A:423:ILE:O	1.59	1.01
1:A:509:LYS:HA	1:A:509:LYS:HE2	1.57	0.85
1:A:40:LEU:HD13	1:A:61:LYS:HB2	1.65	0.77
5:A:708:NAG:O7	5:A:708:NAG:O3	2.05	0.73
1:A:371:VAL:HG12	1:A:516:LYS:NZ	2.03	0.73
1:A:40:LEU:H	1:A:40:LEU:HD12	1.56	0.70
1:A:479:ARG:HB3	4:A:706:PEG:H21	1.76	0.66
1:A:370:ILE:HD12	1:A:370:ILE:C	2.14	0.66
1:A:28:CYS:SG	1:A:41:PHE:CD2	2.89	0.66
1:A:355:LEU:HA	1:A:358:MET:HE3	1.79	0.64
1:A:371:VAL:HG13	1:A:380:GLU:CD	2.18	0.64
1:A:371:VAL:HG12	1:A:516:LYS:HZ1	1.63	0.62
1:A:479:ARG:CB	4:A:706:PEG:H21	2.29	0.61
1:A:459:SER:OG	1:A:460:GLY:N	2.31	0.61
1:A:406:GLN:HG3	1:A:471:THR:CG2	2.18	0.60
1:A:317:ALA:CB	1:A:323:GLU:O	2.49	0.60
1:A:90:SER:O	1:A:91:ALA:HB3	2.03	0.57
1:A:323:GLU:HA	1:A:323:GLU:OE2	2.05	0.55
1:A:40:LEU:HD12	1:A:40:LEU:N	2.16	0.55
1:A:40:LEU:CD1	1:A:61:LYS:HB2	2.35	0.54
1:A:40:LEU:HD22	1:A:68:LEU:HD12	1.89	0.54
1:A:364:THR:HG23	1:A:365:LYS:HE3	1.90	0.54
1:A:367:LYS:HB3	1:A:369:ARG:NH1	2.23	0.53
1:A:29:GLN:HA	1:A:41:PHE:CE1	2.43	0.52
1:A:198:ASP:HB3	1:A:239:LEU:HD12	1.92	0.52
1:A:39:LEU:HD11	1:A:308:ARG:HG2	1.91	0.52
1:A:317:ALA:HB1	1:A:323:GLU:O	2.11	0.51

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:191:VAL:HG22	1:A:251:ILE:HB	1.93	0.50
1:A:474:TYR:CZ	1:A:480:PRO:HG3	2.46	0.49
1:A:367:LYS:CD	1:A:368:PRO:HD2	2.14	0.47
1:A:20:VAL:HB	1:A:156:THR:HG21	1.96	0.47
1:A:126:VAL:HA	1:A:180:LEU:HD11	1.96	0.46
1:A:367:LYS:HD2	1:A:368:PRO:CD	2.14	0.46
1:A:272:PHE:HB3	1:A:355:LEU:HB3	1.97	0.46
1:A:129:HIS:ND1	1:A:155:GLY:HA2	2.31	0.46
1:A:40:LEU:HD13	1:A:61:LYS:CB	2.42	0.45
1:A:532:ARG:NE	1:A:533:GLY:H	2.15	0.45
1:A:252:LYS:O	1:A:253:LYS:CG	2.65	0.44
1:A:325:LYS:HD3	1:A:325:LYS:HA	1.65	0.44
1:A:479:ARG:CB	4:A:706:PEG:C2	2.97	0.43
1:A:371:VAL:HG13	1:A:380:GLU:OE2	2.17	0.43
1:A:87:HIS:C	1:A:87:HIS:HD1	2.22	0.43
1:A:479:ARG:HB3	4:A:706:PEG:C2	2.46	0.42
1:A:415:PHE:CZ	1:A:466:LEU:HD21	2.53	0.42
1:A:263:GLN:H	1:A:263:GLN:HG3	1.35	0.42
1:A:370:ILE:O	1:A:370:ILE:HG23	2.19	0.42
1:A:530:ARG:HD3	1:A:586:ARG:HA	2.01	0.42
1:A:186:ILE:CG2	1:A:257:LEU:HD21	2.49	0.41
1:A:298:ALA:O	1:A:302:LEU:HD23	2.21	0.41
1:A:186:ILE:HG23	1:A:257:LEU:HD21	2.02	0.41
1:A:532:ARG:HD2	1:A:532:ARG:HA	1.27	0.41
1:A:252:LYS:O	1:A:253:LYS:HG2	2.21	0.41
1:A:254:SER:O	1:A:255:LYS:HB2	2.20	0.41
1:A:357:LYS:HA	1:A:357:LYS:HD3	1.92	0.40
1:A:409:LEU:HD12	1:A:409:LEU:HA	1.86	0.40
1:A:564:LYS:HB2	1:A:569:TRP:CZ3	2.56	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	576/607 (95%)	549 (95%)	27 (5%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	520/535 (97%)	471 (91%)	49 (9%)	8	17

All (49) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	8	LYS
1	A	11	CYS
1	A	40	LEU
1	A	84	GLN
1	A	86	SER
1	A	87	HIS
1	A	90	SER
1	A	95	ASP
1	A	136	ARG
1	A	142	GLU
1	A	156	THR
1	A	166	VAL
1	A	173	LYS
1	A	175	CYS
1	A	231	GLU
1	A	240	LYS
1	A	246	LEU
1	A	252	LYS
1	A	253	LYS
1	A	255	LYS
1	A	257	LEU
1	A	261	SER
1	A	263	GLN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	294	LYS
1	A	323	GLU
1	A	325	LYS
1	A	337	SER
1	A	365	LYS
1	A	370	ILE
1	A	387	LEU
1	A	434	SER
1	A	456	MET
1	A	459	SER
1	A	476	ASP
1	A	499	THR
1	A	503	TYR
1	A	508	ASP
1	A	509	LYS
1	A	510	ILE
1	A	515	GLN
1	A	532	ARG
1	A	534	HIS
1	A	541	ILE
1	A	545	TYR
1	A	554	LYS
1	A	577	TRP
1	A	579	GLU
1	A	594	VAL
1	A	604	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

2 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	NAG	B	1	2,1	14,14,15	0.77	0	17,19,21	1.56	3 (17%)
2	NAG	B	2	2	14,14,15	0.65	0	17,19,21	1.06	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	B	1	2,1	-	5/6/23/26	0/1/1/1
2	NAG	B	2	2	-	2/6/23/26	0/1/1/1

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	1	NAG	C1-O5-C5	3.41	116.82	112.19
2	B	1	NAG	O4-C4-C3	-3.23	102.89	110.35
2	B	2	NAG	O5-C1-C2	-2.85	106.78	111.29
2	B	1	NAG	O5-C5-C6	2.84	111.65	107.20

There are no chirality outliers.


All (7) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	1	NAG	C3-C2-N2-C7
2	B	1	NAG	C8-C7-N2-C2
2	B	1	NAG	O7-C7-N2-C2
2	B	1	NAG	C4-C5-C6-O6
2	B	1	NAG	O5-C5-C6-O6
2	B	2	NAG	O5-C5-C6-O6
2	B	2	NAG	C4-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



## 5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 5 are monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	PEG	A	706	-	6,6,6	0.43	0	5,5,5	0.31	0
5	NAG	A	708	1	14,14,15	0.30	0	17,19,21	0.62	0
5	NAG	A	707	1	14,14,15	0.67	0	17,19,21	2.51	4 (23%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	PEG	A	706	-	-	1/4/4/4	-
5	NAG	A	708	1	-	6/6/23/26	0/1/1/1
5	NAG	A	707	1	-	3/6/23/26	0/1/1/1

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	707	NAG	C2-N2-C7	6.99	132.86	122.90
5	A	707	NAG	C8-C7-N2	5.15	124.82	116.10
5	A	707	NAG	C1-O5-C5	3.80	117.34	112.19
5	A	707	NAG	O7-C7-C8	-2.92	116.63	122.06

There are no chirality outliers.

All (10) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	A	707	NAG	C3-C2-N2-C7
5	A	708	NAG	C1-C2-N2-C7
5	A	708	NAG	C8-C7-N2-C2
5	A	707	NAG	C8-C7-N2-C2
5	A	707	NAG	O7-C7-N2-C2
5	A	708	NAG	O7-C7-N2-C2
5	A	708	NAG	C4-C5-C6-O6
4	A	706	PEG	O1-C1-C2-O2
5	A	708	NAG	O5-C5-C6-O6
5	A	708	NAG	C3-C2-N2-C7

There are no ring outliers.

2 monomers are involved in 5 short contacts:



Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	706	PEG	4	0
5	A	708	NAG	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	585/607 (96%)	0.25	34 (5%) <b>23</b> <b>17</b>	41, 70, 113, 177	19 (3%)

All (34) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	321	CYS	4.4
1	A	323	GLU	4.3
1	A	371	VAL	4.2
1	A	324	GLY	4.1
1	A	322	ASN	3.9
1	A	370	ILE	3.5
1	A	509	LYS	3.4
1	A	534	HIS	3.3
1	A	531	TYR	3.0
1	A	111	VAL	2.9
1	A	48	PRO	2.8
1	A	508	ASP	2.8
1	A	368	PRO	2.8
1	A	586	ARG	2.8
1	A	410	THR	2.6
1	A	75	ALA	2.6
1	A	385	VAL	2.5
1	A	533	GLY	2.5
1	A	377	VAL	2.5
1	A	409	LEU	2.4
1	A	400	GLY	2.4
1	A	587	PRO	2.4
1	A	579	GLU	2.4
1	A	317	ALA	2.3
1	A	391	SER	2.3
1	A	378	ARG	2.3
1	A	72	ASN	2.2

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	372	GLY	2.2
1	A	89	ILE	2.1
1	A	578	GLY	2.1
1	A	2	CYS	2.1
1	A	399	GLY	2.1
1	A	470	THR	2.0
1	A	374	THR	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)


There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	NAG	B	1	14/15	0.85	0.14	54,59,73,73	14
2	NAG	B	2	14/15	0.89	0.23	50,57,59,61	14

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.



## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
5	NAG	A	707	14/15	0.64	0.33	101,113,123,123	14
5	NAG	A	708	14/15	0.64	0.48	117,136,148,150	0
3	CL	A	705	1/1	0.77	0.18	95,95,95,95	0
3	CL	A	704	1/1	0.80	0.11	88,88,88,88	0
4	PEG	A	706	7/7	0.83	0.18	54,55,59,60	1
3	CL	A	701	1/1	0.84	0.21	89,89,89,89	0
3	CL	A	703	1/1	0.87	0.51	90,90,90,90	0
3	CL	A	702	1/1	0.88	0.29	80,80,80,80	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.