



# Full wwPDB NMR Structure Validation Report ⓘ

Oct 17, 2021 – 09:47 AM EDT

PDB ID : 1I8H  
Title : SOLUTION STRUCTURE OF PIN1 WW DOMAIN COMPLEXED WITH HUMAN TAU PHOSPHOTHREONINE PEPTIDE  
Authors : Wintjens, R.; Wieruszeski, J.-M.; Drobecq, H.; Lippens, G.; Landrieu, I.  
Deposited on : 2001-03-14

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
RCI : v\_1n\_11\_5\_13\_A (Berjanski et al., 2005)  
PANAV : Wang et al. (2010)  
ShiftChecker : 2.23.2  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.23.2


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*SOLUTION NMR*

The overall completeness of chemical shifts assignment was not calculated.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	158937	12864
Ramachandran outliers	154571	11451
Sidechain outliers	154315	11428

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	13	15% 15% 8% 62%
2	B	39	51% 18% 15% 15%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA and RNA chains that are outliers for geometric criteria:

Mol	Chain	Compound	Res	Total models with violations	
				Chirality	Geometry
1	A	TPO	7	10	-

## 2 Ensemble composition and analysis

This entry contains 10 models. Model 6 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *closest to the average*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:5-A:6, A:8-A:10, B:1-B:33 (38)	0.67	6

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 2 clusters and 1 single-model cluster was found.

Cluster number	Models
1	1, 2, 3, 4, 6, 8, 9
2	5, 10
Single-model clusters	7

### 3 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 823 atoms, of which 407 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called MICROTUBULE-ASSOCIATED PROTEIN TAU.

Mol	Chain	Residues	Atoms					Trace	
			Total	C	H	N	O		P
1	A	13	211	61	110	18	21	1	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	3	SER	ALA	engineered mutation	UNP P10636
A	7	TPO	THR	modified residue	UNP P10636

- Molecule 2 is a protein called PEPTIDYL-PROLYL CIS-TRANS ISOMERASE NIMA-INTERACTING 1.

Mol	Chain	Residues	Atoms					Trace	
			Total	C	H	N	O		S
2	B	39	612	194	297	61	59	1	0

## 4 Residue-property plots

### 4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: MICROTUBULE-ASSOCIATED PROTEIN TAU

Chain A: 



- Molecule 2: PEPTIDYL-PROLYL CIS-TRANS ISOMERASE NIMA-INTERACTING 1

Chain B: 



### 4.2 Scores per residue for each member of the ensemble

Colouring as in section 4.1 above.

#### 4.2.1 Score per residue for model 1

- Molecule 1: MICROTUBULE-ASSOCIATED PROTEIN TAU

Chain A: 



- Molecule 2: PEPTIDYL-PROLYL CIS-TRANS ISOMERASE NIMA-INTERACTING 1

Chain B: 




### 4.2.2 Score per residue for model 2

- Molecule 1: MICROTUBULE-ASSOCIATED PROTEIN TAU




- Molecule 2: PEPTIDYL-PROLYL CIS-TRANS ISOMERASE NIMA-INTERACTING 1




### 4.2.3 Score per residue for model 3

- Molecule 1: MICROTUBULE-ASSOCIATED PROTEIN TAU




- Molecule 2: PEPTIDYL-PROLYL CIS-TRANS ISOMERASE NIMA-INTERACTING 1




### 4.2.4 Score per residue for model 4

- Molecule 1: MICROTUBULE-ASSOCIATED PROTEIN TAU




- Molecule 2: PEPTIDYL-PROLYL CIS-TRANS ISOMERASE NIMA-INTERACTING 1




### 4.2.5 Score per residue for model 5

- Molecule 1: MICROTUBULE-ASSOCIATED PROTEIN TAU




- Molecule 2: PEPTIDYL-PROLYL CIS-TRANS ISOMERASE NIMA-INTERACTING 1




### 4.2.6 Score per residue for model 6 (medoid)

- Molecule 1: MICROTUBULE-ASSOCIATED PROTEIN TAU




- Molecule 2: PEPTIDYL-PROLYL CIS-TRANS ISOMERASE NIMA-INTERACTING 1




### 4.2.7 Score per residue for model 7

- Molecule 1: MICROTUBULE-ASSOCIATED PROTEIN TAU




- Molecule 2: PEPTIDYL-PROLYL CIS-TRANS ISOMERASE NIMA-INTERACTING 1




### 4.2.8 Score per residue for model 8

- Molecule 1: MICROTUBULE-ASSOCIATED PROTEIN TAU




- Molecule 2: PEPTIDYL-PROLYL CIS-TRANS ISOMERASE NIMA-INTERACTING 1




### 4.2.9 Score per residue for model 9

- Molecule 1: MICROTUBULE-ASSOCIATED PROTEIN TAU




- Molecule 2: PEPTIDYL-PROLYL CIS-TRANS ISOMERASE NIMA-INTERACTING 1




### 4.2.10 Score per residue for model 10

- Molecule 1: MICROTUBULE-ASSOCIATED PROTEIN TAU



- Molecule 2: PEPTIDYL-PROLYL CIS-TRANS ISOMERASE NIMA-INTERACTING 1





## 5 Refinement protocol and experimental data overview

The models were refined using the following method: *distance geometry simulated annealing*.

Of the 50 calculated structures, 10 were deposited, based on the following criterion: *structures with the lowest energy*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
X-PLOR	structure solution	3.851
Discover	refinement	2.98

No chemical shift data was provided.

## 6 Model quality i

### 6.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: TPO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#Z>5	RMSZ	#Z>5
1	A	1.36±0.03	0±0/42 ( 0.0± 0.0%)	1.89±0.19	1±0/55 ( 2.0± 0.5%)
2	B	1.31±0.02	0±0/291 ( 0.0± 0.0%)	1.78±0.06	10±2/394 ( 2.7± 0.6%)
All	All	1.32	0/3330 ( 0.0%)	1.80	116/4490 ( 2.6%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	Chirality	Planarity
1	A	0.0±0.0	0.9±0.7
2	B	0.0±0.0	4.1±1.3
All	All	0	50

There are no bond-length outliers.

All unique angle outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
2	B	12	ARG	NE-CZ-NH1	10.08	125.34	120.30	10	8
2	B	9	ARG	NE-CZ-NH1	9.83	125.22	120.30	1	10
1	A	6	ARG	NE-CZ-NH1	9.18	124.89	120.30	1	10
2	B	31	ARG	NE-CZ-NH1	8.25	124.42	120.30	8	10
2	B	16	ARG	NE-CZ-NH1	7.86	124.23	120.30	5	10
2	B	31	ARG	N-CA-CB	-7.41	97.26	110.60	2	3
2	B	31	ARG	NE-CZ-NH2	-6.95	116.82	120.30	10	2
1	A	6	ARG	NE-CZ-NH2	-6.84	116.88	120.30	5	1
2	B	12	ARG	NE-CZ-NH2	-6.71	116.95	120.30	4	4
2	B	16	ARG	NE-CZ-NH2	-6.61	117.00	120.30	9	3

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
2	B	9	ARG	NE-CZ-NH2	-6.55	117.02	120.30	1	2
2	B	6	TRP	CD1-NE1-CE2	-6.43	103.21	109.00	6	10
2	B	17	VAL	CB-CA-C	6.12	123.03	111.40	7	5
2	B	17	VAL	CA-CB-CG1	5.96	119.85	110.90	10	4
2	B	6	TRP	N-CA-CB	-5.84	100.09	110.60	2	10
2	B	29	TRP	CD1-NE1-CE2	-5.74	103.84	109.00	6	9
2	B	24	THR	CA-CB-CG2	5.68	120.35	112.40	8	1
2	B	29	TRP	CA-CB-CG	5.58	124.31	113.70	3	5
2	B	25	ASN	CB-CA-C	5.55	121.49	110.40	5	1
2	B	18	TYR	CB-CG-CD1	-5.52	117.69	121.00	4	3
2	B	29	TRP	CB-CA-C	5.25	120.89	110.40	7	1
2	B	31	ARG	CA-CB-CG	5.23	124.92	113.40	7	1
2	B	18	TYR	CB-CG-CD2	-5.21	117.87	121.00	1	1
2	B	17	VAL	N-CA-CB	5.17	122.89	111.50	1	1
2	B	10	MET	N-CA-CB	-5.07	101.48	110.60	8	1

There are no chirality outliers.

All unique planar outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Group	Models (Total)
2	B	18	TYR	Sidechain	10
2	B	9	ARG	Peptide,Sidechain	8
2	B	6	TRP	Peptide	6
1	A	6	ARG	Sidechain,Peptide	5
2	B	12	ARG	Sidechain,Peptide	4
2	B	10	MET	Peptide	2
2	B	31	ARG	Sidechain	2
2	B	16	ARG	Peptide	2
1	A	10	LYS	Peptide	2
1	A	9	PRO	Peptide	1
2	B	19	TYR	Sidechain	1
2	B	7	GLU	Peptide	1
2	B	29	TRP	Peptide	1

## 6.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	41	49	49	0±0
2	B	280	270	270	0±0
All	All	3210	3190	3190	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 0.

All unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
2:B:9:ARG:O	2:B:17:VAL:HG13	0.50	2.07	7	1
2:B:6:TRP:CZ2	2:B:32:PRO:HG3	0.41	2.51	9	1

## 6.3 Torsion angles [i](#)

### 6.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	5/13 (38%)	3±1 (58±17%)	1±1 (14±16%)	1±0 (28±10%)	0	1
2	B	32/39 (82%)	23±1 (72±4%)	6±2 (17±5%)	3±1 (11±3%)	1	9
All	All	370/520 (71%)	260 (70%)	62 (17%)	48 (13%)	1	6

All 12 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
2	B	17	VAL	10
2	B	18	TYR	9
1	A	9	PRO	5
2	B	25	ASN	5
1	A	8	PRO	5
1	A	5	VAL	4
2	B	11	SER	3
2	B	2	LEU	2
2	B	14	SER	2
2	B	32	PRO	1

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Models (Total)
2	B	26	ALA	1
2	B	31	ARG	1

### 6.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	5/12 (42%)	4±1 (80±15%)	1±1 (20±15%)	4	34
2	B	30/34 (88%)	23±1 (78±4%)	7±1 (22±4%)	3	30
All	All	350/460 (76%)	274 (78%)	76 (22%)	3	30

All 18 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
2	B	8	LYS	10
2	B	21	ASN	10
2	B	29	TRP	9
2	B	7	GLU	7
2	B	17	VAL	7
1	A	6	ARG	5
1	A	10	LYS	5
2	B	10	MET	4
2	B	1	LYS	3
2	B	22	HIS	3
2	B	25	ASN	3
2	B	30	GLU	2
2	B	31	ARG	2
2	B	12	ARG	2
2	B	2	LEU	1
2	B	18	TYR	1
2	B	16	ARG	1
2	B	24	THR	1

### 6.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds for which Mogul statistics could be retrieved, the number of bonds that are observed in the model and the number of bonds that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length is the number of standard deviations the observed value is removed from the expected value. A bond length with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the average root-mean-square of all Z scores of the bond lengths.

Mol	Type	Chain	Res	Link	Bond lengths		
					Counts	RMSZ	#Z>2
1	TPO	A	7	1	8,10,11	1.13±0.11	0±0 (2±5%)

In the following table, the Counts columns list the number of angles for which Mogul statistics could be retrieved, the number of angles that are observed in the model and the number of angles that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond angle is the number of standard deviations the observed value is removed from the expected value. A bond angle with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the average root-mean-square of all Z scores of the bond angles.

Mol	Type	Chain	Res	Link	Bond angles		
					Counts	RMSZ	#Z>2
1	TPO	A	7	1	10,14,16	1.17±0.09	1±0 (12±4%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	TPO	A	7	1	1±0,1,3,4	0±0,9,11,13	-

All unique bond outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)	Models	
								Worst	Total
1	A	7	TPO	CB-CA	2.15	1.58	1.53	1	1
1	A	7	TPO	P-OG1	2.11	1.63	1.59	2	1

All unique angle outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
1	A	7	TPO	O-C-CA	2.80	117.43	124.78	1	10
1	A	7	TPO	CG2-CB-CA	2.29	117.68	113.16	2	2

All unique chiral outliers are listed below.

Mol	Chain	Res	Type	Atoms	Models (Total)
1	A	7	TPO	CB	10

There are no torsion outliers.

There are no ring outliers.

## 6.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 6.7 Other polymers [i](#)

There are no such molecules in this entry.

## 6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 7 Chemical shift validation

No chemical shift data were provided