



# Full wwPDB X-ray Structure Validation Report ⓘ

May 26, 2020 – 02:55 pm BST

PDB ID : 1I9Y  
Title : CRYSTAL STRUCTURE OF INOSITOL POLYPHOSPHATE 5-PHOSPHATASE DOMAIN (IPP5C) OF SPSYNAPTOJANIN  
Authors : Tsujishita, Y.; Guo, S.; Stolz, L.; York, J.D.; Hurley, J.H.  
Deposited on : 2001-03-21  
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	9178 (2.00-2.00)
Ramachandran outliers	138981	9054 (2.00-2.00)
Sidechain outliers	138945	9053 (2.00-2.00)
RSRZ outliers	127900	7900 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	347	

## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 2952 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called PHOSPHATIDYLINOSITOL PHOSPHATE PHOSPHATASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	336	2766	1771	473	513	9	0	1	0


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	186	Total	O	0	0
			186	186		

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: PHOSPHATIDYLINOSITOL PHOSPHATE PHOSPHATASE



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	148.97Å 67.41Å 51.44Å 90.00° 106.40° 90.00°	Depositor
Resolution (Å)	30.00 – 2.00 27.83 – 2.00	Depositor EDS
% Data completeness (in resolution range)	(Not available) (30.00-2.00) 96.0 (27.83-2.00)	Depositor EDS
$R_{merge}$	0.04	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	5.97 (at 1.99Å)	Xtrriage
Refinement program	CNS 1.0	Depositor
R, $R_{free}$	0.195 , 0.225 0.206 , (Not available)	Depositor DCC
$R_{free}$ test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	22.1	Xtrriage
Anisotropy	0.996	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 48.4	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	2952	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	32.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.33% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality i

### 5.1 Standard geometry i

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.87	2/2835 (0.1%)	1.02	16/3834 (0.4%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	817	ARG	CG-CD	-5.86	1.37	1.51
1	A	715	ARG	CZ-NH1	5.07	1.39	1.33

All (16) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	639	ARG	NE-CZ-NH2	-14.56	113.02	120.30
1	A	639	ARG	NE-CZ-NH1	13.86	127.23	120.30
1	A	715	ARG	NE-CZ-NH2	-10.08	115.26	120.30
1	A	715	ARG	NE-CZ-NH1	7.58	124.09	120.30
1	A	817	ARG	CA-CB-CG	-6.54	99.01	113.40
1	A	794	TYR	CA-C-N	-6.52	102.85	117.20
1	A	817	ARG	NE-CZ-NH2	-6.45	117.08	120.30
1	A	794	TYR	C-N-CA	6.12	137.00	121.70
1	A	794	TYR	O-C-N	6.08	132.43	122.70
1	A	721	LEU	CA-CB-CG	5.97	129.03	115.30
1	A	711	ASP	CB-CG-OD1	-5.82	113.06	118.30
1	A	639	ARG	CB-CG-CD	5.40	125.64	111.60
1	A	710	ARG	NE-CZ-NH2	-5.40	117.60	120.30
1	A	584	ASN	N-CA-C	-5.32	96.64	111.00

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	847	GLU	N-CA-C	-5.27	96.77	111.00
1	A	639	ARG	CD-NE-CZ	5.02	130.63	123.60

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	707	TYR	Sidechain
1	A	714	TYR	Sidechain

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2766	0	2687	72	0
2	A	186	0	0	7	0
All	All	2952	0	2687	72	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 13.

All (72) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:583:GLU:HB2	1:A:585:THR:HG23	1.51	0.93
1:A:795:LYS:HG3	1:A:804:ASP:HB2	1.53	0.88
1:A:816:ASP:O	1:A:817:ARG:HG3	1.75	0.87
1:A:809[A]:HIS:C	1:A:810:ARG:N	2.34	0.81
1:A:808:LYS:HE2	1:A:808:LYS:HA	1.63	0.81
1:A:808:LYS:C	1:A:809[B]:HIS:N	2.33	0.80
1:A:583:GLU:HG2	1:A:585:THR:OG1	1.81	0.79
1:A:639:ARG:HD3	2:A:161:HOH:O	1.89	0.73
1:A:795:LYS:HB3	2:A:13:HOH:O	1.87	0.72
1:A:582:PRO:HG2	1:A:583:GLU:H	1.52	0.72
1:A:717:ILE:HG23	1:A:721:LEU:HD22	1.73	0.69

Continued on next page...

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:555:HIS:CD2	1:A:851:VAL:HG22	2.28	0.68
1:A:581:PHE:HE1	1:A:624:LEU:HG	1.59	0.68
1:A:816:ASP:OD2	1:A:839:HIS:HD2	1.78	0.67
1:A:548:ARG:HH22	1:A:861:LEU:HD22	1.60	0.67
1:A:776:THR:OG1	1:A:778:LYS:HG3	1.94	0.66
1:A:795:LYS:HG2	1:A:810:ARG:HB2	1.78	0.65
1:A:717:ILE:CG2	1:A:721:LEU:HD22	2.26	0.65
1:A:577:GLU:H	1:A:577:GLU:CD	2.01	0.64
1:A:808:LYS:O	1:A:809[A]:HIS:C	2.36	0.63
1:A:537:ILE:O	1:A:541:VAL:HG23	1.98	0.63
1:A:699:HIS:HD2	2:A:163:HOH:O	1.80	0.63
1:A:582:PRO:HG2	1:A:583:GLU:N	2.16	0.60
1:A:670:THR:HG23	1:A:680:GLY:HA2	1.83	0.59
1:A:809[A]:HIS:O	1:A:810:ARG:N	2.36	0.57
1:A:750:GLU:CD	1:A:750:GLU:H	2.05	0.57
1:A:869:ARG:O	1:A:873:VAL:HG23	2.06	0.56
1:A:678:ASN:HB2	2:A:24:HOH:O	2.07	0.55
1:A:861:LEU:O	1:A:865:LEU:HD23	2.07	0.54
1:A:548:ARG:NH2	1:A:861:LEU:HD22	2.23	0.54
1:A:869:ARG:O	1:A:869:ARG:HD3	2.08	0.54
1:A:548:ARG:NH2	1:A:861:LEU:HD13	2.23	0.53
1:A:839:HIS:HE1	2:A:177:HOH:O	1.92	0.52
1:A:756:ILE:HG13	1:A:757:ALA:N	2.24	0.52
1:A:572:ALA:HA	1:A:617:TRP:CZ2	2.45	0.51
1:A:644:VAL:HG12	1:A:645:GLY:N	2.25	0.51
1:A:799:GLY:HA2	1:A:835:TYR:HB3	1.93	0.50
1:A:537:ILE:HG23	1:A:869:ARG:CG	2.41	0.50
1:A:794:TYR:CG	1:A:795:LYS:N	2.78	0.50
1:A:809[A]:HIS:CA	1:A:810:ARG:N	2.74	0.50
1:A:808:LYS:C	1:A:809[B]:HIS:CB	2.80	0.50
1:A:785:GLU:OE2	1:A:817:ARG:HG2	2.12	0.49
1:A:565:TYR:CG	1:A:841:PRO:HB3	2.48	0.49
1:A:710:ARG:HB3	1:A:743:TYR:CZ	2.48	0.49
1:A:537:ILE:HG23	1:A:869:ARG:HG3	1.95	0.47
1:A:666:THR:HG23	1:A:721:LEU:HD12	1.95	0.47
1:A:581:PHE:CE1	1:A:624:LEU:HG	2.43	0.47
1:A:697:THR:HA	1:A:738:PHE:O	2.16	0.46
1:A:816:ASP:C	1:A:817:ARG:HG3	2.33	0.46
1:A:582:PRO:CG	1:A:583:GLU:H	2.23	0.46
1:A:870:LYS:O	1:A:874:ARG:HG3	2.15	0.46
1:A:750:GLU:CD	1:A:750:GLU:N	2.69	0.46

*Continued on next page...*



Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:582:PRO:CG	1:A:583:GLU:N	2.78	0.45
1:A:640:SER:HA	1:A:648:LEU:O	2.16	0.45
1:A:785:GLU:HB2	1:A:819:LEU:CD2	2.46	0.45
1:A:756:ILE:HD13	1:A:791:PRO:HB3	1.99	0.45
1:A:644:VAL:HG12	1:A:645:GLY:H	1.82	0.44
1:A:808:LYS:C	1:A:809[B]:HIS:HB3	2.37	0.44
1:A:614:ARG:HD3	1:A:642:GLN:OE1	2.18	0.44
1:A:645:GLY:HA3	2:A:90:HOH:O	2.18	0.44
1:A:672:LEU:HD22	1:A:672:LEU:HA	1.68	0.43
1:A:583:GLU:HB3	2:A:56:HOH:O	2.18	0.43
1:A:551:GLU:O	1:A:857:LYS:HD2	2.18	0.43
1:A:548:ARG:HB3	1:A:551:GLU:OE2	2.19	0.43
1:A:672:LEU:HB2	1:A:712:HIS:ND1	2.33	0.43
1:A:581:PHE:HA	1:A:583:GLU:OE2	2.19	0.42
1:A:828:SER:O	1:A:844:ALA:HA	2.19	0.42
1:A:538:HIS:CD2	1:A:862:PHE:HZ	2.39	0.41
1:A:581:PHE:CZ	1:A:628:LYS:HG3	2.56	0.41
1:A:582:PRO:HG3	1:A:843:TYR:CZ	2.55	0.41
1:A:808:LYS:HE2	1:A:808:LYS:CA	2.39	0.41
1:A:850:ILE:HD12	1:A:850:ILE:HA	1.79	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	329/347 (95%)	315 (96%)	10 (3%)	4 (1%)	<b>13</b>   <b>7</b>

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	804	ASP

Continued on next page...

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	583	GLU
1	A	609	ALA
1	A	644	VAL

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	301/309 (97%)	287 (95%)	14 (5%)	26 22

All (14) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	557	ASN
1	A	580	LEU
1	A	583	GLU
1	A	672	LEU
1	A	685	ARG
1	A	699	HIS
1	A	721	LEU
1	A	756	ILE
1	A	786	LEU
1	A	792	PRO
1	A	795	LYS
1	A	850	ILE
1	A	859	LYS
1	A	869	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	542	ASN
1	A	555	HIS
1	A	578	ASN
1	A	699	HIS

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	827	HIS
1	A	839	HIS
1	A	849	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

### 5.7 Other polymers [i](#)

There are no such residues in this entry.

### 5.8 Polymer linkage issues [i](#)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
1	A	2

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	A	809[A]:HIS	C	810:ARG	N	2.34
1	A	808:LYS	C	809[B]:HIS	N	2.33

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	336/347 (96%)	0.32	34 (10%) <b>7</b> <b>6</b>	15, 26, 71, 77	0

All (34) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	874	ARG	6.0
1	A	583	GLU	5.1
1	A	877	SER	4.9
1	A	608	SER	4.4
1	A	631	SER	4.0
1	A	602	THR	4.0
1	A	871	GLN	3.9
1	A	867	ASN	3.8
1	A	873	VAL	3.8
1	A	633	PRO	3.7
1	A	860	ILE	3.7
1	A	584	ASN	3.5
1	A	868	GLN	3.4
1	A	861	LEU	3.4
1	A	671	GLY	3.2
1	A	807	ASP	3.1
1	A	856	GLU	2.9
1	A	632	GLY	2.9
1	A	876	ALA	2.9
1	A	630	THR	2.8
1	A	872	GLU	2.8
1	A	870	LYS	2.8
1	A	536	PRO	2.7
1	A	725	ARG	2.6
1	A	875	ASP	2.5
1	A	609	ALA	2.5
1	A	878	GLN	2.5

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	592	VAL	2.2
1	A	842	ILE	2.2
1	A	808	LYS	2.2
1	A	551	GLU	2.2
1	A	546	ARG	2.1
1	A	562	VAL	2.0
1	A	866	TYR	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.