



wwPDB X-ray Structure Validation Summary Report ⓘ

Sep 10, 2023 – 08:07 AM EDT

PDB ID : 4IEF
Title : Complex of Porphyromonas gingivalis RgpB pro- and mature domains
Authors : de Diego, I.; Veillard, F.T.; Guevara, T.; Potempa, B.; Sztukowska, M.;
Potempa, J.; Gomis-Ruth, F.X.
Deposited on : 2012-12-13
Resolution : 2.30 Å (reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.35.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.






Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5042 (2.30-2.30)
Clashscore	141614	5643 (2.30-2.30)
Ramachandran outliers	138981	5575 (2.30-2.30)
Sidechain outliers	138945	5575 (2.30-2.30)
RSRZ outliers	127900	4938 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	210	 4% 82% 9% 9%
1	C	210	 7% 85% 8% 7%
1	E	210	 5% 82% 10% 8%
1	G	210	 9% 84% 6% 10%
2	B	439	 2% 88% 8% .

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	D	439	 4% 88% 8% ••
2	F	439	 5% 87% 9% •
2	H	439	 2% 88% 8% •

2 Entry composition i

There are 10 unique types of molecules in this entry. The entry contains 20046 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Gingipain R2 Pro-Domain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	192	1504	958	250	292	4	0	0	0
1	C	195	1527	972	256	295	4	0	0	0
1	E	194	1516	966	252	294	4	0	0	0
1	G	190	1492	951	248	289	4	0	0	0

There are 20 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	20	GLY	-	expression tag	UNP P95493
A	21	PRO	-	expression tag	UNP P95493
A	22	LEU	-	expression tag	UNP P95493
A	23	GLY	-	expression tag	UNP P95493
A	24	SER	-	expression tag	UNP P95493
C	20	GLY	-	expression tag	UNP P95493
C	21	PRO	-	expression tag	UNP P95493
C	22	LEU	-	expression tag	UNP P95493
C	23	GLY	-	expression tag	UNP P95493
C	24	SER	-	expression tag	UNP P95493
E	20	GLY	-	expression tag	UNP P95493
E	21	PRO	-	expression tag	UNP P95493
E	22	LEU	-	expression tag	UNP P95493
E	23	GLY	-	expression tag	UNP P95493
E	24	SER	-	expression tag	UNP P95493
G	20	GLY	-	expression tag	UNP P95493
G	21	PRO	-	expression tag	UNP P95493
G	22	LEU	-	expression tag	UNP P95493
G	23	GLY	-	expression tag	UNP P95493
G	24	SER	-	expression tag	UNP P95493

- Molecule 2 is a protein called Gingipain R2 Mature Domain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	423	Total 3272	C 2058	N 548	O 649	S 17	0	0	0
2	D	424	Total 3280	C 2062	N 550	O 651	S 17	0	0	0
2	F	424	Total 3280	C 2062	N 550	O 651	S 17	0	0	0
2	H	424	Total 3280	C 2062	N 550	O 651	S 17	0	0	0

There are 24 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	663	HIS	-	expression tag	UNP P95493
B	664	HIS	-	expression tag	UNP P95493
B	665	HIS	-	expression tag	UNP P95493
B	666	HIS	-	expression tag	UNP P95493
B	667	HIS	-	expression tag	UNP P95493
B	668	HIS	-	expression tag	UNP P95493
D	663	HIS	-	expression tag	UNP P95493
D	664	HIS	-	expression tag	UNP P95493
D	665	HIS	-	expression tag	UNP P95493
D	666	HIS	-	expression tag	UNP P95493
D	667	HIS	-	expression tag	UNP P95493
D	668	HIS	-	expression tag	UNP P95493
F	663	HIS	-	expression tag	UNP P95493
F	664	HIS	-	expression tag	UNP P95493
F	665	HIS	-	expression tag	UNP P95493
F	666	HIS	-	expression tag	UNP P95493
F	667	HIS	-	expression tag	UNP P95493
F	668	HIS	-	expression tag	UNP P95493
H	663	HIS	-	expression tag	UNP P95493
H	664	HIS	-	expression tag	UNP P95493
H	665	HIS	-	expression tag	UNP P95493
H	666	HIS	-	expression tag	UNP P95493
H	667	HIS	-	expression tag	UNP P95493
H	668	HIS	-	expression tag	UNP P95493

- Molecule 3 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total Cl 1 1	0	0


- Molecule 4 is BARIUM ION (three-letter code: BA) (formula: Ba).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	B	1	Total Ba 1 1	0	0
4	D	1	Total Ba 1 1	0	0
4	F	1	Total Ba 1 1	0	0
4	H	1	Total Ba 1 1	0	0

- Molecule 5 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	B	4	Total Ca 4 4	0	0
5	D	3	Total Ca 3 3	0	0
5	F	4	Total Ca 4 4	0	0
5	H	3	Total Ca 3 3	0	0

- Molecule 6 is 2-AMINO-2-HYDROXYMETHYL-PROPANE-1,3-DIOL (three-letter code: TRS) (formula: C₄H₁₂NO₃).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
6	B	1	8	4	1	3	0	0

- Molecule 7 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Na		
7	F	3	3	3	0	0

- Molecule 8 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Mg		
8	H	1	1	1	0	0

- Molecule 9 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	H	1	Total	C O	0	0
			6	3 3		


- Molecule 10 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
10	A	68	Total	O	0	0
			68	68		
10	B	179	Total	O	0	0
			179	179		
10	C	62	Total	O	0	0
			62	62		
10	D	139	Total	O	0	0
			139	139		
10	E	69	Total	O	0	0
			69	69		
10	F	156	Total	O	0	0
			156	156		
10	G	43	Total	O	0	0
			43	43		
10	H	142	Total	O	0	0
			142	142		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Gingipain R2 Pro-Domain




- Molecule 1: Gingipain R2 Pro-Domain




- Molecule 1: Gingipain R2 Pro-Domain





- Molecule 1: Gingipain R2 Pro-Domain









● Molecule 2: Gingipain R2 Mature Domain




● Molecule 2: Gingipain R2 Mature Domain



● Molecule 2: Gingipain R2 Mature Domain



● Molecule 2: Gingipain R2 Mature Domain



4 Data and refinement statistics i

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	84.42Å 133.14Å 109.83Å 90.00° 90.52° 90.00°	Depositor
Resolution (Å)	33.67 – 2.30 33.61 – 2.30	Depositor EDS
% Data completeness (in resolution range)	99.2 (33.67-2.30) 99.3 (33.61-2.30)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.68 (at 2.29Å)	Xtrriage
Refinement program	BUSTER 2.11.2	Depositor
R, R_{free}	0.189 , 0.225 0.192 , 0.232	Depositor DCC
R_{free} test set	5323 reflections (4.98%)	wwPDB-VP
Wilson B-factor (Å ²)	32.5	Xtrriage
Anisotropy	0.266	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 47.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	0.027 for h,-k,-l	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	20046	wwPDB-VP
Average B, all atoms (Å ²)	39.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.20% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: TRS, BA, CA, CL, CSD, NA, GOL, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.40	0/1531	0.60	0/2075
1	C	0.40	0/1554	0.60	0/2105
1	E	0.39	0/1543	0.59	0/2091
1	G	0.39	0/1519	0.59	0/2058
2	B	0.41	0/3324	0.62	0/4508
2	D	0.40	0/3332	0.63	0/4519
2	F	0.41	0/3332	0.62	0/4519
2	H	0.40	0/3332	0.62	0/4519
All	All	0.40	0/19467	0.61	0/26394

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1504	0	1513	10	0
1	C	1527	0	1540	9	0
1	E	1516	0	1527	13	0
1	G	1492	0	1501	7	0
2	B	3272	0	3219	18	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	D	3280	0	3225	22	0
2	F	3280	0	3225	26	0
2	H	3280	0	3225	17	0
3	A	1	0	0	0	0
4	B	1	0	0	0	0
4	D	1	0	0	0	0
4	F	1	0	0	0	0
4	H	1	0	0	0	0
5	B	4	0	0	0	0
5	D	3	0	0	0	0
5	F	4	0	0	0	0
5	H	3	0	0	0	0
6	B	8	0	12	0	0
7	F	3	0	0	0	0
8	H	1	0	0	0	0
9	H	6	0	6	0	0
10	A	68	0	0	3	0
10	B	179	0	0	1	0
10	C	62	0	0	1	0
10	D	139	0	0	3	0
10	E	69	0	0	5	0
10	F	156	0	0	3	0
10	G	43	0	0	1	0
10	H	142	0	0	3	0
All	All	20046	0	18993	105	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 105 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:217:ILE:HD11	2:D:535:ILE:HG21	1.59	0.85
1:G:103:GLU:HG3	10:G:319:HOH:O	1.86	0.75
2:F:452:GLY:H	2:F:455:HIS:HD2	1.39	0.69
2:F:537:ARG:HD2	2:F:649:TYR:HB2	1.74	0.68
2:B:452:GLY:H	2:B:455:HIS:HD2	1.40	0.68

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	188/210 (90%)	182 (97%)	6 (3%)	0	100	100
1	C	191/210 (91%)	186 (97%)	5 (3%)	0	100	100
1	E	190/210 (90%)	183 (96%)	7 (4%)	0	100	100
1	G	186/210 (89%)	180 (97%)	6 (3%)	0	100	100
2	B	420/439 (96%)	413 (98%)	5 (1%)	2 (0%)	29	35
2	D	421/439 (96%)	412 (98%)	6 (1%)	3 (1%)	22	26
2	F	421/439 (96%)	410 (97%)	9 (2%)	2 (0%)	29	35
2	H	421/439 (96%)	413 (98%)	5 (1%)	3 (1%)	22	26
All	All	2438/2596 (94%)	2379 (98%)	49 (2%)	10 (0%)	34	42

5 of 10 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	449	SER
2	B	474	VAL
2	D	449	SER
2	F	449	SER
2	F	474	VAL

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	170/184 (92%)	164 (96%)	6 (4%)	36	50

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	C	172/184 (94%)	167 (97%)	5 (3%)	42	58
1	E	171/184 (93%)	164 (96%)	7 (4%)	30	43
1	G	169/184 (92%)	166 (98%)	3 (2%)	59	75
2	B	357/372 (96%)	351 (98%)	6 (2%)	60	76
2	D	358/372 (96%)	351 (98%)	7 (2%)	55	72
2	F	358/372 (96%)	351 (98%)	7 (2%)	55	72
2	H	358/372 (96%)	350 (98%)	8 (2%)	52	69
All	All	2113/2224 (95%)	2064 (98%)	49 (2%)	50	67

5 of 49 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	E	131	ASP
2	F	533	ASN
1	E	205	LEU
2	F	252	ASP
2	F	565	TRP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 22 such sidechains are listed below:

Mol	Chain	Res	Type
2	F	510	ASN
1	G	60	GLN
1	G	55	GLN
1	G	139	GLN
2	D	455	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The

Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	CSD	B	473	2	3,7,8	0.76	0	1,8,10	0.72	0
2	CSD	F	473	2	3,7,8	0.84	0	1,8,10	0.18	0
2	CSD	H	473	2	3,7,8	0.82	0	1,8,10	0.43	0
2	CSD	D	473	2	3,7,8	0.81	0	1,8,10	0.20	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	CSD	B	473	2	-	0/2/6/8	-
2	CSD	F	473	2	-	0/2/6/8	-
2	CSD	H	473	2	-	0/2/6/8	-
2	CSD	D	473	2	-	0/2/6/8	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

3 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	473	CSD	1	0
2	F	473	CSD	1	0
2	H	473	CSD	1	0

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry

Of 25 ligands modelled in this entry, 23 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
9	GOL	H	706	5	5,5,5	0.38	0	5,5,5	0.58	0
6	TRS	B	706	-	7,7,7	0.26	0	9,9,9	1.14	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
9	GOL	H	706	5	-	2/4/4/4	-
6	TRS	B	706	-	-	6/9/9/9	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

5 of 8 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
6	B	706	TRS	C1-C-C2-O2
6	B	706	TRS	C3-C-C2-O2
6	B	706	TRS	N-C-C2-O2
9	H	706	GOL	C1-C2-C3-O3
6	B	706	TRS	C2-C-C1-O1

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	192/210 (91%)	0.09	8 (4%) 36 43	26, 38, 59, 97	0
1	C	195/210 (92%)	0.21	15 (7%) 13 17	23, 42, 68, 103	0
1	E	194/210 (92%)	0.27	10 (5%) 27 34	28, 43, 70, 101	0
1	G	190/210 (90%)	0.43	18 (9%) 8 11	31, 45, 67, 109	0
2	B	422/439 (96%)	-0.02	8 (1%) 66 73	19, 32, 55, 80	0
2	D	423/439 (96%)	0.16	19 (4%) 33 40	19, 34, 64, 87	0
2	F	423/439 (96%)	0.16	20 (4%) 31 38	21, 35, 56, 92	0
2	H	423/439 (96%)	-0.01	10 (2%) 59 66	20, 34, 57, 76	0
All	All	2462/2596 (94%)	0.13	108 (4%) 34 41	19, 37, 63, 109	0

The worst 5 of 108 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	E	196	ALA	7.0
1	A	227	ALA	5.8
2	D	594	SER	5.5
1	G	197	GLU	5.3
2	F	532	PRO	5.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	CSD	D	473	8/9	0.93	0.11	28,30,39,41	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	CSD	B	473	8/9	0.95	0.09	28,29,36,36	0
2	CSD	H	473	8/9	0.95	0.11	29,31,36,38	0
2	CSD	F	473	8/9	0.98	0.08	30,32,36,38	0

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
6	TRS	B	706	8/8	0.64	0.26	54,55,56,56	0
5	CA	B	704	1/1	0.78	0.13	51,51,51,51	0
9	GOL	H	706	6/6	0.91	0.16	39,42,46,49	0
8	MG	H	705	1/1	0.93	0.13	50,50,50,50	0
7	NA	F	708	1/1	0.94	0.34	42,42,42,42	0
7	NA	F	706	1/1	0.94	0.22	38,38,38,38	0
7	NA	F	707	1/1	0.94	0.11	63,63,63,63	0
5	CA	D	704	1/1	0.96	0.12	42,42,42,42	0
5	CA	B	705	1/1	0.97	0.24	44,44,44,44	0
5	CA	F	705	1/1	0.97	0.25	55,55,55,55	0
5	CA	F	704	1/1	0.98	0.18	54,54,54,54	0
5	CA	H	703	1/1	0.98	0.09	31,31,31,31	0
5	CA	B	702	1/1	0.99	0.11	27,27,27,27	0
5	CA	B	703	1/1	0.99	0.10	25,25,25,25	0
5	CA	H	702	1/1	0.99	0.03	35,35,35,35	0
3	CL	A	301	1/1	0.99	0.04	34,34,34,34	0
5	CA	H	704	1/1	0.99	0.15	44,44,44,44	0
4	BA	F	701	1/1	0.99	0.06	59,59,59,59	0
5	CA	D	702	1/1	0.99	0.07	28,28,28,28	0
5	CA	D	703	1/1	0.99	0.12	23,23,23,23	0
4	BA	H	701	1/1	0.99	0.05	67,67,67,67	0
5	CA	F	702	1/1	0.99	0.09	32,32,32,32	0
5	CA	F	703	1/1	0.99	0.05	36,36,36,36	0
4	BA	B	701	1/1	1.00	0.06	55,55,55,55	0
4	BA	D	701	1/1	1.00	0.05	63,63,63,63	0

6.5 Other polymers

There are no such residues in this entry.