



# Full wwPDB X-ray Structure Validation Report ⓘ

Jan 3, 2024 – 02:13 pm GMT

PDB ID : 5IFP  
Title : STRUCTURE OF BETA-GALACTOSIDASE FROM ASPERGILLUS NIGER  
Authors : Rico-Diaz, A.; Ramirez-Escudero, M.; Vizoso Vazquez, A.; Cerdan, M.E.; Becerra, M.; Sanz-Aparicio, J.  
Deposited on : 2016-02-26  
Resolution : 1.71 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.4, CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.36  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.71 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	5722 (1.74-1.70)
Clashscore	141614	6152 (1.74-1.70)
Ramachandran outliers	138981	6051 (1.74-1.70)
Sidechain outliers	138945	6051 (1.74-1.70)
RSRZ outliers	127900	5629 (1.74-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	1013	 93%
2	B	8	 12% 88%
2	C	8	 75% 25%
3	D	2	 50% 50%
4	E	6	 33% 50% 17%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

<b>Mol</b>	<b>Type</b>	<b>Chain</b>	<b>Res</b>	<b>Chirality</b>	<b>Geometry</b>	<b>Clashes</b>	<b>Electron density</b>
2	MAN	B	6	-	-	-	X
2	MAN	C	6	-	-	-	X
3	NAG	D	2	-	-	-	X

## 2 Entry composition i

There are 10 unique types of molecules in this entry. The entry contains 8920 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called beta-galactosidase A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	968	7482	4780	1219	1472	11	0	0	0

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1008	HIS	-	expression tag	UNP A2QAN3
A	1009	HIS	-	expression tag	UNP A2QAN3
A	1010	HIS	-	expression tag	UNP A2QAN3
A	1011	HIS	-	expression tag	UNP A2QAN3
A	1012	HIS	-	expression tag	UNP A2QAN3
A	1013	HIS	-	expression tag	UNP A2QAN3

- Molecule 2 is an oligosaccharide called alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-3)-alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	B	8	94	52	2	40	0	0	0
2	C	8	94	52	2	40	0	0	0

- Molecule 3 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
3	D	2	28	16	2	10	0	0	0

- Molecule 4 is an oligosaccharide called alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-3)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
4	E	6	72	40	2	30	0	0	0

- Molecule 5 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula:  $C_8H_{15}NO_6$ ).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
5	A	1	14	8	1	5	0	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	1	Total	C	N	O	0	0
			14	8	1	5		
5	A	1	Total	C	N	O	0	0
			14	8	1	5		
5	A	1	Total	C	N	O	0	0
			14	8	1	5		
5	A	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 6 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).




Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	1	Total	O S	0	0
			5	4 1		
6	A	1	Total	O S	0	0
			5	4 1		

- Molecule 7 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	1	Total Cl	0	0
			1 1		


- Molecule 8 is 2-[BIS-(2-HYDROXY-ETHYL)-AMINO]-2-HYDROXYMETHYL-PROPAN

E-1,3-DIOL (three-letter code: BTB) (formula:  $C_8H_{19}NO_5$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
8	A	1	14	8	1	5	0	0

- Molecule 9 is GLYCEROL (three-letter code: GOL) (formula:  $C_3H_8O_3$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
9	A	1	6	3	3	0	0
9	A	1	6	3	3	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
9	A	1	Total	C	O	0	0
			6	3	3		
9	A	1	Total	C	O	0	0
			6	3	3		
9	A	1	Total	C	O	0	0
			6	3	3		

- Molecule 10 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
10	A	1011	Total	O	0	0
			1011	1011		





- Molecule 4: alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-3)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain E:  33% 50% 17%



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	85.57Å 111.42Å 126.73Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	83.68 – 1.71 39.94 – 1.71	Depositor EDS
% Data completeness (in resolution range)	99.9 (83.68-1.71) 99.9 (39.94-1.71)	Depositor EDS
$R_{merge}$	0.12	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.72 (at 1.71Å)	Xtrriage
Refinement program	REFMAC 5.8.0135	Depositor
R, $R_{free}$	0.150 , 0.171 0.162 , 0.181	Depositor DCC
$R_{free}$ test set	6555 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	11.4	Xtrriage
Anisotropy	0.290	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.38 , 39.2	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	8920	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	13.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.71% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: BTB, CL, MAN, SO4, GOL, NAG, BMA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.34	0/7685	0.62	0/10488

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	7482	0	7177	12	0
2	B	94	0	79	0	0
2	C	94	0	79	0	0
3	D	28	0	25	1	0
4	E	72	0	61	1	0
5	A	84	0	78	0	0
6	A	10	0	0	0	0
7	A	1	0	0	0	0
8	A	14	0	19	0	0
9	A	30	0	40	0	0
10	A	1011	0	0	1	0
All	All	8920	0	7558	12	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 1.

All (12) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:260:TYR:HA	1:A:300:GLN:HB2	1.94	0.49
1:A:791:TYR:CE2	3:D:1:NAG:H82	2.51	0.46
1:A:728:LEU:HD21	10:A:3126:HOH:O	2.17	0.44
1:A:355:TYR:CZ	1:A:357:ASN:HB2	2.52	0.44
1:A:148:PHE:CG	1:A:152:LEU:HD12	2.54	0.43
1:A:96:TYR:CE2	1:A:141:ALA:HB2	2.53	0.43
1:A:51:LYS:NZ	1:A:335:GLN:HE22	2.17	0.42
1:A:632:VAL:HG11	1:A:656:TRP:CE2	2.55	0.41
1:A:355:TYR:HB2	1:A:356:PRO:HD2	2.02	0.41
1:A:390:LYS:NZ	1:A:577:GLN:HE21	2.19	0.41
1:A:940:GLY:HA2	4:E:1:NAG:H83	2.03	0.41
1:A:416:VAL:HG22	1:A:430:VAL:HG22	2.03	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	966/1013 (95%)	935 (97%)	30 (3%)	1 (0%)	51 33

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	422	ASN

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	808/842 (96%)	803 (99%)	5 (1%)	<a href="#">86</a> <a href="#">80</a>

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	63	PHE
1	A	198	GLU
1	A	355	TYR
1	A	553	ARG
1	A	931	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	335	GLN
1	A	349	ASN
1	A	422	ASN
1	A	458	GLN
1	A	466	ASN
1	A	577	GLN
1	A	638	ASN
1	A	648	HIS
1	A	781	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates i

24 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	NAG	B	1	1,2	14,14,15	0.38	0	17,19,21	1.00	1 (5%)
2	NAG	B	2	2	14,14,15	0.29	0	17,19,21	0.76	0
2	BMA	B	3	2	11,11,12	0.33	0	15,15,17	0.78	1 (6%)
2	MAN	B	4	2	11,11,12	0.19	0	15,15,17	1.02	1 (6%)
2	MAN	B	5	2	11,11,12	0.38	0	15,15,17	1.00	1 (6%)
2	MAN	B	6	2	11,11,12	0.43	0	15,15,17	0.94	1 (6%)
2	MAN	B	7	2	11,11,12	0.45	0	15,15,17	0.94	2 (13%)
2	MAN	B	8	2	11,11,12	0.32	0	15,15,17	0.71	1 (6%)
2	NAG	C	1	1,2	14,14,15	0.32	0	17,19,21	1.01	0
2	NAG	C	2	2	14,14,15	0.23	0	17,19,21	0.69	0
2	BMA	C	3	2	11,11,12	0.40	0	15,15,17	0.68	0
2	MAN	C	4	2	11,11,12	0.36	0	15,15,17	1.10	1 (6%)
2	MAN	C	5	2	11,11,12	0.36	0	15,15,17	0.77	0
2	MAN	C	6	2	11,11,12	0.40	0	15,15,17	0.79	1 (6%)
2	MAN	C	7	2	11,11,12	0.30	0	15,15,17	0.72	0
2	MAN	C	8	2	11,11,12	0.36	0	15,15,17	0.80	0
3	NAG	D	1	3,1	14,14,15	0.38	0	17,19,21	0.73	0
3	NAG	D	2	3	14,14,15	0.30	0	17,19,21	0.72	0
4	NAG	E	1	4,1	14,14,15	0.31	0	17,19,21	0.80	1 (5%)
4	NAG	E	2	4	14,14,15	0.30	0	17,19,21	0.78	0
4	BMA	E	3	4	11,11,12	0.47	0	15,15,17	1.01	1 (6%)
4	MAN	E	4	4	11,11,12	0.38	0	15,15,17	0.86	1 (6%)
4	MAN	E	5	4	11,11,12	0.33	0	15,15,17	0.92	1 (6%)
4	MAN	E	6	4	11,11,12	0.36	0	15,15,17	0.70	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	B	1	1,2	-	0/6/23/26	0/1/1/1
2	NAG	B	2	2	-	0/6/23/26	0/1/1/1
2	BMA	B	3	2	-	0/2/19/22	0/1/1/1
2	MAN	B	4	2	-	0/2/19/22	0/1/1/1
2	MAN	B	5	2	-	2/2/19/22	0/1/1/1
2	MAN	B	6	2	-	2/2/19/22	0/1/1/1
2	MAN	B	7	2	-	0/2/19/22	0/1/1/1
2	MAN	B	8	2	-	0/2/19/22	0/1/1/1
2	NAG	C	1	1,2	-	0/6/23/26	0/1/1/1
2	NAG	C	2	2	-	0/6/23/26	0/1/1/1
2	BMA	C	3	2	-	0/2/19/22	0/1/1/1
2	MAN	C	4	2	-	0/2/19/22	0/1/1/1
2	MAN	C	5	2	-	0/2/19/22	0/1/1/1
2	MAN	C	6	2	-	2/2/19/22	0/1/1/1
2	MAN	C	7	2	-	0/2/19/22	0/1/1/1
2	MAN	C	8	2	-	0/2/19/22	0/1/1/1
3	NAG	D	1	3,1	-	2/6/23/26	0/1/1/1
3	NAG	D	2	3	-	0/6/23/26	0/1/1/1
4	NAG	E	1	4,1	-	0/6/23/26	0/1/1/1
4	NAG	E	2	4	-	0/6/23/26	0/1/1/1
4	BMA	E	3	4	-	2/2/19/22	0/1/1/1
4	MAN	E	4	4	-	0/2/19/22	0/1/1/1
4	MAN	E	5	4	-	0/2/19/22	0/1/1/1
4	MAN	E	6	4	-	0/2/19/22	0/1/1/1

There are no bond length outliers.

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	4	MAN	C1-O5-C5	2.84	116.04	112.19
2	B	4	MAN	C1-O5-C5	2.81	115.99	112.19
2	B	5	MAN	C1-O5-C5	2.78	115.95	112.19
4	E	3	BMA	C1-C2-C3	2.25	112.44	109.67
2	B	3	BMA	C1-O5-C5	2.23	115.22	112.19
4	E	1	NAG	C1-C2-N2	-2.21	106.72	110.49
4	E	5	MAN	C1-O5-C5	2.17	115.13	112.19
4	E	4	MAN	C1-O5-C5	2.17	115.13	112.19
2	B	8	MAN	C1-O5-C5	2.15	115.11	112.19
2	B	7	MAN	C1-O5-C5	2.03	114.95	112.19
2	C	6	MAN	C1-O5-C5	2.03	114.95	112.19
2	B	6	MAN	C1-C2-C3	2.02	112.15	109.67
2	B	7	MAN	C1-C2-C3	2.02	112.15	109.67

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	1	NAG	O5-C5-C6	2.02	110.37	107.20

There are no chirality outliers.

All (10) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	C	6	MAN	C4-C5-C6-O6
3	D	1	NAG	O5-C5-C6-O6
2	C	6	MAN	O5-C5-C6-O6
4	E	3	BMA	C4-C5-C6-O6
2	B	5	MAN	C4-C5-C6-O6
2	B	5	MAN	O5-C5-C6-O6
2	B	6	MAN	C4-C5-C6-O6
4	E	3	BMA	O5-C5-C6-O6
3	D	1	NAG	C4-C5-C6-O6
2	B	6	MAN	O5-C5-C6-O6


There are no ring outliers.


2 monomers are involved in 2 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	D	1	NAG	1	0
4	E	1	NAG	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.









## 5.6 Ligand geometry [i](#)

Of 15 ligands modelled in this entry, 1 is monoatomic - leaving 14 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
5	NAG	A	1156	1	14,14,15	0.30	0	17,19,21	0.77	0
9	GOL	A	3005	-	5,5,5	0.55	0	5,5,5	0.89	0
6	SO4	A	2002	-	4,4,4	0.33	0	6,6,6	0.24	0
5	NAG	A	1402	1	14,14,15	0.32	0	17,19,21	0.80	0
5	NAG	A	1522	1	14,14,15	0.30	0	17,19,21	1.07	1 (5%)
9	GOL	A	3003	-	5,5,5	0.36	0	5,5,5	0.26	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
6	SO4	A	2001	-	4,4,4	0.31	0	6,6,6	0.14	0
8	BTB	A	2004	-	13,13,13	0.80	0	7,16,16	0.46	0
5	NAG	A	1478	1	14,14,15	0.34	0	17,19,21	1.08	1 (5%)
9	GOL	A	3001	-	5,5,5	0.31	0	5,5,5	0.23	0
9	GOL	A	3002	-	5,5,5	0.31	0	5,5,5	0.33	0
9	GOL	A	3006	-	5,5,5	0.26	0	5,5,5	0.28	0
5	NAG	A	1777	1	14,14,15	0.29	0	17,19,21	0.61	0
5	NAG	A	1739	1	14,14,15	0.64	0	17,19,21	1.12	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	NAG	A	1156	1	-	0/6/23/26	0/1/1/1
9	GOL	A	3005	-	-	0/4/4/4	-
5	NAG	A	1402	1	-	0/6/23/26	0/1/1/1
5	NAG	A	1522	1	-	0/6/23/26	0/1/1/1
9	GOL	A	3003	-	-	2/4/4/4	-
8	BTB	A	2004	-	-	0/21/21/21	-
5	NAG	A	1478	1	-	2/6/23/26	0/1/1/1
9	GOL	A	3001	-	-	0/4/4/4	-
9	GOL	A	3002	-	-	0/4/4/4	-
9	GOL	A	3006	-	-	2/4/4/4	-
5	NAG	A	1777	1	-	0/6/23/26	0/1/1/1
5	NAG	A	1739	1	-	0/6/23/26	0/1/1/1

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	1478	NAG	O5-C5-C6	3.28	112.35	107.20
5	A	1522	NAG	C3-C4-C5	2.30	114.34	110.24
5	A	1739	NAG	C3-C4-C5	2.20	114.17	110.24

There are no chirality outliers.

All (6) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
9	A	3003	GOL	O1-C1-C2-C3
9	A	3006	GOL	C1-C2-C3-O3
5	A	1478	NAG	O5-C5-C6-O6
5	A	1478	NAG	C4-C5-C6-O6
9	A	3003	GOL	O1-C1-C2-O2
9	A	3006	GOL	O2-C2-C3-O3

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	968/1013 (95%)	-0.11	11 (1%) 80 84	6, 11, 20, 33	0

All (11) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	41	LEU	8.0
1	A	581	ASP	5.2
1	A	422	ASN	5.1
1	A	916	THR	3.7
1	A	1008	HIS	3.7
1	A	873	GLN	3.0
1	A	499	ASP	2.9
1	A	423	SER	2.9
1	A	529	SER	2.7
1	A	702	GLN	2.2
1	A	688	SER	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	NAG	D	2	14/15	0.46	0.46	39,42,44,44	0

*Continued on next page...*






Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	MAN	C	6	11/12	0.59	0.42	35,39,40,41	0
2	MAN	B	6	11/12	0.66	0.41	42,44,45,46	0
4	MAN	E	6	11/12	0.72	0.38	37,40,41,42	0
2	MAN	B	5	11/12	0.78	0.25	33,38,39,40	0
2	MAN	B	7	11/12	0.80	0.36	25,27,30,31	0
2	MAN	C	8	11/12	0.81	0.17	24,25,26,27	0
2	MAN	B	4	11/12	0.83	0.12	22,23,27,28	0
4	BMA	E	3	11/12	0.84	0.20	22,24,27,28	0
3	NAG	D	1	14/15	0.84	0.19	22,27,29,35	0
2	MAN	B	8	11/12	0.86	0.25	27,28,29,29	0
2	MAN	C	4	11/12	0.88	0.17	19,20,22,23	0
4	MAN	E	5	11/12	0.89	0.17	27,28,30,33	0
2	MAN	C	5	11/12	0.89	0.20	24,26,27,30	0
2	BMA	B	3	11/12	0.90	0.10	14,17,19,22	0
2	MAN	C	7	11/12	0.90	0.11	19,20,21,22	0
2	BMA	C	3	11/12	0.91	0.10	14,16,17,17	0
4	MAN	E	4	11/12	0.91	0.10	18,21,22,24	0
4	NAG	E	1	14/15	0.93	0.08	12,13,15,16	0
4	NAG	E	2	14/15	0.93	0.13	15,18,20,20	0
2	NAG	C	2	14/15	0.93	0.09	11,11,13,14	0
2	NAG	C	1	14/15	0.95	0.08	10,10,12,12	0
2	NAG	B	1	14/15	0.96	0.08	9,10,12,12	0
2	NAG	B	2	14/15	0.97	0.07	11,11,12,13	0




The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.

**Electron density around Chain B:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)




**Electron density around Chain C:**


$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



**Electron density around Chain D:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
5	NAG	A	1402	14/15	0.66	0.32	33,37,41,43	0
5	NAG	A	1522	14/15	0.70	0.22	28,31,34,36	0
9	GOL	A	3003	6/6	0.73	0.21	40,40,41,41	0
5	NAG	A	1156	14/15	0.75	0.31	23,25,29,29	0
9	GOL	A	3002	6/6	0.76	0.18	31,31,31,31	0
5	NAG	A	1739	14/15	0.77	0.16	17,19,20,20	0
8	BTB	A	2004	14/14	0.78	0.23	25,29,32,33	0
9	GOL	A	3006	6/6	0.79	0.28	34,36,37,37	0
5	NAG	A	1478	14/15	0.84	0.23	24,28,30,31	0
9	GOL	A	3005	6/6	0.85	0.22	13,15,15,15	0
5	NAG	A	1777	14/15	0.88	0.12	16,18,20,21	0
6	SO4	A	2002	5/5	0.89	0.23	35,37,39,39	0
9	GOL	A	3001	6/6	0.97	0.08	8,8,8,8	0
6	SO4	A	2001	5/5	0.98	0.11	22,23,23,24	0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
7	CL	A	2003	1/1	0.99	0.04	12,12,12,12	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.