

Full wwPDB NMR Structure Validation Report (i)

May 29, 2020 – 07:10 am BST

PDB ID : 5IIV

Title : GCN4p pH 1.5

Authors : Brady, M.R.; Kaplan, A.R.; Alexandrescu, A.T.

Deposited on : 2016-03-01

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at

https://www.wwpdb.org/validation/2017/NMRValidationReportHelp with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (1)) were used in the production of this report:

Cyrange : Kirchner and Güntert (2011)

NmrClust : Kelley et al. (1996)

MolProbity: 4.02b-467

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

RCI : v 1n 11 5 13 A (Berjanski et al., 2005)

PANAV : Wang et al. (2010)

ShiftChecker: 2.11

Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.11

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $SOLUTION\ NMR$

The overall completeness of chemical shifts assignment is 35%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$\begin{array}{c} \text{Whole archive} \\ (\#\text{Entries}) \end{array}$	$rac{ ext{NMR archive}}{ ext{(\#Entries)}}$
Clashscore	158937	12864
Ramachandran outliers	154571	11451
Sidechain outliers	154315	11428

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Mol	Chain	Length	Quality of chain			
1	A	33	70%	9%	21%	
1	В	33	70%	9%	21%	



2 Ensemble composition and analysis (i)

This entry contains 20 models. Model 2 is the overall representative, medoid model (most similar to other models).

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues						
Well-defined core	Well-defined core Residue range (total) Backbone RMSD (Å) Medoid model					
1	A:7-A:32, B:7-B:32 (52)	0.36	2			

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 3 clusters and 4 single-model clusters were found.

Cluster number	Models
1	1, 2, 4, 5, 10, 11, 12, 13, 14, 15, 19
2	8, 18, 20
3	6, 17
Single-model clusters	3; 7; 9; 16



3 Entry composition (i)

There is only 1 type of molecule in this entry. The entry contains 1096 atoms, of which 560 are hydrogens and 0 are deuteriums.

• Molecule 1 is a protein called General control protein GCN4.

Mol	Chain	Residues		${f Atoms}$					Trace
1	Α	33	Total	С	Н	N	О	S	0
1	$\begin{array}{c c} 1 & A & \end{array}$	აა	548	167	280	46	54	1	
1	D	33	Total	С	Н	N	О	S	0
	$\begin{array}{c c} I & B \end{array}$	_ აა	548	167	280	46	54	1	U

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	${f Comment}$	Reference
A	1	GLY	_	expression tag	UNP P03069
A	2	SER	_	expression tag	UNP P03069
В	1	GLY	_	expression tag	UNP P03069
В	2	SER	-	expression tag	UNP P03069




4 Residue-property plots (i)

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA and DNA chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

• Molecule 1: General control protein GCN4




4.2 Scores per residue for each member of the ensemble

Colouring as in section 4.1 above.

4.2.1 Score per residue for model 1

• Molecule 1: General control protein GCN4





9%

21%

4.2.2 Score per residue for model 2 (medoid)

• Molecule 1: General control protein GCN4

70%

61 82 83 83 84 84 16 16 16 17 127

Chain B:

4.2.3 Score per residue for model 3

• Molecule 1: General control protein GCN4

Chain A: 55% 21% • 21% • 21%

• Molecule 1: General control protein GCN4

Chain B: 55% 21% · 21%

4.2.4 Score per residue for model 4

• Molecule 1: General control protein GCN4

Chain A: 52% 24% · 21%

• Molecule 1: General control protein GCN4

Chain B: 52% 24% · 21%





4.2.5 Score per residue for model 5

• Molecule 1: General control protein GCN4

Chain A: 58% 21% 21%

• Molecule 1: General control protein GCN4

Chain B: 58% 21% 21%



4.2.6 Score per residue for model 6

• Molecule 1: General control protein GCN4

Chain A: 58% 18% • 21%

• Molecule 1: General control protein GCN4

Chain B: 58% 18% • 21%



4.2.7 Score per residue for model 7

• Molecule 1: General control protein GCN4

Chain A: 55% 21% · 21%

• Molecule 1: General control protein GCN4

Chain B: 55% 21% · 21%





4.2.8 Score per residue for model 8

• Molecule 1: General control protein GCN4

Chain A: 64% 15% 21%

• Molecule 1: General control protein GCN4

Chain B: 58% 21% 21%

8.82 6.64 6.65

4.2.9 Score per residue for model 9

• Molecule 1: General control protein GCN4

Chain A: 61% 18% 21%

• Molecule 1: General control protein GCN4

Chain B: 61% 18% 21%

4.2.10 Score per residue for model 10

• Molecule 1: General control protein GCN4

Chain A: 61% 18% 21%

• Molecule 1: General control protein GCN4

Chain B: 61% 18% 21%



4.2.11 Score per residue for model 11

• Molecule 1: General control protein GCN4

Chain A: 67% 9% · 21%



• Molecule 1: General control protein GCN4

Chain B: 67% 9% · 21%



4.2.12 Score per residue for model 12

• Molecule 1: General control protein GCN4

Chain A: 64% 15% 21%



• Molecule 1: General control protein GCN4

Chain B: 64% 15% 21%



4.2.13 Score per residue for model 13

• Molecule 1: General control protein GCN4

Chain A: 70% 9% 21%



• Molecule 1: General control protein GCN4

Chain B: 70% 9% 21%





4.2.14 Score per residue for model 14

• Molecule 1: General control protein GCN4

Chain A: 64% 12% • 21%

61 M3 M3 M4 M4 M4 M7 M17 V18 H19 H19

• Molecule 1: General control protein GCN4

Chain B: 64% 12% • 21%

4.2.15 Score per residue for model 15

• Molecule 1: General control protein GCN4

Chain A: 76% . 21%

K4 K4 K4 K6 L6 K9 K9

• Molecule 1: General control protein GCN4

Chain B: 76% . 21%

4.2.16 Score per residue for model 16

• Molecule 1: General control protein GCN4

Chain A: 61% 18% 21%

• Molecule 1: General control protein GCN4


Chain B: 61% 18% 21%






4.2.17 Score per residue for model 17

• Molecule 1: General control protein GCN4




4.2.18 Score per residue for model 18

• Molecule 1: General control protein GCN4



4.2.19 Score per residue for model 19

• Molecule 1: General control protein GCN4






4.2.20 Score per residue for model 20

• Molecule 1: General control protein GCN4

Chain A: 73% 6% 21%

• Molecule 1: General control protein GCN4

Chain B: 73% 6% 21%





Refinement protocol and experimental data overview (i) 5



Of the? calculated structures, 20 were deposited, based on the following criterion:?.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
X-PLOR NIH	structure calculation	
ARIA	refinement	

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	$input_cs.cif$
Number of chemical shift lists	1
Total number of shifts	333
Number of shifts mapped to atoms	333
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	35%

No validations of the models with respect to experimental NMR restraints is performed at this time.



6 Model quality (i)

6.1 Standard geometry (i)

There are no covalent bond-length or bond-angle outliers.

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

There are no planarity outliers.

6.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	214	223	222	3±3
1	В	214	223	222	4±3
All	All	8560	8920	8880	109

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	$\operatorname{Clash}(\mathring{\mathrm{A}})$	$\mathbf{Distance}(\mathbf{\mathring{A}})$	Models	
Atom-1	A tom-2	Clash(A)	Distance(A)	Worst	Total
1:A:24:VAL:HG12	1:B:24:VAL:HG12	0.69	1.63	7	1
1:A:14:LEU:HA	1:A:17:ASN:ND2	0.62	2.09	12	1
1:B:14:LEU:HA	1:B:17:ASN:ND2	0.62	2.10	12	1
1:A:14:LEU:HA	1:A:17:ASN:OD1	0.59	1.97	4	2
1:B:14:LEU:HA	1:B:17:ASN:OD1	0.59	1.97	4	2
1:A:24:VAL:HG22	1:B:24:VAL:HG22	0.57	1.77	9	3
1:B:23:GLU:O	1:B:26:ARG:HG2	0.56	1.99	17	5
1:A:23:GLU:O	1:A:26:ARG:HG2	0.56	1.99	17	5
1:A:27:LEU:HB3	1:B:27:LEU:HB3	0.56	1.76	11	13
1:A:20:LEU:HD11	1:B:20:LEU:HD11	0.52	1.79	17	1
1:A:17:ASN:HD22	1:A:18:TYR:N	0.52	2.03	6	2
1:B:25:ALA:O	1:B:29:LYS:HG2	0.51	2.06	7	1
1:B:17:ASN:HD22	1:B:18:TYR:N	0.51	2.04	6	3

Continued on next page...



Continued from previous page...

Atom 1	A tom 2	Clash(Å)	Distance (Å)	Models	
Atom-1	Atom-2	Clash(A)	$\operatorname{Distance}(\operatorname{\AA})$	Worst	Total
1:A:25:ALA:O	1:A:29:LYS:HG2	0.51	2.06	7	1
1:A:27:LEU:O	1:A:31:VAL:HG23	0.50	2.07	19	4
1:A:26:ARG:O	1:A:30:LEU:HD23	0.49	2.07	11	1
1:B:26:ARG:O	1:B:30:LEU:HD23	0.49	2.07	11	1
1:B:27:LEU:O	1:B:31:VAL:HG23	0.49	2.08	19	4
1:B:19:HIS:O	1:B:23:GLU:HG2	0.49	2.08	14	3
1:A:19:HIS:O	1:A:23:GLU:HG2	0.48	2.08	14	3
1:A:24:VAL:HG12	1:B:24:VAL:CG1	0.48	2.36	7	1
1:B:26:ARG:O	1:B:30:LEU:HD13	0.47	2.10	13	3
1:A:24:VAL:CG1	1:B:24:VAL:HG12	0.47	2.37	7	1
1:A:26:ARG:O	1:A:30:LEU:HD13	0.46	2.10	13	3
1:B:20:LEU:O	1:B:24:VAL:HG23	0.46	2.11	8	2
1:A:20:LEU:O	1:A:24:VAL:HG23	0.45	2.11	8	2
1:B:20:LEU:HD13	1:B:21:GLU:N	0.45	2.25	17	1
1:A:26:ARG:O	1:A:30:LEU:HG	0.45	2.12	10	2
1:A:20:LEU:HD13	1:A:21:GLU:N	0.45	2.25	17	1
1:B:26:ARG:O	1:B:30:LEU:HG	0.45	2.12	10	2
1:B:12:GLU:O	1:B:15:SER:HB3	0.45	2.12	18	1
1:A:17:ASN:HA	1:A:20:LEU:HB3	0.45	1.89	17	1
1:A:20:LEU:HD13	1:A:20:LEU:C	0.44	2.33	17	1
1:B:20:LEU:HD13	1:B:20:LEU:C	0.44	2.33	17	1
1:B:17:ASN:HA	1:B:20:LEU:HB3	0.44	1.90	17	1
1:A:12:GLU:O	1:A:15:SER:HB3	0.44	2.13	18	1
1:B:17:ASN:O	1:B:21:GLU:HG3	0.43	2.13	17	1
1:A:10:VAL:O	1:A:14:LEU:HD23	0.43	2.14	17	1
1:A:20:LEU:HD11	1:B:20:LEU:CD1	0.43	2.44	17	1
1:A:25:ALA:O	1:A:29:LYS:HE3	0.43	2.13	3	1
1:B:25:ALA:O	1:B:29:LYS:HE3	0.43	2.12	3	1
1:B:15:SER:O	1:B:19:HIS:HB2	0.43	2.13	7	1
1:A:17:ASN:O	1:A:21:GLU:HG3	0.43	2.13	17	1
1:A:20:LEU:CD1	1:B:20:LEU:HD11	0.43	2.44	17	1
1:B:13:LEU:O	1:B:16:LYS:HG2	0.42	2.14	16	1
1:B:10:VAL:O	1:B:14:LEU:HD23	0.42	2.14	17	1
1:B:24:VAL:O	1:B:28:LYS:HG2	0.42	2.14	1	1
1:A:13:LEU:O	1:A:16:LYS:HG2	0.42	2.14	16	1
1:A:24:VAL:O	1:A:28:LYS:HG2	0.42	2.15	1	1
1:B:23:GLU:HA	1:B:26:ARG:HG3	0.42	1.92	9	1
1:A:10:VAL:HG22	1:B:10:VAL:HG22	0.42	1.90	4	1
1:B:23:GLU:O	1:B:27:LEU:HD13	0.42	2.15	5	1
1:A:27:LEU:HD23	1:B:28:LYS:N	0.42	2.29	9	1
1:A:20:LEU:HD11	1:B:20:LEU:CG	0.42	2.45	17	1

Continued on next page...



Continued from previous page...

Atom-1	Atom-2	$\operatorname{Clash}(\mathring{\mathrm{A}})$	$\mathbf{Distance}(\mathbf{\mathring{A}})$	Models	
Atom-1	A tom-2	Clash(A)	Distance(A)	Worst	Total
1:A:15:SER:O	1:A:19:HIS:HB2	0.42	2.14	7	1
1:A:23:GLU:HA	1:A:26:ARG:HG3	0.42	1.92	9	1
1:A:20:LEU:CG	1:B:20:LEU:HD11	0.41	2.45	17	1
1:A:23:GLU:O	1:A:27:LEU:HD13	0.41	2.15	5	1
1:B:12:GLU:O	1:B:16:LYS:HG3	0.41	2.16	8	1
1:A:20:LEU:HD11	1:B:20:LEU:HD21	0.41	1.93	17	1
1:A:19:HIS:O	1:A:22:ASN:HB2	0.40	2.16	4	1
1:B:19:HIS:O	1:B:22:ASN:HB2	0.40	2.16	4	1
1:A:27:LEU:HB2	1:B:27:LEU:HD13	0.40	1.91	6	1
1:A:20:LEU:HD21	1:B:20:LEU:HD11	0.40	1.94	17	1

6.3 Torsion angles (i)

6.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Pe	erce	${f ntiles}$
1	A	26/33~(79%)	$25\pm0 \ (98\pm2\%)$	0±0 (1±1%)	0±0 (2±2%)		14	59
1	В	26/33~(79%)	$25\pm0 \ (98\pm2\%)$	0±0 (1±1%)	0±0 (2±2%)		14	59
All	All	1040/1320 (79%)	1018 (98%)	6 (1%)	16 (2%)		14	59

All 2 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	32	GLY	8
1	В	32	GLY	8

6.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.



Mol	Chain	Analysed	Rotameric	Outliers	Percen	$_{ m tiles}$
1	A	$24/30 \ (80\%)$	$23\pm1 \ (95\pm5\%)$	1±1 (5±5%)	29	78
1	В	24/30 (80%)	$23\pm1 \ (95\pm5\%)$	1±1 (5±5%)	29	78
All	All	$960/1200 \; (80\%)$	914 (95%)	46 (5%)	29	78

All 24 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	8	ASP	4
1	В	8	ASP	4
1	В	17	ASN	3
1	A	22	ASN	3
1	В	22	ASN	3
1	A	17	ASN	3
1	В	23	GLU	2
1	A	20	LEU	2
1	A	7	GLU	2
1	В	20	LEU	2
1	A	23	GLU	2
1	A	26	ARG	2
1	В	26	ARG	2
1	В	7	GLU	2
1	В	9	LYS	1
1	A	14	LEU	1
1	A	9	LYS	1
1	A	24	VAL	1
1	A	13	LEU	1
1	В	14	LEU	1
1	В	13	LEU	1
1	В	29	LYS	1
1	В	24	VAL	1
1	A	29	LYS	1

6.3.3 RNA (i)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.



6.5 Carbohydrates (i)

There are no carbohydrates in this entry.

6.6 Ligand geometry (i)

There are no ligands in this entry.

6.7 Other polymers (i)

There are no such molecules in this entry.

6.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



7 Chemical shift validation (i)

The completeness of assignment taking into account all chemical shift lists is 35% for the well-defined parts and 35% for the entire structure.

7.1 Chemical shift list 1

File name: input cs.cif

Chemical shift list name: 15_cshift.txt

7.1.1 Bookkeeping (i)

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	333
Number of shifts mapped to atoms	333
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	0

7.1.2 Chemical shift referencing (i)

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	$\text{Correction} \pm \text{precision}, \textit{ppm}$	Suggested action
$^{13}\mathrm{C}_{\alpha}$	32	-1.10 ± 0.58	None needed (imprecise)
$^{13}C_{\beta}$	31	0.62 ± 0.15	Should be applied
¹³ C′	31	-0.88 ± 0.40	Should be applied
^{15}N	31	0.53 ± 0.36	None needed (imprecise)

7.1.3 Completeness of resonance assignments (i)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 35%, i.e. 241 atoms were assigned a chemical shift out of a possible 686. 0 out of 16 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	$^{1}\mathbf{H}$	$^{13}\mathbf{C}$	$^{15}{ m N}$
Backbone	128/260~(49%)	51/104 (49%)	51/104 (49%)	26/52~(50%)
Sidechain	113/394 (29%)	77/228 (34%)	34/148 (23%)	2/18 (11%)

Continued on next page...



Continued from previous page...

	Total	$^{1}\mathrm{H}$	$^{13}\mathbf{C}$	$^{15}{ m N}$
Aromatic	0/32~(0%)	0/16 (0%)	0/12 (0%)	0/4~(0%)
Overall	241/686 (35%)	$128/348 \ (37\%)$	85/264 (32%)	$28/74 \ (38\%)$

The following table shows the completeness of the chemical shift assignments for the full structure. The overall completeness is 35%, i.e. 300 atoms were assigned a chemical shift out of a possible 856. 0 out of 18 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	$^{1}\mathrm{H}$	$^{13}\mathbf{C}$	$^{15}{ m N}$
Backbone	$156/330 \ (47\%)$	62/132~(47%)	63/132~(48%)	31/66 (47%)
Sidechain	$144/494 \ (29\%)$	$99/288 \; (34\%)$	42/184~(23%)	3/22 (14%)
Aromatic	$0/32 \; (0\%)$	$0/16 \; (0\%)$	$0/12 \; (0\%)$	0/4 (0%)
Overall	300/856 $(35%)$	161/436 (37%)	$105/328 \ (32\%)$	34/92 (37%)

7.1.4 Statistically unusual chemical shifts (i)

There are no statistically unusual chemical shifts.

7.1.5 Random Coil Index (RCI) plots (i)

The image below reports random coil index values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition.

Random coil index (RCI) for chain A:

