



Full wwPDB X-ray Structure Validation Report ⓘ

May 22, 2020 – 11:00 pm BST

PDB ID : 4IK5
Title : High resolution structure of Delta-REST-GCaMP3
Authors : Chen, Y.; Song, X.; Miao, L.; Zhu, Y.; Ji, G.
Deposited on : 2012-12-25
Resolution : 2.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4661 (2.50-2.50)
Clashscore	141614	5346 (2.50-2.50)
Ramachandran outliers	138981	5231 (2.50-2.50)
Sidechain outliers	138945	5233 (2.50-2.50)
RSRZ outliers	127900	4559 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	414	

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 3439 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called RCaMP, Green fluorescent protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	392	3129	1965	526	624	14	0	1	0

There are 23 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	59	LEU	-	linker	UNP K4DIE3
A	60	GLU	-	linker	UNP K4DIE3
A	65	LYS	MET	engineered mutation	UNP P42212
A	75	ALA	VAL	engineered mutation	UNP P42212
A	87	GLY	SER	engineered mutation	UNP P42212
A	92	TYR	ASP	engineered mutation	UNP P42212
A	115	VAL	THR	engineered mutation	UNP P42212
A	118	LYS	ALA	engineered mutation	UNP P42212
A	143	LEU	HIS	engineered mutation	UNP P42212
A	151	GLY	-	linker	UNP P42212
A	152	GLY	-	linker	UNP P42212
A	153	THR	-	linker	UNP P42212
A	154	GLY	-	linker	UNP P42212
A	155	GLY	-	linker	UNP P42212
A	156	SER	-	linker	UNP P42212
A	157	MET	-	linker	UNP P42212
A	158	VAL	-	linker	UNP P42212
A	221	LEU	PHE	engineered mutation	UNP P42212
A	222	CRO	SER	chromophore	UNP P42212
A	222	CRO	TYR	chromophore	UNP P42212
A	222	CRO	GLY	chromophore	UNP P42212
A	250	ILE	VAL	engineered mutation	UNP P42212
A	372	THR	ILE	engineered mutation	UNP K4DIE3

- Molecule 2 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	4	Total 4	Ca 4	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	306	Total 306	O 306	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: RCaMP, Green fluorescent protein



4 Data and refinement statistics

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, α , β , γ	120.02Å 120.02Å 96.88Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	46.95 – 2.50 46.95 – 2.50	Depositor EDS
% Data completeness (in resolution range)	98.9 (46.95-2.50) 98.9 (46.95-2.50)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.48 (at 2.51Å)	Xtrriage
Refinement program	REFMAC 5.5.0110	Depositor
R, R_{free}	0.176 , 0.228 0.173 , 0.223	Depositor DCC
R_{free} test set	1244 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å ²)	42.0	Xtrriage
Anisotropy	0.024	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 38.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	3439	wwPDB-VP
Average B, all atoms (Å ²)	42.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.41% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CA, CRO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.10	2/3164 (0.1%)	1.00	11/4260 (0.3%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	385	GLU	CG-CD	5.83	1.60	1.51
1	A	174	GLU	CG-CD	-5.20	1.44	1.51

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	55	ARG	NE-CZ-NH2	-12.78	113.91	120.30
1	A	55	ARG	NE-CZ-NH1	8.73	124.67	120.30
1	A	353	MET	CG-SD-CE	-8.58	86.48	100.20
1	A	88	VAL	CB-CA-C	-6.15	99.72	111.40
1	A	362	ASP	CB-CG-OD1	5.75	123.48	118.30
1	A	279	ARG	CG-CD-NE	-5.74	99.74	111.80
1	A	279	ARG	NE-CZ-NH1	-5.73	117.43	120.30
1	A	366	ASP	CB-CG-OD2	-5.61	113.25	118.30
1	A	303	ARG	NE-CZ-NH1	5.50	123.05	120.30
1	A	397	ASP	CB-CG-OD1	5.29	123.06	118.30
1	A	320	LEU	CA-CB-CG	5.20	127.26	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3129	0	3029	72	0
2	A	4	0	0	0	0
3	A	306	0	0	33	0
All	All	3439	0	3029	72	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

All (72) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:306:LEU:HA	3:A:905:HOH:O	1.08	1.24
1:A:143:LEU:C	3:A:728:HOH:O	1.77	1.19
1:A:222:CRO:CA3	1:A:225:VAL:N	2.22	1.02
1:A:55:ARG:HG2	1:A:373:MET:HG3	1.45	0.98
1:A:42:LYS:HA	3:A:904:HOH:O	1.66	0.94
1:A:389:GLU:HG3	3:A:650:HOH:O	1.69	0.92
1:A:346:THR:HG22	1:A:349:GLU:H	1.34	0.91
1:A:120:SER:O	3:A:886:HOH:O	1.92	0.87
1:A:42:LYS:HA	3:A:879:HOH:O	1.74	0.87
1:A:389:GLU:CG	3:A:650:HOH:O	2.27	0.78
1:A:131:VAL:HG23	3:A:886:HOH:O	1.83	0.78
1:A:247:GLU:HG2	3:A:752:HOH:O	1.82	0.78
1:A:96:GLN:HB2	3:A:836:HOH:O	1.85	0.76
1:A:346:THR:HG22	1:A:349:GLU:HG3	1.70	0.74
1:A:222:CRO:HA31	1:A:225:VAL:N	2.00	0.74
1:A:209:LYS:HE3	3:A:661:HOH:O	1.89	0.72
1:A:346:THR:CG2	1:A:349:GLU:H	2.03	0.69
1:A:388:ARG:NH1	3:A:818:HOH:O	2.26	0.68
1:A:121:LYS:HA	3:A:886:HOH:O	1.97	0.65
1:A:350:LEU:HA	1:A:353:MET:HE3	1.79	0.64
1:A:42:LYS:N	3:A:879:HOH:O	2.33	0.62
1:A:40:ARG:C	1:A:42:LYS:H	2.05	0.60
1:A:346:THR:CG2	1:A:349:GLU:HG3	2.32	0.60
1:A:70:LYS:HB3	1:A:98:THR:HG23	1.84	0.58

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:350:LEU:HD23	1:A:353:MET:HE1	1.84	0.58
1:A:131:VAL:CG2	3:A:886:HOH:O	2.46	0.56
1:A:222:CRO:O3	1:A:225:VAL:N	2.36	0.56
1:A:88:VAL:HG13	1:A:381:THR:HB	1.87	0.56
1:A:162:GLU:HG3	3:A:882:HOH:O	2.05	0.56
1:A:306:LEU:HD12	1:A:306:LEU:H	1.71	0.56
1:A:407:LEU:O	1:A:411:MET:HG2	2.05	0.56
1:A:104:PRO:HB2	3:A:859:HOH:O	2.06	0.55
1:A:350:LEU:HD23	1:A:353:MET:CE	2.37	0.55
1:A:76:ASN:ND2	3:A:865:HOH:O	2.39	0.55
1:A:42:LYS:CA	3:A:879:HOH:O	2.42	0.54
1:A:41:ARG:C	3:A:879:HOH:O	2.46	0.54
1:A:302:THR:O	3:A:862:HOH:O	2.19	0.53
1:A:55:ARG:NH2	1:A:356:GLU:OE2	2.37	0.52
1:A:222:CRO:HD1	1:A:222:CRO:N2	2.24	0.52
1:A:305:GLN:HG3	3:A:772:HOH:O	2.10	0.52
1:A:121:LYS:CA	3:A:886:HOH:O	2.56	0.52
1:A:346:THR:HG22	1:A:349:GLU:N	2.14	0.51
1:A:70:LYS:O	1:A:98:THR:HG22	2.10	0.51
1:A:131:VAL:CB	3:A:886:HOH:O	2.60	0.49
1:A:290:ASP:HA	3:A:648:HOH:O	2.13	0.49
1:A:69:GLN:HG3	3:A:757:HOH:O	2.12	0.49
1:A:328:THR:HG21	3:A:868:HOH:O	2.13	0.49
1:A:122:ASP:C	1:A:122:ASP:OD1	2.52	0.48
1:A:376:ARG:HD3	3:A:889:HOH:O	2.12	0.48
1:A:188:GLY:HA2	1:A:202:LYS:O	2.14	0.47
1:A:97:ASN:HA	1:A:250:ILE:O	2.15	0.47
1:A:98:THR:HG22	1:A:99:PRO:HD2	1.97	0.46
1:A:305:GLN:HA	1:A:305:GLN:NE2	2.31	0.46
1:A:419:THR:O	1:A:422:GLU:N	2.49	0.46
1:A:131:VAL:HB	3:A:886:HOH:O	2.16	0.45
1:A:70:LYS:O	1:A:98:THR:CG2	2.65	0.45
1:A:392:ARG:CG	3:A:611:HOH:O	2.66	0.44
1:A:392:ARG:HD2	3:A:806:HOH:O	2.18	0.44
1:A:40:ARG:C	1:A:42:LYS:N	2.70	0.44
1:A:174:GLU:HG2	3:A:863:HOH:O	2.17	0.43
1:A:392:ARG:CZ	3:A:660:HOH:O	2.67	0.43
1:A:367:PHE:HB3	1:A:368:PRO:HD3	2.00	0.42
1:A:302:THR:O	1:A:303:ARG:HG3	2.19	0.42
1:A:346:THR:HG22	1:A:349:GLU:CG	2.42	0.42
1:A:72:GLY:HA3	1:A:97:ASN:O	2.19	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:302:THR:CG2	1:A:379:LYS:HE3	2.49	0.41
1:A:307:THR:O	1:A:309:GLU:N	2.53	0.41
1:A:267:ALA:HA	1:A:279:ARG:O	2.21	0.41
1:A:394:PHE:O	1:A:396:LYS:N	2.52	0.41
1:A:214:TRP:N	1:A:215:PRO:CD	2.84	0.41
1:A:345:PRO:HA	1:A:349:GLU:OE2	2.21	0.41
1:A:222:CRO:C3	1:A:225:VAL:CA	2.94	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	386/414 (93%)	369 (96%)	12 (3%)	5 (1%)	12 21

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	41	ARG
1	A	42	LYS
1	A	306	LEU
1	A	395	ASP
1	A	303	ARG

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	338/354 (96%)	312 (92%)	26 (8%)	13 25

All (26) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	41	ARG
1	A	42	LYS
1	A	46	THR
1	A	57	SER
1	A	59	LEU
1	A	74	LYS
1	A	88	VAL
1	A	98	THR
1	A	126	LYS
1	A	159	SER
1	A	160	LYS
1	A	174	GLU
1	A	200	THR
1	A	289	GLU
1	A	303	ARG
1	A	304	ASP
1	A	306	LEU
1	A	309	GLU
1	A	315	LYS
1	A	320	LEU
1	A	332	LYS
1	A	346	THR
1	A	373	MET
1	A	392	ARG
1	A	417	LYS
1	A	441	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	305	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	CRO	A	222	1	23,23,24	3.85	6 (26%)	30,32,34	4.02	14 (46%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CRO	A	222	1	-	2/12/31/32	0/2/2/2

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	222	CRO	CB2-CA2	16.96	1.49	1.35
1	A	222	CRO	C1-N2	3.92	1.38	1.32
1	A	222	CRO	O2-C2	3.39	1.30	1.23
1	A	222	CRO	C2-N3	-2.62	1.33	1.39
1	A	222	CRO	CA1-C1	-2.50	1.47	1.51
1	A	222	CRO	CA2-C2	-2.02	1.46	1.48

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	222	CRO	O2-C2-CA2	-16.06	121.94	130.96
1	A	222	CRO	CA2-C2-N3	10.51	108.34	103.37
1	A	222	CRO	C2-N3-C1	-4.37	105.76	107.97
1	A	222	CRO	CG2-CB2-CA2	-3.47	125.69	129.94
1	A	222	CRO	CA3-N3-C1	-3.13	123.41	127.16
1	A	222	CRO	CA1-C1-N3	-3.07	121.07	124.75
1	A	222	CRO	CA3-N3-C2	3.03	130.74	123.80

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	222	CRO	N3-C1-N2	2.94	113.49	111.45
1	A	222	CRO	O2-C2-N3	2.70	129.72	124.35
1	A	222	CRO	O3-C3-CA3	-2.70	118.25	126.39
1	A	222	CRO	CE2-CD2-CG2	-2.47	118.03	121.25
1	A	222	CRO	C2-CA2-N2	-2.22	107.38	108.93
1	A	222	CRO	CB2-CA2-C2	2.14	124.83	122.28
1	A	222	CRO	C1-CA1-N1	-2.11	106.54	109.96

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	222	CRO	C3-CA3-N3-C1
1	A	222	CRO	C3-CA3-N3-C2

There are no ring outliers.

1 monomer is involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	222	CRO	5	0

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 4 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	391/414 (94%)	-0.42	4 (1%) 82 84	25, 37, 72, 94	0

All (4) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	305	GLN	3.4
1	A	306	LEU	3.2
1	A	417	LYS	2.9
1	A	304	ASP	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	CRO	A	222	22/23	0.99	0.11	21,28,34,35	0

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	CA	A	502	1/1	0.99	0.13	29,29,29,29	0
2	CA	A	504	1/1	0.99	0.04	50,50,50,50	0
2	CA	A	501	1/1	1.00	0.12	30,30,30,30	0
2	CA	A	503	1/1	1.00	0.05	50,50,50,50	0

6.5 Other polymers [i](#)

There are no such residues in this entry.