Full wwPDB X-ray Structure Validation Report (i)

Oct 19, 2023 – 02:59 AM EDT

PDB ID	:	2ILN
Title	:	Crystal structure of the Bowman-Birk inhibitor from snail medic seeds in complex with bovine trypsin
Authors	:	Capaldi, S.; Perduca, M.; Faggion, B.; Carrizo, M.E.; Tava, A.; Ragona, L.; Monaco, H.L.
Deposited on	:	2006-10-03
Resolution	:	2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.36
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.36

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	8085 (2.00-2.00)
Clashscore	141614	9178 (2.00-2.00)
Ramachandran outliers	138981	9054 (2.00-2.00)
Sidechain outliers	138945	9053 (2.00-2.00)
RSRZ outliers	127900	7900 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	223	3% 81% 17% •
1	B	223	% 81% 17% •
2	I	62	18% 60% 21% • • 15%

2 Entry composition (i)

There are 3 unique types of molecules in this entry. The entry contains 4045 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Cationic trypsin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	223	1629	1012	279	324	14	0	0	0
1	B	223	1629	1012	279	324	14	0	0	0

- Molecule 2 is a protein called Bowman-Birk type proteinase inhibitor.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	I	53	411	249	72	77	13	0	0	0

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
3	A	169	169	169	0	0
3	B	175	175	175	0	0
3	I	32	32	32	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Cationic trypsin



• Molecule 1: Cationic trypsin



• Molecule 2: Bowman-Birk type proteinase inhibitor



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 41	Depositor
Cell constants a, b, c, α, β, γ	54.47Å 54.47Å 180.08Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	25.00 – 2.00 29.27 – 2.00	Depositor EDS
% Data completeness (in resolution range)	95.6 (25.00-2.00) 95.6 (29.27-2.00)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.09	Depositor
$< I/\sigma(I) >$ [1]	2.07 (at 2.00Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.232 , 0.287 0.230 , 0.281	Depositor DCC
R_{free} test set	1680 reflections (4.99%)	wwPDB-VP
Wilson B-factor (Å^2)	24.5	Xtriage
Anisotropy	0.926	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.35 , 35.7	EDS
L-test for twinning[2]	$< \|L\| > = 0.53$, $< L^2 > = 0.37$	Xtriage
Estimated twinning fraction	0.437 for h,-k,-l	Xtriage
F_o,F_c correlation	0.95	EDS
Total number of atoms	4045	wwPDB-VP
Average B, all atoms (Å^2)	29.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.62% of the height of the origin peak. No significant pseudotranslation is detected.*

[1]Intensities estimated from amplitudes.

[2]Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\|>5$	RMSZ	$\#\|Z\|>5$
1	A	0.46	0/1660	0.62	0/2250
1	B	0.48	0/1660	0.61	0/2250
2	I	0.52	0/422	0.80	1/572 (0.2%)
All	All	0.48	0/3742	0.64	1/5072 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
2	I	57	CYS	N-CA-C	5.27	125.23	111.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1629	0	1588	34	0
1	B	1629	0	1588	35	0
2	I	411	0	370	18	0
3	A	169	0	0	4	0
3	B	175	0	0	2	0
3	I	32	0	0	0	0
All	All	4045	0	3546	82	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 11.

All (82) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:I:31:CYS:HA	2:I:32:HIS:HB2	1.31	1.09
2:I:58:TYR:HB3	2:I:59:PRO:HD2	1.46	0.98
1:B:100:ASN:HD21	1:B:179:ASN:HD22	1.19	0.90
2:I:58:TYR:HB3	2:I:59:PRO:CD	2.03	0.89
1:A:100:ASN:HD21	1:A:179:ASN:HD22	1.21	0.88
1:A:100:ASN:ND2	1:A:179:ASN:HD22	1.74	0.86
2:I:28:ARG:HH21	2:I:32:HIS:HE1	1.24	0.82
1:B:48:ASN:HD22	1:B:50:GLN:H	1.30	0.77
1:A:48:ASN:HD22	1:A:50:GLN:H	1.33	0.76
2:I:58:TYR:CB	2:I:59:PRO:HD2	2.16	0.75
1:B:64:GLN:HE22	1:B:66:ARG:HH21	1.36	0.72
1:B:100:ASN:ND2	1:B:179:ASN:HD22	1.88	0.71
1:A:100:ASN:HD21	1:A:179:ASN:ND2	1.88	0.71
2:I:58:TYR:CB	2:I:59:PRO:CD	2.70	0.69
1:B:151:TYR:CE1	2:I:44:PHE:CZ	2.81	0.69
1:A:64:GLN:HE22	1:A:66:ARG:HH21	1.41	0.68
1:B:64:GLN:NE2	1:B:66:ARG:HE	1.92	0.66
2:I:31:CYS:HA	2:I:32:HIS:CB	2.16	0.66
1:B:143:ASN:HD22	1:B:144:THR:N	1.94	0.64
1:B:143:ASN:HD21	1:B:150:SER:H	1.45	0.63
1:B:100:ASN:HD21	1:B:179:ASN:ND2	1.94	0.63
2:I:37:SER:HB3	2:I:51:TYR:CE2	2.34	0.62
1:A:48:ASN:ND2	1:A:50:GLN:H	1.96	0.61
1:B:48:ASN:ND2	1:B:50:GLN:H	1.96	0.60
1:A:149:THR:HG23	1:A:151:TYR:CE2	2.37	0.60
1:B:143:ASN:HD22	1:B:144:THR:H	1.48	0.59
1:B:151:TYR:CE1	2:I:44:PHE:HZ	2.20	0.59
1:A:149:THR:HG23	1:A:151:TYR:HE2	1.69	0.58
1:A:165:ASP:HB3	3:A:398:HOH:O	2.04	0.57
1:B:103:ILE:HD13	1:B:104:MET:N	2.20	0.56
1:B:143:ASN:ND2	1:B:150:SER:H	2.03	0.56
1:A:87:LYS:HD2	3:A:403:HOH:O	2.04	0.56
1:A:115:ASN:HD22	1:A:117:ARG:H	1.54	0.55
1:A:217:SER:HB3	1:A:224:LYS:HD2	1.87	0.55
1:A:143:ASN:HD22	1:A:144:THR:N	2.06	0.54
2:I:13:PRO:HD2	2:I:23:GLN:O	2.08	0.53
1:A:210:GLN:NE2	3:A:404:HOH:O	2.42	0.53
1:A:101:ASN:HA	1:A:234:TYR:OH	2.09	0.52

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:240:GLN:HG3	3:A:304:HOH:O	2.08	0.52
2:I:28:ARG:HH21	2:I:32:HIS:CE1	2.14	0.52
1:A:158:LEU:HD11	1:A:188:LYS:HB3	1.91	0.52
1:B:138:ILE:HG12	1:B:199:VAL:HG22	1.93	0.51
1:B:103:ILE:HD12	1:B:237:TRP:HZ3	1.75	0.50
1:B:48:ASN:HD22	1:B:48:ASN:C	2.14	0.49
1:B:158:LEU:HD11	1:B:188:LYS:HB3	1.94	0.49
1:B:64:GLN:HE21	1:B:66:ARG:HE	1.57	0.49
1:A:143:ASN:HD21	1:A:150:SER:H	1.59	0.48
1:B:213:VAL:HA	1:B:228:TYR:CD1	2.47	0.48
1:B:48:ASN:HD21	1:B:51:TRP:HD1	1.60	0.48
1:A:149:THR:CG2	1:A:151:TYR:HE2	2.27	0.48
1:B:69:GLU:HB2	3:B:246:HOH:O	2.14	0.47
1:A:124:PRO:HD3	1:A:209:LEU:O	2.15	0.47
1:B:151:TYR:HE1	2:I:44:PHE:CZ	2.32	0.47
1:B:121:ILE:HD13	1:B:122:SER:N	2.29	0.47
2:I:10:ASP:HA	2:I:60:SER:HB3	1.97	0.47
1:A:211:GLY:HA2	1:A:229:THR:O	2.15	0.47
1:B:184:TYR:CE1	1:B:188:LYS:HE3	2.51	0.45
1:B:91:HIS:CG	1:B:92:PRO:HD2	2.51	0.45
1:B:162:ILE:HD13	1:B:181:PHE:CE1	2.52	0.45
1:A:97:ASN:ND2	1:B:173:PRO:O	2.50	0.45
1:A:91:HIS:CG	1:A:92:PRO:HD2	2.52	0.45
1:A:143:ASN:HD22	1:A:144:THR:H	1.64	0.45
1:A:64:GLN:NE2	1:A:66:ARG:HE	2.14	0.45
1:A:27:VAL:HG13	1:A:29:TYR:CZ	2.52	0.44
1:B:100:ASN:ND2	1:B:179:ASN:HB2	2.32	0.44
1:A:146:SER:HB3	1:A:221(A):ALA:HB3	1.99	0.44
2:I:31:CYS:CA	2:I:32:HIS:HB2	2.23	0.44
1:A:55:ALA:O	1:A:104:MET:HE2	2.17	0.44
2:I:58:TYR:N	2:I:58:TYR:CD1	2.86	0.43
1:A:115:ASN:ND2	1:A:117:ARG:H	2.17	0.43
1:A:143:ASN:ND2	1:A:150:SER:H	2.17	0.43
1:A:213:VAL:HA	1:A:228:TYR:CD1	2.53	0.43
1:B:160:ALA:HB1	1:B:184(A):GLY:HA2	2.01	0.43
1:A:210:GLN:HE21	1:A:210:GLN:HA	1.84	0.43
1:B:115:ASN:ND2	1:B:117:ARG:H	2.17	0.42
1:B:72:ASN:OD1	1:B:153:ASP:HB3	2.19	0.42
1:B:233:ASN:ND2	3:B:387:HOH:O	2.53	0.41
2:I:23:GLN:HB3	2:I:53:ILE:HG13	2.01	0.41
1:A:115:ASN:ND2	1:A:118:VAL:H	2.19	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:210:GLN:HA	1:B:210:GLN:HE21	1.86	0.41
1:B:101:ASN:HA	1:B:234:TYR:OH	2.21	0.40
1:A:195:SER:OG	2:I:16:ARG:C	2.60	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	221/223 (99%)	215 (97%)	5 (2%)	1 (0%)	29	23
1	B	221/223 (99%)	217 (98%)	4 (2%)	0	100	100
2	I	51/62 (82%)	43 (84%)	6 (12%)	2 (4%)	3	1
All	All	493/508 (97%)	475 (96%)	15 (3%)	3 (1%)	25	19

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	I	32	HIS
2	I	58	TYR
1	A	152	PRO

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	184/184 (100%)	179 (97%)	5 (3%)	44	46
1	B	184/184 (100%)	176 (96%)	8 (4%)	29	26
2	I	52/60 (87%)	49 (94%)	3 (6%)	20	15
All	All	420/428 (98%)	404 (96%)	16 (4%)	33	31

All (16) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	48	ASN
1	A	109	LYS
1	A	115	ASN
1	A	143	ASN
1	A	165	ASP
1	B	48	ASN
1	B	89	ILE
1	B	103	ILE
1	B	121	ILE
1	B	143	ASN
1	B	147	SER
1	B	201	CYS
1	B	244	SER
2	I	55	ASP
2	I	57	CYS
2	I	58	TYR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (21) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	30	GLN
1	A	48	ASN
1	A	64	GLN
1	A	74	ASN
1	A	100	ASN
1	A	115	ASN
1	A	143	ASN
1	A	175	GLN
1	A	210	GLN
1	A	233	ASN
1	B	30	GLN
1	B	48	ASN
1	B	64	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	74	ASN
1	B	100	ASN
1	B	115	ASN
1	B	135	GLN
1	B	143	ASN
1	B	210	GLN
1	B	233	ASN
2	I	32	HIS

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

There are no ligands in this entry.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å^2)	Q<0.9
1	A	223/223 (100%)	0.31	7 (3%) 49 48	20, 27, 35, 48	0
1	B	223/223 (100%)	0.31	3 (1%) 77 76	18, 28, 37, 45	0
2	I	53/62 (85%)	1.09	11 (20%) 1 0	21, 36, 54, 58	0
All	All	499/508 (98%)	0.39	21 (4%) 36 35	18, 28, 42, 58	0

All (21) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	147	SER	6.0
2	I	59	PRO	5.6
2	I	58	TYR	5.4
2	I	57	CYS	5.0
1	A	148	GLY	4.5
2	I	31	CYS	4.2
2	I	60	SER	3.9
2	I	51	TYR	3.9
2	I	32	HIS	3.8
1	B	147	SER	3.6
2	I	56	PHE	3.6
2	I	11	PHE	3.5
1	A	152	PRO	3.4
2	I	33	SER	3.3
1	A	146	SER	2.8
1	A	151	TYR	2.6
2	I	44	PHE	2.5
1	A	149	THR	2.4
1	A	154	VAL	2.2
1	B	223	ASN	2.2
1	B	89	ILE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

There are no ligands in this entry.

6.5 Other polymers (i)

There are no such residues in this entry.