Full wwPDB NMR Structure Validation Report (i)

Feb 14, 2022 – 12:07 AM EST

PDB ID	:	1IMT
Title	:	MAMBA INTESTINAL TOXIN 1, NMR, 39 STRUCTURES
Authors	:	Boisbouvier, J.; Albrand, J.-P.; Blackledge, M.; Jaquinod, M.; Schweitz, H.; Lazdunski, M.; Marion, D.
Deposited on	:	1998-04-14

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/NMRValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity	:	4.02b-467
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
RCI	:	v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV	:	Wang et al. (2010)
ShiftChecker	:	2.26
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.26

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
SOLUTION NMR

The overall completeness of chemical shifts assignment was not calculated.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	158937	12864
Ramachandran outliers	154571	11451
Sidechain outliers	154315	11428

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Mol	Chain	Length	Quality of chain
1	A	80	82% 12% 5%

2 Ensemble composition and analysis (i)

This entry contains 39 models. Model 2 is the overall representative, medoid model (most similar to other models).

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	**Residue range (total)**	**Backbone RMSD (Å)**	**Medoid model**
1	A:5-A:80 (76)	0.61	2

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 7 clusters and 5 single-model clusters were found.

Cluster number	Models
1	2, 5, 9, 10, 19, 27, 30, 31, 35, 37
2	7, 14, 18, 22, 25, 29, 34, 36
3	1, 12, 21, 33
4	13, 23, 32
5	16, 28, 38
6	3, 4, 17
7	6, 15, 24
Single-model clusters	8; 11; 20; 26; 39

3 Entry composition (i)

There is only 1 type of molecule in this entry. The entry contains 1172 atoms, of which 584 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called INTESTINAL TOXIN 1.

Mol	Chain	Residues	Atoms						Trace
			Total	C	H	N	O	S	
1	A	80	1172	362	584	109	106	11	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	?	-	LYS	deletion	UNP P25687

4 Residue-property plots (i)

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

• Molecule 1: INTESTINAL TOXIN 1

Chain A: 82% 12% 5%



4.2 Scores per residue for each member of the ensemble

Colouring as in section 4.1 above.

4.2.1 Score per residue for model 1

• Molecule 1: INTESTINAL TOXIN 1

Chain A: 89% 6% 5%



4.2.2 Score per residue for model 2 (medoid)

• Molecule 1: INTESTINAL TOXIN 1

Chain A: 84% 10% • 5%



4.2.3 Score per residue for model 3

• Molecule 1: INTESTINAL TOXIN 1

Chain A: 88% 8% 5%



4.2.4 Score per residue for model 4

• Molecule 1: INTESTINAL TOXIN 1

Chain A: 84% 10% • 5%



4.2.5 Score per residue for model 5

• Molecule 1: INTESTINAL TOXIN 1

Chain A: 81% 14% 5%



4.2.6 Score per residue for model 6

• Molecule 1: INTESTINAL TOXIN 1

Chain A: 81% 12% • 5%

4.2.7 Score per residue for model 7

• Molecule 1: INTESTINAL TOXIN 1

Chain A: 81% 12% • 5%



4.2.8 Score per residue for model 8

- Molecule 1: INTESTINAL TOXIN 1

Chain A: 82% 10% • 5%



4.2.9 Score per residue for model 9

- Molecule 1: INTESTINAL TOXIN 1

Chain A: 81% 14% 5%

4.2.10 Score per residue for model 10

- Molecule 1: INTESTINAL TOXIN 1

Chain A: 85% 10% 5%

4.2.11 Score per residue for model 11

- Molecule 1: INTESTINAL TOXIN 1

Chain A: 85% 10% 5%

4.2.12 Score per residue for model 12

- Molecule 1: INTESTINAL TOXIN 1

Chain A: 81% 12% • 5%



4.2.13 Score per residue for model 13

• Molecule 1: INTESTINAL TOXIN 1

Chain A: 86% 9% 5%



4.2.14 Score per residue for model 14

• Molecule 1: INTESTINAL TOXIN 1

Chain A: 81% 14% 5%

4.2.15 Score per residue for model 15

• Molecule 1: INTESTINAL TOXIN 1

Chain A: 79% 16% 5%



4.2.16 Score per residue for model 16

• Molecule 1: INTESTINAL TOXIN 1

Chain A: 85% 10% 5%



4.2.17 Score per residue for model 17

• Molecule 1: INTESTINAL TOXIN 1

Chain A: 85% 10% 5%

4.2.18 Score per residue for model 18

• Molecule 1: INTESTINAL TOXIN 1

Chain A: 82% 12% 5%



4.2.19 Score per residue for model 19

• Molecule 1: INTESTINAL TOXIN 1

Chain A: 84% 11% 5%



4.2.20 Score per residue for model 20

• Molecule 1: INTESTINAL TOXIN 1

Chain A: 84% 9% • 5%



4.2.21 Score per residue for model 21

• Molecule 1: INTESTINAL TOXIN 1

Chain A: 85% 10% 5%

4.2.22 Score per residue for model 22

• Molecule 1: INTESTINAL TOXIN 1

Chain A: 86% 8% • 5%



4.2.23 Score per residue for model 23

- Molecule 1: INTESTINAL TOXIN 1

Chain A: 78% 16% • 5%



4.2.24 Score per residue for model 24

- Molecule 1: INTESTINAL TOXIN 1

Chain A: 81% 14% 5%

4.2.25 Score per residue for model 25

- Molecule 1: INTESTINAL TOXIN 1

Chain A: 85% 9% • 5%



4.2.26 Score per residue for model 26

- Molecule 1: INTESTINAL TOXIN 1

Chain A: 85% 10% 5%

4.2.27 Score per residue for model 27

- Molecule 1: INTESTINAL TOXIN 1

Chain A: 82% 9% • • 5%

4.2.28 Score per residue for model 28

• Molecule 1: INTESTINAL TOXIN 1

Chain A: 82% 10% • 5%



4.2.29 Score per residue for model 29

• Molecule 1: INTESTINAL TOXIN 1

Chain A: 85% 10% 5%



4.2.30 Score per residue for model 30

• Molecule 1: INTESTINAL TOXIN 1

Chain A: 82% 12% 5%



4.2.31 Score per residue for model 31

• Molecule 1: INTESTINAL TOXIN 1

Chain A: 86% 9% 5%



4.2.32 Score per residue for model 32

• Molecule 1: INTESTINAL TOXIN 1

Chain A: 78% 15% • 5%

4.2.33 Score per residue for model 33

• Molecule 1: INTESTINAL TOXIN 1

Chain A: 85% 6% • 5%



4.2.34 Score per residue for model 34

• Molecule 1: INTESTINAL TOXIN 1

Chain A: 82% 12% 5%



4.2.35 Score per residue for model 35

• Molecule 1: INTESTINAL TOXIN 1

Chain A: 85% 10% 5%



4.2.36 Score per residue for model 36

• Molecule 1: INTESTINAL TOXIN 1

Chain A: 82% 12% 5%

4.2.37 Score per residue for model 37

• Molecule 1: INTESTINAL TOXIN 1

Chain A: 84% 11% 5%

4.2.38 Score per residue for model 38

- Molecule 1: INTESTINAL TOXIN 1

Chain A:



4.2.39 Score per residue for model 39

- Molecule 1: INTESTINAL TOXIN 1

Chain A:



5 Refinement protocol and experimental data overview (i)

The models were refined using the following method: *RESTRAINED SIMULATED ANNEALING AND RESTRAINED MOLECULAR DYNAMICS ENERGY REFINEMENT*.

Of the 110 calculated structures, 39 were deposited, based on the following criterion: *LEAST RESTRAINT VIOLATION AND BEST PHYSICAL ENERGY*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
Discover	refinement	
Discover	structure solution	

No chemical shift data was provided.

6 Model quality (i)

6.1 Standard geometry (i)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#Z>5	RMSZ	#Z>5
1	A	0.50±0.01	0±0/574 (0.0± 0.0%)	0.91±0.03	0±0/773 (0.0± 0.0%)
All	All	0.50	0/22386 (0.0%)	0.91	2/30147 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	Chirality	Planarity
1	A	0.0±0.0	0.6±0.7
All	All	0	23

There are no bond-length outliers.

All unique angle outliers are listed below.

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})	Models	
								Worst	Total
1	A	28	VAL	CA-CB-CG1	5.65	119.38	110.90	39	2

There are no chirality outliers.

All unique planar outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Group	Models (Total)
1	A	29	ARG	Sidechain	6
1	A	57	HIS	Sidechain	5
1	A	54	ARG	Sidechain	4
1	A	75	PHE	Sidechain	3
1	A	46	HIS	Sidechain	2
1	A	9	ARG	Sidechain	2
1	A	50	PHE	Sidechain	1

6.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	561	551	551	0±1
All	All	21879	21489	21489	12

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 0.

All unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:23:LEU:HD22	1:A:57:HIS:CE1	0.76	2.16	9	2
1:A:23:LEU:HD13	1:A:57:HIS:NE2	0.56	2.16	23	4
1:A:23:LEU:HG	1:A:57:HIS:CE1	0.43	2.48	12	1
1:A:22:SER:CB	1:A:28:VAL:HG13	0.42	2.44	39	1
1:A:69:GLN:H	1:A:69:GLN:CD	0.42	2.18	33	1
1:A:69:GLN:CD	1:A:69:GLN:N	0.41	2.74	9	1
1:A:59:CYS:HB3	1:A:60:PRO:CD	0.40	2.46	12	1
1:A:23:LEU:HD12	1:A:57:HIS:CE1	0.40	2.51	27	1

6.3 Torsion angles (i)

6.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	75/80 (94%)	60±3 (80±4%)	12±3 (16±4%)	3±2 (4±2%)	5 31
All	All	2925/3120 (94%)	2347 (80%)	461 (16%)	117 (4%)	5 31

All 19 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	63	PRO	23
1	A	64	ASN	21
1	A	37	SER	10
1	A	15	LYS	9
1	A	25	ILE	8
1	A	5	GLY	6
1	A	46	HIS	6
1	A	31	CYS	6
1	A	26	LYS	5
1	A	14	GLY	4
1	A	47	LYS	4
1	A	48	ILE	3
1	A	74	LYS	3
1	A	10	ASP	3
1	A	45	SER	2
1	A	6	ALA	1
1	A	23	LEU	1
1	A	79	SER	1
1	A	57	HIS	1

6.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	65/68 (96%)	59±1 (91±2%)	6±1 (9±2%)	13 60
All	All	2535/2652 (96%)	2311 (91%)	224 (9%)	13 60

All 26 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	9	ARG	39
1	A	18	CYS	30
1	A	69	GLN	23
1	A	34	VAL	21
1	A	47	LYS	17
1	A	29	ARG	16
1	A	46	HIS	13

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Models (Total)
1	A	71	SER	9
1	A	28	VAL	8
1	A	57	HIS	8
1	A	41	CYS	6
1	A	13	CYS	4
1	A	59	CYS	4
1	A	80	LYS	4
1	A	25	ILE	4
1	A	36	THR	3
1	A	75	PHE	3
1	A	54	ARG	2
1	A	7	CYS	2
1	A	65	LEU	2
1	A	50	PHE	1
1	A	15	LYS	1
1	A	45	SER	1
1	A	64	ASN	1
1	A	55	MET	1
1	A	23	LEU	1

6.3.3 RNA (i)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates (i)

There are no monosaccharides in this entry.

6.6 Ligand geometry (i)

There are no ligands in this entry.

6.7 Other polymers (i)

There are no such molecules in this entry.

6.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

7 Chemical shift validation (i)

No chemical shift data were provided