

wwPDB X-ray Structure Validation Summary Report (i)

Nov 15, 2023 – 12:08 AM JST

PDB ID : 6IM1

Title: Crystal structure of a highly thermostable carbonic anhydrase from Perse-

phonella marina EX-H1

Authors: Jin, M.S.; Kim, S.; Sung, J.; Yeon, J.; Choi, S.H.

Deposited on : 2018-10-21

Resolution : 2.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at

https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity: 4.02b-467

Mogul : 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : 1.13

EDS : 2.36

buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

 $Refmac \quad : \quad 5.8.0158$

CCP4 : 7.0.044 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.36

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X- $RAY\ DIFFRACTION$

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive	Similar resolution
Metric	$(\# \mathrm{Entries})$	$(\# ext{Entries}, ext{ resolution range}(ext{Å}))$
R_{free}	130704	8085 (2.00-2.00)
Clashscore	141614	9178 (2.00-2.00)
Ramachandran outliers	138981	9054 (2.00-2.00)
Sidechain outliers	138945	9053 (2.00-2.00)
RSRZ outliers	127900	7900 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain		
1	A	243	2%	20/	00/
1	A	240	83%	9%	9%
1	В	243	86%	6%	9%
1	\mathbf{C}	243	81%	10%	9%
1	C	210	6%	1070	370
1	D	243	84%	7%	9%
1	Б	0.49	5%		
1	Е	243	79%	13%	9%
1	F	243	81%	10%	9%



2 Entry composition (i)

There are 5 unique types of molecules in this entry. The entry contains 11390 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Carbonic anhydrase (Carbonate dehydratase).

Mol	Chain	Residues		At	oms			ZeroOcc	AltConf	Trace
1	A	222	Total	С	N	О	S	0	0	0
1	A	222	1803	1152	315	326	10	O	0	
1	В	222	Total	С	N	О	S	0	0	0
1	Б	222	1803	1152	315	326	10	U	U	
1	\mathbf{C}	222	Total	С	N	О	S	0	0	0
1		222	1803	1152	315	326	10	U	U	0
1	D	222	Total	С	N	О	S	0	0	0
1	D	222	1803	1152	315	326	10	U	0	
1	Е	222	Total	С	N	О	S	0	0	0
1	12	222	1803	1152	315	326	10	U	0	
1	F	222	Total	С	N	О	S	0	0	0
1	I'	222	1802	1152	315	325	10	U	U	

• Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn) (labeled as "Ligand of Interest" by depositor).


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total Zn 1 1	0	0
2	В	1	Total Zn 1 1	0	0
2	С	1	Total Zn 1 1	0	0
2	D	1	Total Zn 1 1	0	0
2	E	1	Total Zn 1 1	0	0
2	F	1	Total Zn 1 1	0	0

• Molecule 3 is CALCIUM ION (three-letter code: CA) (formula: Ca) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	6	Total Ca 6 6	0	0
3	В	6	Total Ca 6 6	0	0
3	D	3	Total Ca 3 3	0	0

 \bullet Molecule 4 is TETRAETHYLENE GLYCOL (three-letter code: PG4) (formula: $C_8H_{18}O_5)$ (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C O 13 8 5	0	0
4	В	1	Total C O 13 8 5	0	0
4	E	1	Total C O 13 8 5	0	0
4	F	1	Total C O 13 8 5	0	0

• Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	141	Total O 141 141	0	0
5	В	126	Total O 126 126	0	0

Continued on next page...



Continued from previous page...


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	С	53	Total O 53 53	0	0
5	D	54	Total O 54 54	0	0
5	Е	61	Total O 61 61	0	0
5	F	65	Total O 65 65	0	0



3 Residue-property plots (i)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Carbonic anhydrase (Carbonate dehydratase)






• Molecule 1: Carbonic anhydrase (Carbonate dehydratase)



• Molecule 1: Carbonic anhydrase (Carbonate dehydratase)





4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants	118.92Å 120.07Å 124.07Å	Depositor
a, b, c, α , β , γ	90.00° 90.00° 90.00°	Depositor
Resolution (Å)	124.07 - 2.00	Depositor
Resolution (A)	40.55 - 2.00	EDS
% Data completeness	$99.5 \ (124.07 - 2.00)$	Depositor
(in resolution range)	99.6 (40.55-2.00)	EDS
R_{merge}	(Not available)	Depositor
R_{sum}	(Not available)	Depositor
$< I/\sigma(I) > 1$	3.89 (at 2.00Å)	Xtriage
Refinement program	REFMAC 5.8.0135	Depositor
R, R_{free}	0.204 , 0.236	Depositor
it, it free	0.214 , 0.242	DCC
R_{free} test set	5670 reflections (4.76%)	wwPDB-VP
Wilson B-factor (Å ²)	37.2	Xtriage
Anisotropy	0.026	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	$0.36 \; , 48.4$	EDS
L-test for twinning ²	$< L > = 0.51, < L^2> = 0.34$	Xtriage
	0.000 for -h,l,k	
	0.000 for -l,-k,-h	
Estimated twinning fraction	0.000 for k,h,-l	Xtriage
	0.000 for k,l,h	
	0.000 for l,h,k	
F_o, F_c correlation	0.96	EDS
Total number of atoms	11390	wwPDB-VP
Average B, all atoms (\mathring{A}^2)	48.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The analyses of the Patterson function reveals a significant off-origin peak that is 59.62 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.7186e-05. The detected translational NCS is most likely also responsible for the elevated intensity ratio.

²Theoretical values of <|L|>, $<L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: CA, ZN, PG4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond	angles
IVIOI	Chain	RMSZ	# Z > 5	RMSZ	# Z > 5
1	A	1.12	0/1852	0.93	0/2494
1	В	1.09	0/1852	0.94	0/2494
1	С	0.92	0/1852	0.89	0/2494
1	D	0.90	0/1852	0.89	0/2494
1	Е	1.00	0/1852	0.87	0/2494
1	F	0.99	0/1851	0.87	0/2492
All	All	1.01	0/11111	0.90	0/14962

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1803	0	1776	13	0
1	В	1803	0	1777	16	0
1	С	1803	0	1777	22	0
1	D	1803	0	1777	11	0
1	Е	1803	0	1777	28	0
1	F	1802	0	1777	14	0
2	A	1	0	0	0	0

Continued on next page...



Continued from previous page...

Mol	Chain		H(model)	H(added)	Clashes	Symm-Clashes
2	В	1	0	0	0	0
2	С	1	0	0	0	0
2	D	1	0	0	0	0
2	Е	1	0	0	0	0
2	F	1	0	0	0	0
3	A	6	0	0	0	0
3	В	6	0	0	0	0
3	D	3	0	0	0	0
4	A	13	0	18	0	0
4	В	13	0	18	0	0
4	Е	13	0	18	0	0
4	F	13	0	18	0	0
5	A	141	0	0	4	0
5	В	126	0	0	3	0
5	С	53	0	0	1	0
5	D	54	0	0	0	0
5	Ε	61	0	0	4	0
5	F	65	0	0	0	0
All	All	11390	0	10733	103	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 103 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	$egin{aligned} & ext{Interatomic} \ & ext{distance} \ & ext{(Å)} \end{aligned}$	$egin{aligned} ext{Clash} \ ext{overlap } (ext{Å}) \end{aligned}$
1:C:159:MET:HE1	1:C:223:ILE:HD12	1.17	1.17
1:E:159:MET:HE1	1:E:223:ILE:HD12	1.22	1.16
1:C:159:MET:CE	1:C:223:ILE:HD12	1.82	1.08
1:E:159:MET:CE	1:E:223:ILE:HD12	1.84	1.06
1:C:44:CYS:HG	1:C:197:CYS:HG	1.05	0.89

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.



The Analysed column shows the r	number of residu	ues for which	the backbone	conformation	was
analysed, and the total number of	residues.				

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percei	ntiles
1	A	$220/243\ (90\%)$	215 (98%)	5 (2%)	0	100	100
1	В	$220/243\ (90\%)$	214 (97%)	6 (3%)	0	100	100
1	C	$220/243\ (90\%)$	214 (97%)	6 (3%)	0	100	100
1	D	$220/243\ (90\%)$	214 (97%)	6 (3%)	0	100	100
1	E	$220/243\ (90\%)$	215 (98%)	5 (2%)	0	100	100
1	F	$220/243\ (90\%)$	217 (99%)	3 (1%)	0	100	100
All	All	$1320/1458 \; (90\%)$	1289 (98%)	31 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	193/209 (92%)	189 (98%)	4 (2%)	53 57
1	В	193/209 (92%)	190 (98%)	3 (2%)	62 67
1	С	193/209 (92%)	190 (98%)	3 (2%)	62 67
1	D	193/209 (92%)	190 (98%)	3 (2%)	62 67
1	Е	193/209 (92%)	189 (98%)	4 (2%)	53 57
1	F	193/209 (92%)	189 (98%)	4 (2%)	53 57
All	All	1158/1254 (92%)	1137 (98%)	21 (2%)	59 63

5 of 21 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	Ε	88	TYR
1	F	24	SER
1	F	222	LYS
1	F	88	TYR
1	Е	156	TRP



Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	В	236	ASN
1	D	236	ASN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

Of 25 ligands modelled in this entry, 21 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Tuno	Chain	Res Link Bond lengths		Bond angles					
IVIOI	Type	Chain	Res	Lilik	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	PG4	A	305	-	12,12,12	0.73	0	11,11,11	0.81	0
4	PG4	В	306	-	12,12,12	0.89	0	11,11,11	0.83	0
4	PG4	F	302	-	12,12,12	0.95	0	11,11,11	1.09	0
4	PG4	Е	302	-	12,12,12	0.85	0	11,11,11	0.85	1 (9%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.



Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	PG4	A	305	-	-	2/10/10/10	-
4	PG4	В	306	-	-	3/10/10/10	-
4	PG4	F	302	-	-	4/10/10/10	-
4	PG4	Е	302	-	-	1/10/10/10	-

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\mathbf{Observed}(^o)$	$\operatorname{Ideal}(^{o})$
4	Ε	302	PG4	O3-C4-C3	2.17	120.18	110.39

There are no chirality outliers.

5 of 10 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	A	305	PG4	O1-C1-C2-O2
4	F	302	PG4	O2-C3-C4-O3
4	A	305	PG4	O2-C3-C4-O3
4	В	306	PG4	O2-C3-C4-O3
4	Е	302	PG4	O3-C5-C6-O4

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers (i)

There are no such residues in this entry.



5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<rsrz></rsrz>	# RSRZ > 2	$OWAB(A^2)$	Q < 0.9
1	A	222/243 (91%)	-0.10	4 (1%) 68 66	23, 33, 59, 83	0
1	В	222/243 (91%)	-0.07	6 (2%) 54 53	23, 36, 61, 80	0
1	С	222/243 (91%)	0.46	18 (8%) 12 11	32, 54, 82, 102	0
1	D	222/243 (91%)	0.47	15 (6%) 17 16	32, 52, 85, 114	0
1	E	222/243 (91%)	0.20	12 (5%) 25 24	32, 47, 78, 105	0
1	F	222/243 (91%)	0.27	12 (5%) 25 24	31, 47, 78, 98	0
All	All	1332/1458 (91%)	0.20	67 (5%) 28 28	23, 45, 78, 114	0

The worst 5 of 67 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	С	22	GLY	4.7
1	D	169	LEU	4.5
1	F	169	LEU	4.5
1	Е	22	GLY	4.4
1	D	171	HIS	4.3

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.




6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\operatorname{B-factors}({ ext{\AA}}^2)$	Q < 0.9
3	CA	A	303	1/1	0.69	0.15	66,66,66,66	0
4	PG4	A	305	13/13	0.72	0.35	63,73,83,87	0
4	PG4	В	306	13/13	0.72	0.29	66,72,83,90	0
4	PG4	Ε	302	13/13	0.74	0.21	61,67,73,75	0
3	CA	D	303	1/1	0.81	0.12	101,101,101,101	0
3	CA	A	308	1/1	0.81	0.08	78,78,78,78	0
4	PG4	F	302	13/13	0.81	0.20	51,63,71,79	0
3	CA	В	308	1/1	0.82	0.07	78,78,78,78	0
3	CA	В	304	1/1	0.82	0.12	60,60,60,60	0
3	CA	A	307	1/1	0.84	0.09	60,60,60,60	0
3	CA	В	305	1/1	0.88	0.06	$65,\!65,\!65,\!65$	0
3	CA	D	304	1/1	0.90	0.09	73,73,73,73	0
3	CA	D	302	1/1	0.93	0.06	53,53,53,53	0
3	CA	A	304	1/1	0.95	0.05	64,64,64,64	0
3	CA	В	307	1/1	0.96	0.07	45,45,45,45	0
3	CA	В	302	1/1	0.97	0.06	39,39,39,39	0
3	CA	A	302	1/1	0.97	0.05	40,40,40,40	0
3	CA	В	303	1/1	0.98	0.06	54,54,54,54	0
2	ZN	F	301	1/1	0.98	0.09	46,46,46,46	0
2	ZN	D	301	1/1	0.98	0.04	48,48,48,48	0
3	CA	A	306	1/1	0.98	0.07	41,41,41,41	0
2	ZN	В	301	1/1	0.99	0.06	34,34,34,34	0
2	ZN	С	301	1/1	0.99	0.06	51,51,51,51	0
2	ZN	A	301	1/1	0.99	0.07	32,32,32,32	0
2	ZN	Е	301	1/1	0.99	0.07	44,44,44,44	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.









Electron density around PG4 A 305: $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_{o}\text{-}\mathrm{DF}_{c}$ (at 3 rmsd) in purple (negative) and green (positive)




Electron density around PG4 B 306:


 $2 {
m mF}_o {
m -DF}_c$ (at 0.7 rmsd) in gray ${
m mF}_o {
m -DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)













Electron density around CA D 303: $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_{o}\text{-}\mathrm{DF}_{c}$ (at 3 rmsd) in purple (negative) and green (positive)



Electron density around CA A 308: $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray ${\rm mF}_o\text{-}{\rm DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)









Electron density around CA B 308: $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_{o}\text{-}\mathrm{DF}_{c}$ (at 3 rmsd) in purple (negative) and green (positive)

















Electron density around CA D 304: $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_{o}\text{-}\mathrm{DF}_{c}$ (at 3 rmsd) in purple (negative) and green (positive)









Electron density around CA A 304: $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_{o}\text{-}\mathrm{DF}_{c}$ (at 3 rmsd) in purple (negative) and green (positive)

































Electron density around CA A 306: $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_{o}\text{-}\mathrm{DF}_{c}$ (at 3 rmsd) in purple (negative) and green (positive)
















6.5 Other polymers (i)

There are no such residues in this entry.

