Full wwPDB X-ray Structure Validation Report (i)

Nov 15, 2023 – 12:10 AM JST

PDB ID	:	6IM7
Title	:	CueO-12.1 multicopper oxidase
Authors	:	Wongsantichon, J.; Robinson, R.; Ghadessy, F.
Deposited on	:	2018-10-22
Resolution	:	1.97 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.36
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.36

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 1.97 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	11647 (2.00-1.96)
Clashscore	141614	1014 (1.98-1.98)
Ramachandran outliers	138981	1006 (1.98-1.98)
Sidechain outliers	138945	1006 (1.98-1.98)
RSRZ outliers	127900	11410 (2.00-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	510	

2 Entry composition (i)

There are 3 unique types of molecules in this entry. The entry contains 3926 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Blue copper oxidase CueO,12.1 peptide,Blue copper oxidase CueO.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	461	3573	2269	623	657	24	0	5	0

There are 23 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	358	ILE	MET	engineered mutation	UNP P36649
A	517	LYS	-	expression tag	UNP P36649
A	518	ASP	-	expression tag	UNP P36649
A	519	PRO	-	expression tag	UNP P36649
A	520	ASN	-	expression tag	UNP P36649
A	521	SER	-	expression tag	UNP P36649
A	522	SER	-	expression tag	UNP P36649
A	523	SER	-	expression tag	UNP P36649
A	524	VAL	-	expression tag	UNP P36649
A	525	ASP	-	expression tag	UNP P36649
A	526	LYS	-	expression tag	UNP P36649
A	527	LEU	-	expression tag	UNP P36649
A	528	ALA	-	expression tag	UNP P36649
A	529	ALA	-	expression tag	UNP P36649
A	530	ALA	-	expression tag	UNP P36649
A	531	LEU	-	expression tag	UNP P36649
A	532	GLU	-	expression tag	UNP P36649
A	533	HIS	-	expression tag	UNP P36649
A	534	HIS	-	expression tag	UNP P36649
A	535	HIS	-	expression tag	UNP P36649
A	536	HIS	-	expression tag	UNP P36649
A	537	HIS	-	expression tag	UNP P36649
A	538	HIS	-	expression tag	UNP P36649

- Molecule 2 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Ca		
2	A	1	1	1	0	0

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
3	A	352	352	352	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Blue copper oxidase CueO,12.1 peptide,Blue copper oxidase CueO

Chain A:



4 Data and refinement statistics (i)

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α, β, γ	76.11Å 100.73Å 59.04Å 90.00° 96.17° 90.00°	Depositor
Resolution (Å)	29.35 – 1.97 29.35 – 1.97	Depositor EDS
% Data completeness (in resolution range)	99.1 (29.35-1.97) 97.2 (29.35-1.97)	Depositor EDS
R_{merge}	0.04	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	7.86 (at 1.96Å)	Xtriage
Refinement program	PHENIX 1.13_2998, PHENIX 1.13_2998	Depositor
R, R_{free}	0.140 , 0.184 0.140 , 0.184	Depositor DCC
R_{free} test set	1564 reflections (5.06%)	wwPDB-VP
Wilson B-factor (Å^2)	22.8	Xtriage
Anisotropy	0.007	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.32 , 48.2	EDS
L-test for twinning[2]	$< \|L\| > = 0.49$, $< L^2 > = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o,F_c correlation	0.97	EDS
Total number of atoms	3926	wwPDB-VP
Average B, all atoms (Å^2)	27.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.23% of the height of the origin peak. No significant pseudotranslation is detected.*

[1]Intensities estimated from amplitudes.

[2]Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\| >5$	RMSZ	$\#\|Z\| >5$
1	A	0.36	0/3670	0.57	0/4984

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3573	0	3551	20	0
2	A	1	0	0	0	0
3	A	352	0	0	6	0
All	All	3926	0	3551	20	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (20) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:56:GLN:HG3	1:A:63:THR:HG23	1.81	0.62

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:271[B]:SER:OG	3:A:701:HOH:O	2.18	0.55
1:A:355:MET:SD	1:A:441:MET:HG3	2.48	0.53
1:A:100:LEU:HD21	1:A:140:PHE:HB2	1.92	0.52
1:A:166:GLU:OE1	3:A:702:HOH:O	2.19	0.52
1:A:453:ARG:NH1	3:A:715:HOH:O	2.42	0.52
1:A:279:GLU:OE2	1:A:470:LYS:NZ	2.37	0.49
1:A:410:ILE:HG21	1:A:512:LEU:HD23	1.98	0.46
1:A:82:ARG:NH1	3:A:719:HOH:O	2.46	0.46
1:A:228[A]:ARG:NH2	1:A:290:ASN:HB3	2.31	0.45
1:A:337:LEU:HD21	1:A:468:GLY:HA2	1.97	0.45
1:A:321:PRO:HG3	3:A:721:HOH:O	2.18	0.44
1:A:276:LEU:HD11	1:A:474:LYS:HB2	2.00	0.44
1:A:303:MET:HE3	1:A:441:MET:SD	2.59	0.43
1:A:485:LYS:HE2	1:A:487:ASN:OD1	2.20	0.41
1:A:130:ASN:OD1	3:A:703:HOH:O	2.22	0.41
1:A:452:PHE:HB3	1:A:484:VAL:HG12	2.02	0.41
1:A:140:PHE:CZ	1:A:158:GLY:HA3	2.56	0.41
1:A:102:TRP:HB3	1:A:105:LEU:HD22	2.03	0.40
1:A:364:MET:O	1:A:368:MET:HG2	2.20	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	460/510 (90%)	446 (97%)	14 (3%)	0	100 100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	385/421 (91%)	377 (98%)	8 (2%)	53 47

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	56	GLN
1	A	130	ASN
1	A	199	GLN
1	A	221	TYR
1	A	280	ARG
1	A	321	PRO
1	A	332	ASP
1	A	360	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	428	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

Of 1 ligands modelled in this entry, 1 is monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å^2)	Q<0.9
1	A	461/510 (90%)	-0.22	11 (2%) 59 61	15, 23, 51, 72	0

All (11) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	403	ASP	3.2
1	A	365	GLN	3.1
1	A	339	ALA	2.8
1	A	373	ASP	2.8
1	A	377	ALA	2.6
1	A	376	MET	2.4
1	A	61	GLY	2.4
1	A	42	THR	2.3
1	A	358	ILE	2.2
1	A	518	ASP	2.0
1	A	404	PHE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
2	CA	A	601	1/1	0.99	0.02	22,22,22,22	1

6.5 Other polymers (i)

There are no such residues in this entry.