Full wwPDB X-ray Structure Validation Report (i)

Mar 23, 2024 – 10:29 AM EDT

PDB ID	:	5INE
Title	:	Crystal structure of the prefusion glycoprotein of LCMV
Authors	:	Hastie, K.M.; Saphire, E.O.
Deposited on	:	2016-03-07
Resolution	:	3.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.36.1
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.36.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 3.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1659 (3.60-3.40)
Clashscore	141614	1036 (3.58-3.42)
Ramachandran outliers	138981	1005 (3.58-3.42)
Sidechain outliers	138945	1006 (3.58-3.42)
RSRZ outliers	127900	1559 (3.60-3.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	438	5% 71% 8% 21%
1	B	438	4% 69% 8% 23%
2	C	4	50% 50%
2	D	4	50% 50%
2	J	4	50% 50%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	M	4	25% 75%
2	N	4	75% 25%
3	E	2	100%
4	F	3	100%
4	G	3	100%
4	H	3	67% 33%
4	I	3	33% 67%
4	K	3	67% 33%
4	L	3	100%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	MAN	M	4	-	-	-	X
3	NAG	E	2	-	-	-	X
4	NAG	H	2	-	-	-	X
4	NAG	I	2	-	-	-	X
4	BMA	L	3	-	-	-	X
5	NAG	A	521	-	-	-	X
5	NAG	A	522	-	-	-	X

2 Entry composition (i)

There are 5 unique types of molecules in this entry. The entry contains 6102 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Pre-glycoprotein polyprotein GP complex.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	348	2800	1772	480	523	25	0	0	0
1	B	339	2720	1720	469	506	25	0	0	0

- Molecule 2 is an oligosaccharide called alpha-D-mannopyranose-(1-3)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	C	4	50	28	2	20	0	0	0
2	D	4	50	28	2	20	0	0	0
2	J	4	50	28	2	20	0	0	0
2	M	4	50	28	2	20	0	0	0
2	N	4	50	28	2	20	0	0	0

- Molecule 3 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
3	E	2	28	16	2	10	0	0	0

- Molecule 4 is an oligosaccharide called beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
4	F	3	39	22	2	15	0	0	0
4	G	3	39	22	2	15	0	0	0
4	H	3	39	22	2	15	0	0	0
4	I	3	39	22	2	15	0	0	0
4	K	3	39	22	2	15	0	0	0
4	L	3	39	22	2	15	0	0	0

- Molecule 5 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: $C_8H_{15}NO_6$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
5	A	1	14	8	1	5	0	0
			Total	C	N	O		
5	A	1	14	8	1	5	0	0
			Total	C	N	O		
5	A	1	14	8	1	5	0	0
			Total	C	N	O		
5	B	1	14	8	1	5	0	0
			Total	C	N	O		
5	B	1	14	8	1	5	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Pre-glycoprotein polyprotein GP complex

Chain A:



• Molecule 1: Pre-glycoprotein polyprotein GP complex

Chain B:



• Molecule 2: alpha-D-mannopyranose-(1-3)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain C:

• Molecule 2: alpha-D-mannopyranose-(1-3)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain D:

• Molecule 2: alpha-D-mannopyranose-(1-3)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain J:

• Molecule 2: alpha-D-mannopyranose-(1-3)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain M: 

• Molecule 2: alpha-D-mannopyranose-(1-3)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain N:

• Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain E: 

• Molecule 4: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain F: 

• Molecule 4: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain G: 100%

NAG1 NAG2 BMA3

• Molecule 4: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain H: 67% 33%

NAG1 NAG2 BMA3

• Molecule 4: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain I: 33% 67%

NAG1 NAG2 BMA3

• Molecule 4: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain K: 67% 33%

NAG1 NAG2 BMA3

• Molecule 4: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain L: 100%

NAG1 NAG2 BMA3

4 Data and refinement statistics (i)

Property	Value	Source
Space group	I 4 3 2	Depositor
Cell constants a, b, c, α, β, γ	262.33Å 262.33Å 262.33Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	45.00 – 3.50 47.89 – 3.50	Depositor EDS
% Data completeness (in resolution range)	100.0 (45.00-3.50) 100.0 (47.89-3.50)	Depositor EDS
R_{merge}	0.03	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	1.00 (at 3.48Å)	Xtriage
Refinement program	BUSTER-TNT 2.10.1	Depositor
R, R_{free}	0.211 , 0.231 0.231 , 0.242	Depositor DCC
R_{free} test set	1003 reflections (5.09%)	wwPDB-VP
Wilson B-factor (Å^2)	170.4	Xtriage
Anisotropy	0.000	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.34 , 162.9	EDS
L-test for twinning[2]	$< \|L\| > = 0.46$, $< L^2 > = 0.29$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o,F_c correlation	0.93	EDS
Total number of atoms	6102	wwPDB-VP
Average B, all atoms (Å^2)	185.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.39% of the height of the origin peak. No significant pseudotranslation is detected.*

[1] Intensities estimated from amplitudes.

[2] Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: MAN, BMA, NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\|>5$	RMSZ	$\#\|Z\|>5$
1	A	0.37	0/2867	0.59	0/3871
1	B	0.36	0/2784	0.60	0/3756
All	All	0.36	0/5651	0.59	0/7627

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2800	0	2665	14	0
1	B	2720	0	2589	15	0
2	C	50	0	43	0	0
2	D	50	0	43	0	0
2	J	50	0	43	0	0
2	M	50	0	43	0	0
2	N	50	0	43	0	0
3	E	28	0	25	0	0
4	F	39	0	34	0	0
4	G	39	0	34	0	0
4	H	39	0	34	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	I	39	0	34	0	0
4	K	39	0	34	0	0
4	L	39	0	34	0	0
5	A	42	0	39	0	0
5	B	28	0	26	0	0
All	All	6102	0	5763	28	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (28) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:273:SER:HB2	1:A:279:GLU:HA	1.74	0.69
1:B:167:THR:HG22	1:B:231:GLN:HG2	1.78	0.66
1:A:167:THR:HG22	1:A:231:GLN:HG2	1.79	0.64
1:A:77:VAL:HG21	1:A:299:PHE:HZ	1.65	0.60
1:A:110:LEU:HB3	1:A:230:ILE:HG12	1.93	0.50
1:B:110:LEU:HB3	1:B:230:ILE:HG12	1.93	0.50
1:A:302:THR:HB	1:A:306:LYS:HE3	1.95	0.48
1:B:414:ASP:HA	1:B:417:ILE:HD12	1.96	0.47
1:B:128:ALA:CB	1:B:162:PHE:HB2	2.45	0.47
1:A:128:ALA:CB	1:A:162:PHE:HB2	2.45	0.46
1:A:152:ASN:HA	1:B:265:ALA:HA	1.97	0.45
1:A:172:LEU:HB3	1:A:188:ARG:HG3	1.99	0.45
1:A:117:ILE:HD13	1:A:229:ILE:HG21	1.98	0.45
1:B:117:ILE:HD13	1:B:229:ILE:HG21	1.98	0.45
1:B:392:TRP:CD1	1:B:399:TYR:HB3	2.52	0.44
1:B:172:LEU:HB3	1:B:188:ARG:HG3	1.99	0.44
1:A:108:LEU:HD13	1:A:138:LEU:HD22	2.00	0.43
1:B:88:MET:HB3	1:B:244:PRO:HB3	2.01	0.43
1:B:108:LEU:HD13	1:B:138:LEU:HD22	2.00	0.43
1:A:229:ILE:HG22	1:A:231:GLN:HG3	2.02	0.42
1:B:77:VAL:HG21	1:B:299:PHE:HZ	1.84	0.42
1:A:392:TRP:CD1	1:A:399:TYR:HB3	2.55	0.41
1:A:361:LEU:O	1:A:365:MET:HG2	2.21	0.41
1:B:77:VAL:HG21	1:B:299:PHE:CZ	2.56	0.41
1:A:247:MET:O	1:A:251:LEU:HG	2.21	0.40
1:B:229:ILE:HG22	1:B:231:GLN:HG3	2.02	0.40
1:B:247:MET:O	1:B:251:LEU:HG	2.22	0.40
1:B:312:ASP:C	1:B:355:LEU:HD11	2.42	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	340/438 (78%)	313 (92%)	26 (8%)	1 (0%)	41	75
1	B	329/438 (75%)	309 (94%)	19 (6%)	1 (0%)	41	75
All	All	669/876 (76%)	622 (93%)	45 (7%)	2 (0%)	41	75

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	396	ASN
1	A	281	PRO

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	311/384 (81%)	299 (96%)	12 (4%)	32	64
1	B	300/384 (78%)	288 (96%)	12 (4%)	31	64
All	All	611/768 (80%)	587 (96%)	24 (4%)	32	64

All (24) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	71	VAL

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	84	LEU
1	A	90	ASN
1	A	110	LEU
1	A	123	CYS
1	A	137	THR
1	A	184	CYS
1	A	239	CYS
1	A	267	THR
1	A	296	LEU
1	A	304	VAL
1	A	332	PHE
1	B	71	VAL
1	B	84	LEU
1	B	90	ASN
1	B	110	LEU
1	B	123	CYS
1	B	137	THR
1	B	184	CYS
1	B	239	CYS
1	B	296	LEU
1	B	332	PHE
1	B	396	ASN
1	B	418	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	136	HIS
1	B	415	ASN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

40 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
2	NAG	C	1	2,1	14,14,15	0.28	0	17,19,21	0.69	0
2	NAG	C	2	2	14,14,15	0.28	0	17,19,21	0.55	0
2	BMA	C	3	2	11,11,12	0.39	0	15,15,17	1.15	1 (6%)
2	MAN	C	4	2	11,11,12	0.37	0	15,15,17	0.99	1 (6%)
2	NAG	D	1	2,1	14,14,15	0.27	0	17,19,21	0.62	0
2	NAG	D	2	2	14,14,15	0.26	0	17,19,21	1.07	2 (11%)
2	BMA	D	3	2	11,11,12	0.40	0	15,15,17	0.89	0
2	MAN	D	4	2	11,11,12	0.39	0	15,15,17	1.18	1 (6%)
3	NAG	E	1	3,1	14,14,15	0.29	0	17,19,21	0.89	1 (5%)
3	NAG	E	2	3	14,14,15	0.32	0	17,19,21	0.65	1 (5%)
4	NAG	F	1	1,4	14,14,15	0.27	0	17,19,21	0.63	0
4	NAG	F	2	4	14,14,15	0.26	0	17,19,21	0.58	0
4	BMA	F	3	4	11,11,12	0.33	0	15,15,17	0.51	0
4	NAG	G	1	1,4	14,14,15	0.25	0	17,19,21	0.63	0
4	NAG	G	2	4	14,14,15	0.29	0	17,19,21	0.57	0
4	BMA	G	3	4	11,11,12	0.34	0	15,15,17	0.50	0
4	NAG	H	1	1,4	14,14,15	0.26	0	17,19,21	0.89	2 (11%)
4	NAG	H	2	4	14,14,15	0.28	0	17,19,21	0.68	0
4	BMA	H	3	4	11,11,12	0.34	0	15,15,17	0.64	0
4	NAG	I	1	1,4	14,14,15	0.26	0	17,19,21	0.90	2 (11%)
4	NAG	I	2	4	14,14,15	0.27	0	17,19,21	0.68	0
4	BMA	I	3	4	11,11,12	0.34	0	15,15,17	0.65	1 (6%)
2	NAG	J	1	2,1	14,14,15	0.30	0	17,19,21	0.67	0
2	NAG	J	2	2	14,14,15	0.30	0	17,19,21	0.50	0
2	BMA	J	3	2	11,11,12	0.58	0	15,15,17	1.46	3 (20%)
2	MAN	J	4	2	11,11,12	0.43	0	15,15,17	0.98	1 (6%)
4	NAG	K	1	1,4	14,14,15	0.28	0	17,19,21	0.67	0
4	NAG	K	2	4	14,14,15	0.29	0	17,19,21	1.13	2 (11%)
4	BMA	K	3	4	11,11,12	0.39	0	15,15,17	0.58	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
4	NAG	L	1	1,4	14,14,15	0.29	0	17,19,21	0.89	1 (5%)
4	NAG	L	2	4	14,14,15	0.35	0	17,19,21	0.61	1 (5%)
4	BMA	L	3	4	11,11,12	0.39	0	15,15,17	0.76	1 (6%)
2	NAG	M	1	2,1	14,14,15	0.28	0	17,19,21	0.64	1 (5%)
2	NAG	M	2	2	14,14,15	0.27	0	17,19,21	0.69	1 (5%)
2	BMA	M	3	2	11,11,12	0.33	0	15,15,17	0.56	0
2	MAN	M	4	2	11,11,12	0.39	0	15,15,17	0.83	1 (6%)
2	NAG	N	1	2,1	14,14,15	0.27	0	17,19,21	0.63	0
2	NAG	N	2	2	14,14,15	0.29	0	17,19,21	0.55	0
2	BMA	N	3	2	11,11,12	0.35	0	15,15,17	0.57	0
2	MAN	N	4	2	11,11,12	0.36	0	15,15,17	0.87	1 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	C	1	2,1	-	0/6/23/26	0/1/1/1
2	NAG	C	2	2	-	2/6/23/26	0/1/1/1
2	BMA	C	3	2	-	0/2/19/22	0/1/1/1
2	MAN	C	4	2	-	2/2/19/22	0/1/1/1
2	NAG	D	1	2,1	-	0/6/23/26	0/1/1/1
2	NAG	D	2	2	-	1/6/23/26	0/1/1/1
2	BMA	D	3	2	-	0/2/19/22	0/1/1/1
2	MAN	D	4	2	-	1/2/19/22	0/1/1/1
3	NAG	E	1	3,1	-	0/6/23/26	0/1/1/1
3	NAG	E	2	3	-	0/6/23/26	0/1/1/1
4	NAG	F	1	1,4	-	0/6/23/26	0/1/1/1
4	NAG	F	2	4	-	1/6/23/26	0/1/1/1
4	BMA	F	3	4	-	0/2/19/22	0/1/1/1
4	NAG	G	1	1,4	-	0/6/23/26	0/1/1/1
4	NAG	G	2	4	-	2/6/23/26	0/1/1/1
4	BMA	G	3	4	-	0/2/19/22	0/1/1/1
4	NAG	H	1	1,4	-	0/6/23/26	0/1/1/1
4	NAG	H	2	4	-	0/6/23/26	0/1/1/1
4	BMA	H	3	4	-	0/2/19/22	0/1/1/1
4	NAG	I	1	1,4	-	0/6/23/26	0/1/1/1
4	NAG	I	2	4	-	0/6/23/26	0/1/1/1
4	BMA	I	3	4	-	0/2/19/22	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	J	1	2,1	-	0/6/23/26	0/1/1/1
2	NAG	J	2	2	-	2/6/23/26	0/1/1/1
2	BMA	J	3	2	-	0/2/19/22	0/1/1/1
2	MAN	J	4	2	-	2/2/19/22	0/1/1/1
4	NAG	K	1	1,4	-	0/6/23/26	0/1/1/1
4	NAG	K	2	4	-	1/6/23/26	0/1/1/1
4	BMA	K	3	4	-	0/2/19/22	0/1/1/1
4	NAG	L	1	1,4	-	0/6/23/26	0/1/1/1
4	NAG	L	2	4	-	0/6/23/26	0/1/1/1
4	BMA	L	3	4	-	0/2/19/22	0/1/1/1
2	NAG	M	1	2,1	-	0/6/23/26	0/1/1/1
2	NAG	M	2	2	-	1/6/23/26	0/1/1/1
2	BMA	M	3	2	-	0/2/19/22	0/1/1/1
2	MAN	M	4	2	-	0/2/19/22	0/1/1/1
2	NAG	N	1	2,1	-	0/6/23/26	0/1/1/1
2	NAG	N	2	2	-	2/6/23/26	0/1/1/1
2	BMA	N	3	2	-	0/2/19/22	0/1/1/1
2	MAN	N	4	2	-	0/2/19/22	0/1/1/1

There are no bond length outliers.

All (25) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
2	J	3	BMA	C1-C2-C3	4.31	114.97	109.67
2	D	4	MAN	C1-O5-C5	3.66	117.14	112.19
4	K	2	NAG	O5-C1-C2	-3.55	105.67	111.29
2	C	4	MAN	C1-O5-C5	3.26	116.61	112.19
2	C	3	BMA	C1-C2-C3	3.24	113.65	109.67
2	N	4	MAN	C1-O5-C5	2.97	116.22	112.19
2	D	2	NAG	O5-C1-C2	-2.84	106.81	111.29
2	J	4	MAN	C1-O5-C5	2.80	115.99	112.19
4	I	1	NAG	O5-C1-C2	-2.53	107.29	111.29
4	H	1	NAG	O5-C1-C2	-2.49	107.36	111.29
4	L	1	NAG	O5-C1-C2	-2.48	107.38	111.29
3	E	1	NAG	O5-C1-C2	-2.44	107.44	111.29
4	I	1	NAG	C1-O5-C5	2.38	115.42	112.19
2	J	3	BMA	C3-C4-C5	2.31	114.36	110.24
2	M	4	MAN	C1-O5-C5	2.31	115.33	112.19
3	E	2	NAG	C1-O5-C5	2.31	115.32	112.19
4	H	1	NAG	C1-O5-C5	2.30	115.31	112.19
4	L	3	BMA	C1-O5-C5	2.30	115.31	112.19

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
2	D	2	NAG	C1-C2-N2	2.30	114.41	110.49
2	J	3	BMA	C2-C3-C4	2.28	114.84	110.89
4	L	2	NAG	C1-O5-C5	2.21	115.18	112.19
4	K	2	NAG	C1-C2-N2	2.18	114.21	110.49
2	M	2	NAG	C1-O5-C5	2.14	115.09	112.19
2	M	1	NAG	C1-O5-C5	2.03	114.94	112.19
4	I	3	BMA	C1-O5-C5	2.01	114.92	112.19

There are no chirality outliers.

All (17) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	C	4	MAN	O5-C5-C6-O6
2	C	2	NAG	O5-C5-C6-O6
2	J	4	MAN	O5-C5-C6-O6
2	J	2	NAG	O5-C5-C6-O6
2	C	4	MAN	C4-C5-C6-O6
2	J	4	MAN	C4-C5-C6-O6
2	N	2	NAG	O5-C5-C6-O6
4	G	2	NAG	O5-C5-C6-O6
4	K	2	NAG	O5-C5-C6-O6
2	C	2	NAG	C4-C5-C6-O6
4	F	2	NAG	O5-C5-C6-O6
2	D	2	NAG	O5-C5-C6-O6
2	M	2	NAG	O5-C5-C6-O6
2	J	2	NAG	C4-C5-C6-O6
2	D	4	MAN	O5-C5-C6-O6
2	N	2	NAG	C4-C5-C6-O6
4	G	2	NAG	C4-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.

Oligosaccharide Chain C



Bond lengths



Bond angles



Torsions



Rings

Oligosaccharide Chain D



Bond lengths



Bond angles



Torsions



Rings

Oligosaccharide Chain J



Bond lengths



Bond angles



Torsions



Rings

Oligosaccharide Chain M



Bond lengths



Bond angles



Torsions



Rings

Oligosaccharide Chain N



Oligosaccharide Chain E



Oligosaccharide Chain F



Bond lengths



Bond angles



Torsions



Rings

Oligosaccharide Chain G



Bond lengths



Bond angles



Torsions



Rings

Oligosaccharide Chain H



Oligosaccharide Chain I



Oligosaccharide Chain K



Oligosaccharide Chain L



5.6 Ligand geometry (i)

5 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
5	NAG	B	511	1	14,14,15	0.30	0	17,19,21	0.59	0
5	NAG	A	522	1	14,14,15	0.28	0	17,19,21	0.66	0
5	NAG	A	509	1	14,14,15	0.29	0	17,19,21	0.57	0
5	NAG	B	523	1	14,14,15	0.28	0	17,19,21	0.63	0
5	NAG	A	521	1	14,14,15	0.29	0	17,19,21	0.64	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	NAG	B	511	1	-	0/6/23/26	0/1/1/1
5	NAG	A	522	1	-	0/6/23/26	0/1/1/1
5	NAG	A	509	1	-	0/6/23/26	0/1/1/1
5	NAG	B	523	1	-	0/6/23/26	0/1/1/1
5	NAG	A	521	1	-	0/6/23/26	0/1/1/1

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
5	A	521	NAG	C1-O5-C5	2.08	115.02	112.19

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å^2)	Q<0.9
1	A	348/438 (79%)	0.47	21 (6%)	21	19	126, 169, 220, 261	0
1	B	339/438 (77%)	0.45	19 (5%)	24	22	134, 186, 236, 253	0
All	All	687/876 (78%)	0.46	40 (5%)	23	20	126, 176, 232, 261	0

All (40) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	183	GLN	7.5
1	A	217	THR	6.3
1	A	261	THR	5.5
1	A	254	GLN	5.4
1	B	254	GLN	5.2
1	A	253	ALA	4.8
1	B	264	LEU	4.8
1	A	257	THR	4.5
1	B	259	PHE	4.1
1	B	258	LYS	3.6
1	A	185	ARG	3.6
1	A	151	GLY	3.6
1	B	150	ARG	3.4
1	B	253	ALA	3.3
1	B	265	ALA	3.0
1	B	257	THR	3.0
1	A	152	ASN	3.0
1	A	113	THR	2.9
1	B	65	PRO	2.8
1	B	378	LEU	2.8
1	A	148	SER	2.8
1	A	169	GLN	2.4
1	B	78	GLU	2.4
1	B	392	TRP	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	229	ILE	2.3
1	A	373	SER	2.3
1	A	184	CYS	2.3
1	B	382	LYS	2.3
1	A	112	PHE	2.3
1	A	425	TYR	2.3
1	A	258	LYS	2.3
1	A	227	TYR	2.2
1	B	315	PHE	2.2
1	B	400	LEU	2.2
1	B	424	ASP	2.2
1	B	74	PHE	2.2
1	A	117	ILE	2.1
1	B	419	GLU	2.1
1	A	210	TRP	2.1
1	A	150	ARG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
2	MAN	J	4	11/12	0.18	0.37	265,266,267,267	0
4	BMA	G	3	11/12	0.55	0.29	214,222,225,228	0
2	MAN	D	4	11/12	0.61	0.39	264,264,265,265	0
2	MAN	M	4	11/12	0.68	0.47	257,263,264,265	0
2	MAN	N	4	11/12	0.71	0.39	234,240,243,245	0
4	BMA	L	3	11/12	0.72	0.49	254,254,255,255	0
4	NAG	H	2	14/15	0.74	0.44	223,226,229,232	0
4	BMA	I	3	11/12	0.74	0.38	254,257,261,263	0
2	BMA	D	3	11/12	0.74	0.26	259,262,263,264	0
3	NAG	E	2	14/15	0.75	0.47	249,251,252,253	0
2	NAG	J	2	14/15	0.76	0.32	235,246,251,254	0
4	NAG	I	2	14/15	0.76	0.41	243,245,249,252	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
2	BMA	J	3	11/12	0.77	0.37	256,259,263,265	0
4	NAG	K	2	14/15	0.78	0.35	232,234,240,245	0
2	BMA	M	3	11/12	0.78	0.25	250,254,260,261	0
3	NAG	E	1	14/15	0.80	0.46	231,236,241,245	0
2	NAG	C	2	14/15	0.83	0.23	239,247,251,252	0
4	BMA	H	3	11/12	0.83	0.27	233,234,237,237	0
4	BMA	F	3	11/12	0.84	0.16	220,223,224,225	0
4	NAG	L	2	14/15	0.84	0.52	244,250,253,254	0
2	NAG	M	2	14/15	0.84	0.19	233,235,241,245	0
2	BMA	N	3	11/12	0.85	0.31	226,237,239,240	0
4	BMA	K	3	11/12	0.86	0.41	242,250,254,254	0
2	NAG	C	1	14/15	0.88	0.30	239,242,246,246	0
4	NAG	H	1	14/15	0.88	0.43	205,210,219,222	0
4	NAG	I	1	14/15	0.88	0.29	229,231,236,240	0
4	NAG	G	2	14/15	0.90	0.33	187,194,205,209	0
4	NAG	L	1	14/15	0.90	0.30	231,234,238,241	0
4	NAG	F	1	14/15	0.90	0.20	194,197,208,211	0
2	BMA	C	3	11/12	0.90	0.23	250,252,255,256	0
4	NAG	F	2	14/15	0.91	0.18	213,215,217,218	0
2	NAG	D	1	14/15	0.91	0.20	230,235,239,242	0
2	NAG	M	1	14/15	0.91	0.17	211,217,224,229	0
2	NAG	N	2	14/15	0.91	0.32	203,208,217,220	0
2	NAG	D	2	14/15	0.91	0.20	247,252,257,260	0
2	MAN	C	4	11/12	0.92	0.18	256,257,260,262	0
4	NAG	G	1	14/15	0.92	0.35	166,170,177,183	0
2	NAG	J	1	14/15	0.92	0.29	234,237,240,240	0
4	NAG	K	1	14/15	0.93	0.21	216,221,225,227	0
2	NAG	N	1	14/15	0.93	0.39	181,187,192,198	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around Chain C:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around Chain D:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around Chain J:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around Chain M:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around Chain N:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around Chain E:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around Chain F:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around Chain G:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around Chain H:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around Chain I:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around Chain K:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around Chain L:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
5	NAG	A	522	14/15	0.64	0.42	218,227,235,236	0
5	NAG	B	511	14/15	0.65	0.36	228,234,239,241	0
5	NAG	A	521	14/15	0.70	0.47	245,249,250,251	0
5	NAG	A	509	14/15	0.77	0.27	230,235,236,236	0
5	NAG	B	523	14/15	0.80	0.22	213,226,235,235	0

6.5 Other polymers (i)

There are no such residues in this entry.