



Full wwPDB X-ray Structure Validation Report ⓘ

Jun 7, 2020 – 11:37 am BST

PDB ID : 6IO0
Title : Human IDH1 R132C mutant complexed with compound A.
Authors : Suzuki, M.; Baba, D.; Hanzawa, H.
Deposited on : 2018-10-29
Resolution : 2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4898 (2.20-2.20)
Clashscore	141614	5594 (2.20-2.20)
Ramachandran outliers	138981	5503 (2.20-2.20)
Sidechain outliers	138945	5504 (2.20-2.20)
RSRZ outliers	127900	4800 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	414	<p>4% 85% 11% .</p>
1	B	414	<p>3% 78% 18% .</p>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	GOL	B	502	-	-	X	-

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 6667 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Isocitrate dehydrogenase [NADP] cytoplasmic.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	400	Total 3080	C 1968	N 511	O 582	S 19	0	1	0
1	B	400	Total 3095	C 1975	N 514	O 587	S 19	0	1	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	132	CYS	ARG	engineered mutation	UNP O75874
B	132	CYS	ARG	engineered mutation	UNP O75874

- Molecule 2 is (2E)-3-{3-[3-(2,6-dichlorophenyl)-5-(propan-2-yl)-1,2-oxazole-4-carbonyl]-1-methyl-1H-indol-7-yl}prop-2-enoic acid (three-letter code: AOU) (formula: C₂₅H₂₀Cl₂N₂O₄).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
			Total	C	Cl	N			O
2	A	1	Total 33	C 25	Cl 2	N 2	O 4	0	0

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
			Total	C	Cl	N			O
2	B	1	33	25	2	2	4	0	0

- Molecule 3 is NADPH DIHYDRO-NICOTINAMIDE-ADENINE-DINUCLEOTIDE PHOSPHATE (three-letter code: NDP) (formula: $C_{21}H_{30}N_7O_{17}P_3$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
3	A	1	48	21	7	17	3	0	0
3	B	1	48	21	7	17	3	0	0

- Molecule 4 is CITRIC ACID (three-letter code: CIT) (formula: $C_6H_8O_7$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			13	6	7		
4	A	1	Total	C	O	0	0
			13	6	7		
4	B	1	Total	C	O	0	0
			13	6	7		

- Molecule 5 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			6	3	3		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total 6	C 3	O 3	0	0
5	B	1	Total 6	C 3	O 3	0	0
5	B	1	Total 6	C 3	O 3	0	0
5	B	1	Total 6	C 3	O 3	0	0
5	B	1	Total 6	C 3	O 3	0	0
5	B	1	Total 6	C 3	O 3	0	0


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	121	Total 121	O 121	0	0
6	B	128	Total 128	O 128	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Isocitrate dehydrogenase [NADP] cytoplasmic



- Molecule 1: Isocitrate dehydrogenase [NADP] cytoplasmic



4 Data and refinement statistics

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, α , β , γ	82.75Å 82.75Å 296.42Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	25.00 – 2.20 24.67 – 2.20	Depositor EDS
% Data completeness (in resolution range)	94.3 (25.00-2.20) 94.4 (24.67-2.20)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.75 (at 2.19Å)	Xtrriage
Refinement program	REFMAC 5.8.0232	Depositor
R, R_{free}	0.190 , 0.255 0.190 , 0.254	Depositor DCC
R_{free} test set	2526 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	40.3	Xtrriage
Anisotropy	0.042	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 43.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.45$, $\langle L^2 \rangle = 0.28$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	6667	wwPDB-VP
Average B, all atoms (Å ²)	43.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.20% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, AOU, CIT, NDP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.38	0/3148	0.74	0/4259
1	B	0.39	0/3163	0.77	4/4279 (0.1%)
All	All	0.38	0/6311	0.76	4/8538 (0.0%)

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	338	ARG	CG-CD-NE	6.89	126.26	111.80
1	B	338	ARG	NE-CZ-NH2	-6.70	116.95	120.30
1	B	119	ARG	NE-CZ-NH1	-5.36	117.62	120.30
1	B	338	ARG	NE-CZ-NH1	5.28	122.94	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3080	0	2968	33	0
1	B	3095	0	2980	62	0
2	A	33	0	0	0	0
2	B	33	0	0	2	0
3	A	48	0	26	3	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	B	48	0	26	2	0
4	A	26	0	10	3	0
4	B	13	0	5	1	0
5	A	12	0	16	2	0
5	B	30	0	40	7	0
6	A	121	0	0	3	0
6	B	128	0	0	4	0
All	All	6667	0	6071	93	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (93) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:29:LYS:NZ	1:A:399:ASP:OD1	1.67	1.27
1:A:109:ARG:NH1	1:A:279:ASP:OD1	1.95	1.00
1:B:112:ILE:HA	5:B:502:GOL:H12	1.47	0.94
3:A:502:NDP:H42N	4:A:503:CIT:H22	1.50	0.93
1:B:113:ILE:HG12	5:B:502:GOL:H11	1.51	0.92
1:B:129:ILE:HB	1:B:266:ILE:HD12	1.74	0.70
1:B:410:ALA:O	1:B:411:GLN:HG3	1.91	0.70
1:A:185:GLN:HG2	6:A:683:HOH:O	1.91	0.70
1:A:6:SER:HB3	5:A:505:GOL:H31	1.72	0.70
1:B:323:GLN:O	5:B:504:GOL:H32	1.91	0.70
1:A:17:GLU:HG2	1:A:316:TYR:CD2	2.34	0.63
1:B:116:ASN:HB3	6:B:702:HOH:O	2.00	0.62
1:B:158:PRO:HG2	1:B:163:GLN:HG2	1.82	0.62
1:A:210:SER:HA	1:A:249:ARG:O	2.01	0.61
1:A:284:GLY:HA2	1:B:259:MET:HE3	1.83	0.60
1:B:338:ARG:HG3	1:B:338:ARG:HH11	1.67	0.60
1:A:127:PRO:HD2	1:A:205:TRP:CH2	2.37	0.59
1:B:132:CYS:SG	1:B:276:VAL:HG21	2.43	0.58
1:B:113:ILE:H	5:B:502:GOL:C1	2.17	0.57
1:B:292[B]:THR:HG22	1:B:293:SER:H	1.68	0.57
1:A:194:HIS:O	1:A:198:GLN:HG2	2.03	0.57
3:B:507:NDP:H42N	4:B:508:CIT:H22	1.86	0.57
1:B:220:ASP:OD1	1:B:270:LYS:HE3	2.05	0.57
1:B:22:ILE:HD11	1:B:327:THR:HB	1.87	0.56
1:A:6:SER:CB	5:A:505:GOL:H31	2.35	0.56
1:B:106:THR:HG23	1:B:132:CYS:O	2.06	0.56

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:259:MET:HE3	2:B:506:AOU:C6	2.35	0.55
1:A:239:PHE:CD2	1:A:244:ILE:HG13	2.41	0.55
1:B:383:LEU:N	1:B:384:PRO:CD	2.69	0.55
1:A:12:GLU:HG2	1:A:13:MET:N	2.21	0.54
1:B:210:SER:HA	1:B:249:ARG:O	2.07	0.54
3:A:502:NDP:C4N	4:A:503:CIT:H22	2.32	0.53
1:A:79:ASP:H	1:A:82:ARG:HB2	1.74	0.53
1:A:127:PRO:HG2	1:A:205:TRP:HH2	1.75	0.52
1:A:163:GLN:HA	6:A:658:HOH:O	2.09	0.52
1:B:112:ILE:CA	5:B:502:GOL:H12	2.30	0.51
1:A:75:THR:O	3:A:502:NDP:H2N	2.11	0.51
1:A:338:ARG:HH11	1:A:338:ARG:HG3	1.75	0.51
1:B:361:GLU:OE1	1:B:408:LYS:NZ	2.32	0.51
1:B:224:LYS:HG3	1:B:248:HIS:CD2	2.46	0.50
1:B:233:LYS:C	1:B:234:GLN:HG2	2.32	0.50
1:B:113:ILE:H	5:B:502:GOL:H12	1.77	0.50
1:A:383:LEU:N	1:A:384:PRO:CD	2.75	0.49
1:B:158:PRO:HD2	1:B:163:GLN:O	2.12	0.49
1:B:132:CYS:SG	1:B:276:VAL:CG2	3.01	0.49
1:B:4:LYS:HB2	1:B:34:TYR:O	2.12	0.49
1:B:212:LYS:O	1:B:214:THR:N	2.46	0.49
1:B:397:PHE:CE2	1:B:401:LEU:HD11	2.47	0.48
1:B:32:PHE:N	1:B:33:PRO:CD	2.76	0.48
1:A:169:VAL:O	1:A:170:HIS:HB2	2.14	0.47
1:B:242:GLN:O	1:B:243:LYS:HB2	2.13	0.47
1:A:113:ILE:HD11	1:A:127:PRO:HA	1.96	0.47
1:B:334:PHE:O	1:B:338:ARG:HG2	2.15	0.47
1:A:284:GLY:HA2	1:B:259:MET:CE	2.44	0.46
1:A:318:MET:HE1	6:A:716:HOH:O	2.13	0.46
1:B:201:LEU:HA	1:B:201:LEU:HD23	1.77	0.46
1:B:224:LYS:HG3	1:B:248:HIS:CG	2.50	0.46
1:A:155:THR:HG22	6:B:636:HOH:O	2.16	0.46
1:A:32:PHE:N	1:A:33:PRO:CD	2.80	0.45
1:B:113:ILE:N	5:B:502:GOL:H12	2.31	0.45
1:B:376:LEU:O	1:B:379:CYS:HB2	2.16	0.45
1:A:219:TYR:HB2	1:B:143:ASP:HB2	1.97	0.45
1:B:180:MET:CE	1:B:182:MET:HG3	2.47	0.45
1:B:185:GLN:HB2	6:B:695:HOH:O	2.17	0.44
1:A:22:ILE:HD11	1:A:327:THR:HB	2.00	0.44
1:B:383:LEU:HB3	1:B:384:PRO:HD3	1.99	0.44
1:A:270:LYS:HE2	1:A:270:LYS:HB3	1.72	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:67:HIS:O	1:B:68:ASN:HB2	2.18	0.43
1:B:24:GLU:O	1:B:28:GLU:HG2	2.19	0.43
1:B:288:LEU:HG	1:B:309:HIS:HB3	2.00	0.43
1:B:75:THR:O	3:B:507:NDP:H2N	2.18	0.43
1:B:183:TYR:C	1:B:183:TYR:CD1	2.92	0.43
1:B:233:LYS:C	1:B:234:GLN:CG	2.86	0.43
1:B:129:ILE:HB	1:B:266:ILE:CD1	2.44	0.43
1:B:24:GLU:OE1	1:B:27:LYS:NZ	2.46	0.43
1:B:29:LYS:HE3	1:B:399:ASP:OD1	2.19	0.43
1:A:77:THR:OG1	4:A:503:CIT:O5	2.32	0.43
1:A:182:MET:HE3	1:A:182:MET:HB3	1.92	0.42
1:A:155:THR:O	1:B:152:VAL:HA	2.19	0.42
1:B:122:SER:HB2	6:B:685:HOH:O	2.19	0.42
1:B:233:LYS:O	1:B:234:GLN:HG2	2.19	0.42
1:A:209:LEU:HA	1:A:268:ALA:O	2.19	0.42
1:B:160:ASP:OD1	1:B:160:ASP:C	2.58	0.42
1:B:193:ALA:HB2	1:B:227:PHE:CE1	2.55	0.42
1:B:193:ALA:HA	1:B:227:PHE:CZ	2.56	0.41
1:B:23:TRP:CD2	1:B:73:CYS:HB2	2.56	0.41
1:A:78:PRO:HD2	1:A:92:TRP:O	2.21	0.41
1:B:246:TYR:CD1	1:B:246:TYR:C	2.94	0.41
1:B:68:ASN:O	1:B:302:THR:HA	2.21	0.41
1:B:259:MET:CE	2:B:506:AOU:C6	2.98	0.41
1:B:169:VAL:O	1:B:170:HIS:HB2	2.21	0.40
1:A:284:GLY:CA	1:B:259:MET:HE3	2.49	0.40
1:B:383:LEU:N	1:B:384:PRO:HD2	2.37	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	395/414 (95%)	382 (97%)	13 (3%)	0	100	100
1	B	395/414 (95%)	377 (95%)	17 (4%)	1 (0%)	41	46
All	All	790/828 (95%)	759 (96%)	30 (4%)	1 (0%)	51	60

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	213	ASN

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	314/350 (90%)	306 (98%)	8 (2%)	47	60
1	B	316/350 (90%)	309 (98%)	7 (2%)	52	65
All	All	630/700 (90%)	615 (98%)	15 (2%)	49	62

All (15) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	28	GLU
1	A	94	SER
1	A	199	MET
1	A	247	GLU
1	A	270	LYS
1	A	278	SER
1	A	279	ASP
1	A	312	VAL
1	B	65	LYS
1	B	133	HIS
1	B	184	ASN
1	B	237	SER
1	B	270	LYS
1	B	278	SER
1	B	338	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (11) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	171	ASN
1	A	198	GLN
1	A	283	GLN
1	A	404	ASN
1	B	14	GLN
1	B	90	GLN
1	B	163	GLN
1	B	234	GLN
1	B	283	GLN
1	B	323	GLN
1	B	404	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

14 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	GOL	B	503	-	5,5,5	0.11	0	5,5,5	0.35	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	CIT	A	503	-	3,12,12	0.61	0	3,17,17	0.84	0
3	NDP	A	502	-	45,52,52	0.78	0	53,80,80	0.96	1 (1%)
2	AOU	B	506	-	29,36,36	1.60	4 (13%)	32,53,53	1.61	4 (12%)
5	GOL	B	505	-	5,5,5	0.16	0	5,5,5	0.45	0
5	GOL	B	504	-	5,5,5	0.20	0	5,5,5	0.63	0
3	NDP	B	507	-	45,52,52	0.62	0	53,80,80	0.99	5 (9%)
5	GOL	A	506	-	5,5,5	0.10	0	5,5,5	0.36	0
5	GOL	A	505	-	5,5,5	0.12	0	5,5,5	0.40	0
5	GOL	B	501	-	5,5,5	0.08	0	5,5,5	0.47	0
5	GOL	B	502	-	5,5,5	0.17	0	5,5,5	0.55	0
4	CIT	A	504	-	3,12,12	0.48	0	3,17,17	1.64	1 (33%)
2	AOU	A	501	-	29,36,36	1.39	4 (13%)	32,53,53	1.43	4 (12%)
4	CIT	B	508	-	3,12,12	0.84	0	3,17,17	0.68	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	GOL	B	503	-	-	2/4/4/4	-
4	CIT	A	503	-	-	2/6/16/16	-
3	NDP	A	502	-	-	10/30/77/77	0/5/5/5
2	AOU	B	506	-	-	0/7/21/21	0/4/4/4
5	GOL	B	505	-	-	4/4/4/4	-
5	GOL	B	504	-	-	3/4/4/4	-
3	NDP	B	507	-	-	5/30/77/77	0/5/5/5
5	GOL	A	506	-	-	0/4/4/4	-
5	GOL	A	505	-	-	4/4/4/4	-
5	GOL	B	501	-	-	1/4/4/4	-
5	GOL	B	502	-	-	2/4/4/4	-
4	CIT	A	504	-	-	1/6/16/16	-
2	AOU	A	501	-	-	1/7/21/21	0/4/4/4
4	CIT	B	508	-	-	1/6/16/16	-

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	506	AOU	C1-C13	4.17	1.46	1.42

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	506	AOU	C10-C34	-3.65	1.46	1.50
2	A	501	AOU	C1-C20	-3.28	1.46	1.50
2	B	506	AOU	C10-C14	3.12	1.43	1.40
2	A	501	AOU	C1-C13	2.84	1.45	1.42
2	A	501	AOU	C40-C4	2.30	1.41	1.36
2	B	506	AOU	C1-C20	-2.11	1.47	1.50
2	A	501	AOU	C8-N1	2.07	1.41	1.38

All (15) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	506	AOU	C35-C36-C37	5.66	139.12	129.87
2	A	501	AOU	C35-C36-C37	4.24	136.80	129.87
2	A	501	AOU	C21-C19-C18	-3.16	116.93	123.69
4	A	504	CIT	C3-C2-C1	2.72	119.35	114.98
2	B	506	AOU	C39-C37-C36	2.71	115.65	110.96
3	B	507	NDP	C3N-C7N-N7N	-2.44	113.33	117.67
3	B	507	NDP	C3B-C2B-C1B	-2.36	98.46	102.89
3	A	502	NDP	O2B-P2B-O1X	-2.35	100.34	109.39
3	B	507	NDP	C5A-C6A-N6A	2.33	123.90	120.35
2	A	501	AOU	C14-C10-C15	2.26	118.43	116.05
2	B	506	AOU	C4-C13-C1	-2.24	132.43	135.63
3	B	507	NDP	O3D-C3D-C4D	-2.18	104.73	111.05
2	A	501	AOU	C10-C14-CL2	-2.17	116.96	119.74
3	B	507	NDP	O3B-C3B-C2B	2.15	117.26	111.17
2	B	506	AOU	C6-C15-C10	-2.01	120.21	122.35

There are no chirality outliers.

All (36) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	B	503	GOL	C1-C2-C3-O3
3	A	502	NDP	C5D-O5D-PN-O1N
3	A	502	NDP	C5D-O5D-PN-O2N
5	B	505	GOL	O1-C1-C2-O2
5	B	505	GOL	O1-C1-C2-C3
5	B	505	GOL	C1-C2-C3-O3
5	A	505	GOL	O1-C1-C2-C3
5	A	505	GOL	C1-C2-C3-O3
5	B	501	GOL	O1-C1-C2-C3
5	B	502	GOL	C1-C2-C3-O3
5	B	504	GOL	O1-C1-C2-C3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
5	B	505	GOL	O2-C2-C3-O3
5	B	504	GOL	O1-C1-C2-O2
5	A	505	GOL	O2-C2-C3-O3
5	B	502	GOL	O2-C2-C3-O3
5	B	503	GOL	O2-C2-C3-O3
5	A	505	GOL	O1-C1-C2-O2
3	A	502	NDP	O4B-C4B-C5B-O5B
5	B	504	GOL	O2-C2-C3-O3
3	A	502	NDP	C3B-C4B-C5B-O5B
3	A	502	NDP	C2D-C1D-N1N-C2N
4	A	503	CIT	C6-C3-C4-C5
4	B	508	CIT	C6-C3-C4-C5
3	A	502	NDP	O4D-C1D-N1N-C2N
3	B	507	NDP	O4D-C1D-N1N-C2N
3	B	507	NDP	C2D-C1D-N1N-C2N
4	A	504	CIT	C1-C2-C3-O7
3	B	507	NDP	C3B-C4B-C5B-O5B
3	A	502	NDP	O4D-C1D-N1N-C6N
3	A	502	NDP	C2B-O2B-P2B-O3X
3	A	502	NDP	C5D-O5D-PN-O3
3	B	507	NDP	C2B-O2B-P2B-O3X
4	A	503	CIT	O7-C3-C4-C5
3	B	507	NDP	C1B-C2B-O2B-P2B
2	A	501	AOU	C9-C11-C18-C19
3	A	502	NDP	C2D-C1D-N1N-C6N


There are no ring outliers.



8 monomers are involved in 17 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	503	CIT	3	0
3	A	502	NDP	3	0
2	B	506	AOU	2	0
5	B	504	GOL	1	0
3	B	507	NDP	2	0
5	A	505	GOL	2	0
5	B	502	GOL	6	0
4	B	508	CIT	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will

also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	400/414 (96%)	-0.05	17 (4%) 35 33	27, 41, 69, 106	0
1	B	400/414 (96%)	-0.14	11 (2%) 53 51	23, 39, 66, 100	0
All	All	800/828 (96%)	-0.09	28 (3%) 44 42	23, 41, 69, 106	0

All (28) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	241	ALA	4.8
1	A	276	VAL	4.5
1	B	214	THR	3.8
1	B	213	ASN	3.8
1	B	215	ILE	3.7
1	B	412	ALA	3.2
1	A	283	GLN	2.8
1	B	410	ALA	2.7
1	A	215	ILE	2.7
1	B	207	LEU	2.7
1	A	277	GLN	2.7
1	A	413	LYS	2.5
1	B	159	SER	2.5
1	A	278	SER	2.4
1	B	237	SER	2.4
1	A	209	LEU	2.4
1	A	321	LYS	2.3
1	B	348	ASN	2.2
1	A	333	ILE	2.2
1	B	133	HIS	2.2
1	A	208	TYR	2.2
1	A	322	GLY	2.2
1	A	286	GLY	2.1
1	B	3	LYS	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	268	ALA	2.1
1	A	330	ILE	2.1
1	A	329	PRO	2.0
1	A	227	PHE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)



In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	CIT	A	504	13/13	0.64	0.32	56,80,103,115	0
5	GOL	B	504	6/6	0.73	0.22	58,63,71,72	0
5	GOL	A	506	6/6	0.76	0.12	69,81,82,85	0
5	GOL	B	505	6/6	0.81	0.17	60,74,75,77	0
5	GOL	B	502	6/6	0.83	0.18	55,61,64,64	0
5	GOL	B	503	6/6	0.89	0.21	37,64,67,71	0
5	GOL	A	505	6/6	0.90	0.17	49,64,72,74	0
4	CIT	B	508	13/13	0.90	0.20	67,78,87,111	0
4	CIT	A	503	13/13	0.91	0.29	89,93,96,97	0
3	NDP	A	502	48/48	0.92	0.15	45,59,69,80	0
5	GOL	B	501	6/6	0.95	0.21	57,60,61,61	0
2	AOU	B	506	33/33	0.95	0.09	31,41,48,65	0
2	AOU	A	501	33/33	0.96	0.11	40,50,58,64	0
3	NDP	B	507	48/48	0.99	0.07	25,31,35,38	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around NDP A 502:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around AOU B 506:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around AOU A 501:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around NDP B 507:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers

There are no such residues in this entry.