



# wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 18, 2024 – 02:10 PM JST

PDB ID : 8IOF  
Title : Crystal structure of N-methyl-Cis-4-hydroxy-D-proline dehydratase in Clostridium sp. FS41  
Authors : Jiang, L.; Zhang, Y.  
Deposited on : 2023-03-11  
Resolution : 2.33 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.36  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.33 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	2096 (2.36-2.32)
Clashscore	141614	2193 (2.36-2.32)
Ramachandran outliers	138981	2159 (2.36-2.32)
Sidechain outliers	138945	2160 (2.36-2.32)
RSRZ outliers	127900	2067 (2.36-2.32)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	793	
1	B	793	
1	C	793	
1	D	793	

## 2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 26256 atoms, of which 40 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Benzylsuccinate synthase alpha subunit.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	791	Total 6224	C 3924	N 1049	O 1200	S 51	0	0	0
1	B	791	Total 6224	C 3924	N 1049	O 1200	S 51	0	0	0
1	C	792	Total 6230	C 3927	N 1050	O 1202	S 51	0	0	0
1	D	791	Total 6224	C 3924	N 1049	O 1200	S 51	0	0	0

- Molecule 2 is (2R,4R)-1-methyl-4-hydroxyl-pyrrolidine-2-carboxylic acid (three-letter code: 87I) (formula: C<sub>6</sub>H<sub>11</sub>NO<sub>3</sub>) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	H	N	O		
2	A	1	Total 20	C 6	H 10	N 1	O 3	0	0
2	B	1	Total 20	C 6	H 10	N 1	O 3	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	C	1	Total	C	H	N	O	0	0
			20	6	10	1	3		
2	D	1	Total	C	H	N	O	0	0
			20	6	10	1	3		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	345	Total	O	0	0
			345	345		
3	B	319	Total	O	0	0
			319	319		
3	C	315	Total	O	0	0
			315	315		
3	D	295	Total	O	0	0
			295	295		

### 3 Residue-property plots i

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Benzylsuccinate synthase alpha subunit



- Molecule 1: Benzylsuccinate synthase alpha subunit











## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	78.97Å 103.60Å 114.25Å 83.13° 75.71° 79.84°	Depositor
Resolution (Å)	39.43 – 2.33 39.49 – 2.33	Depositor EDS
% Data completeness (in resolution range)	96.0 (39.43-2.33) 96.3 (39.49-2.33)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.83 (at 2.34Å)	Xtrriage
Refinement program	PHENIX (1.14_3247: ???)	Depositor
R, $R_{free}$	0.190 , 0.266 0.191 , 0.266	Depositor DCC
$R_{free}$ test set	2000 reflections (1.38%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	9.2	Xtrriage
Anisotropy	0.729	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 34.6	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.85	EDS
Total number of atoms	26256	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	13.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.47% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 87I

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.54	3/6346 (0.0%)	0.62	3/8571 (0.0%)
1	B	0.66	11/6346 (0.2%)	0.64	2/8571 (0.0%)
1	C	0.45	0/6352	0.58	0/8579
1	D	0.51	1/6346 (0.0%)	0.60	1/8571 (0.0%)
All	All	0.55	15/25390 (0.1%)	0.61	6/34292 (0.0%)

The worst 5 of 15 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	269	GLU	CD-OE2	-8.72	1.16	1.25
1	B	268	TYR	CZ-OH	-6.90	1.26	1.37
1	B	43	TRP	CE3-CZ3	-6.58	1.27	1.38
1	B	269	GLU	CD-OE1	-6.39	1.18	1.25
1	D	165	GLY	C-O	-6.19	1.13	1.23

The worst 5 of 6 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	167	ARG	NE-CZ-NH1	14.46	127.53	120.30
1	A	680	ASP	O-C-N	10.35	139.26	122.70
1	B	167	ARG	NE-CZ-NH2	-8.81	115.89	120.30
1	A	680	ASP	CA-C-N	-8.07	99.44	117.20
1	D	287	ASP	CB-CG-OD1	6.98	124.58	118.30

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	6224	0	6107	286	0
1	B	6224	0	6107	328	0
1	C	6230	0	6112	278	0
1	D	6224	0	6107	349	0
2	A	10	10	0	2	0
2	B	10	10	0	0	0
2	C	10	10	0	1	0
2	D	10	10	0	2	0
3	A	345	0	0	25	1
3	B	319	0	0	51	0
3	C	315	0	0	43	1
3	D	295	0	0	54	0
All	All	26216	40	24433	1234	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 25.

The worst 5 of 1234 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:342:GLN:HE21	1:D:378:THR:HG21	1.06	1.10
1:B:459:VAL:CG2	1:B:561:ILE:HD12	1.84	1.08
1:C:274:MET:SD	3:C:1193:HOH:O	2.13	1.06
2:D:801:87I:O02	3:D:902:HOH:O	1.74	1.05
1:D:411:ASP:OD1	3:D:903:HOH:O	1.75	1.05

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:1122:HOH:O	3:C:1184:HOH:O[1_556]	2.04	0.16

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	789/793 (100%)	753 (95%)	31 (4%)	5 (1%)	25	26
1	B	789/793 (100%)	749 (95%)	34 (4%)	6 (1%)	19	20
1	C	790/793 (100%)	743 (94%)	42 (5%)	5 (1%)	25	26
1	D	789/793 (100%)	737 (93%)	48 (6%)	4 (0%)	29	31
All	All	3157/3172 (100%)	2982 (94%)	155 (5%)	20 (1%)	25	26

5 of 20 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	158	GLY
1	B	158	GLY
1	C	158	GLY
1	C	434	GLY
1	C	474	GLN

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	665/667 (100%)	648 (97%)	17 (3%)	46	56
1	B	665/667 (100%)	643 (97%)	22 (3%)	38	46
1	C	666/667 (100%)	654 (98%)	12 (2%)	59	70
1	D	665/667 (100%)	652 (98%)	13 (2%)	55	66
All	All	2661/2668 (100%)	2597 (98%)	64 (2%)	49	59

5 of 64 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	296	LYS
1	D	385	LYS
1	B	339	MET
1	B	316	TRP
1	D	460	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 15 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	739	GLN
1	D	739	GLN
1	D	342	GLN
1	D	744	ASN
1	D	704	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	87I	D	801	-	9,10,10	6.17	3 (33%)	12,14,14	1.65	2 (16%)
2	87I	C	801	-	9,10,10	6.16	4 (44%)	12,14,14	1.38	2 (16%)
2	87I	B	801	-	9,10,10	6.54	4 (44%)	12,14,14	1.34	2 (16%)
2	87I	A	801	-	9,10,10	6.40	4 (44%)	12,14,14	0.99	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	87I	D	801	-	-	2/4/16/16	0/1/1/1
2	87I	C	801	-	-	0/4/16/16	0/1/1/1
2	87I	B	801	-	-	4/4/16/16	0/1/1/1
2	87I	A	801	-	-	4/4/16/16	0/1/1/1

The worst 5 of 15 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	801	87I	C08-N04	13.97	1.66	1.47
2	A	801	87I	C08-N04	13.74	1.65	1.47
2	D	801	87I	C08-N04	13.01	1.64	1.47
2	C	801	87I	C08-N04	12.92	1.64	1.47
2	B	801	87I	C08-C07	-12.91	1.33	1.52

The worst 5 of 6 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	801	87I	C06-C07-C08	4.01	108.24	103.31
2	D	801	87I	C07-C06-C05	-2.93	100.22	103.97
2	C	801	87I	C06-C07-C08	2.62	106.53	103.31
2	C	801	87I	C09-N04-C08	-2.50	109.06	112.45
2	B	801	87I	C06-C07-C08	-2.14	100.67	103.31

There are no chirality outliers.

5 of 10 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	801	87I	N04-C05-C10-O02
2	B	801	87I	N04-C05-C10-O03

*Continued on next page...*

*Continued from previous page...*


Mol	Chain	Res	Type	Atoms
2	A	801	87I	N04-C05-C10-O02
2	A	801	87I	N04-C05-C10-O03
2	B	801	87I	C06-C05-C10-O03

There are no ring outliers.


3 monomers are involved in 5 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	D	801	87I	2	0
2	C	801	87I	1	0
2	A	801	87I	2	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.











## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	791/793 (99%)	0.31	19 (2%) 59 68	3, 10, 25, 43	0
1	B	791/793 (99%)	0.35	15 (1%) 66 75	3, 11, 24, 37	0
1	C	792/793 (99%)	0.40	27 (3%) 45 55	4, 12, 25, 44	0
1	D	791/793 (99%)	0.36	23 (2%) 51 61	4, 12, 28, 41	0
All	All	3165/3172 (99%)	0.36	84 (2%) 54 64	3, 11, 26, 44	0

The worst 5 of 84 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	2	SER	4.5
1	B	268	TYR	3.9
1	B	166	TRP	3.7
1	B	232	ALA	3.6
1	C	232	ALA	3.6

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)


There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	87I	C	801	10/10	0.83	0.26	18,28,36,40	0
2	87I	D	801	10/10	0.88	0.22	23,35,42,44	0
2	87I	B	801	10/10	0.91	0.19	22,26,34,34	0
2	87I	A	801	10/10	0.92	0.20	15,22,28,28	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






**Electron density around 87I D 801:**


$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



**Electron density around 87I B 801:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





## 6.5 Other polymers [i](#)

There are no such residues in this entry.