

Full wwPDB X-ray Structure Validation Report (i)

Mar 18, 2024 – 05:15 PM JST

PDB ID : 8IT7

Title: Phosphoglycerate mutase 1 complexed with a compound

Authors : Zhou, L.; Jiang, L.L.

Deposited on : 2023-03-22

Resolution : 2.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity : 4.02b-467

Mogul : 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : 1.13

EDS : 2.36

buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

Refmac : 5.8.0158

CCP4 : 7.0.044 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.36

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X- $RAY\ DIFFRACTION$

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive $(\# \mathrm{Entries})$	$\begin{array}{c} {\rm Similar\ resolution} \\ (\#{\rm Entries},{\rm resolution\ range}({\rm \AA})) \end{array}$
R_{free}	130704	3140 (2.80-2.80)
Clashscore	141614	3569 (2.80-2.80)
Ramachandran outliers	138981	3498 (2.80-2.80)
Sidechain outliers	138945	3500 (2.80-2.80)
RSRZ outliers	127900	3078 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain		
1	В	262	63%	26%	11%
1	С	262	68%	23%	10%



2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 3851 atoms, of which 17 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Phosphoglycerate mutase 1.

\mathbf{Mol}	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace		
1	В	233	Total	_	N	О	S	0	1	0
	Б	200	1882	1199	332	345	6		1	
1	C	237	Total	С	N	Ο	S	0	0	0
1		231	1906	1212	336	352	6		U	

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
В	255	LEU	-	expression tag	UNP P18669
В	256	GLU	-	expression tag	UNP P18669
В	257	HIS	-	expression tag	UNP P18669
В	258	HIS	-	expression tag	UNP P18669
В	259	HIS	-	expression tag	UNP P18669
В	260	HIS	-	expression tag	UNP P18669
В	261	HIS	-	expression tag	UNP P18669
В	262	HIS	-	expression tag	UNP P18669
С	255	LEU	-	expression tag	UNP P18669
С	256	GLU	-	expression tag	UNP P18669
С	257	HIS	-	expression tag	UNP P18669
С	258	HIS	-	expression tag	UNP P18669
С	259	HIS	-	expression tag	UNP P18669
С	260	HIS	-	expression tag	UNP P18669
С	261	HIS	-	expression tag	UNP P18669
С	262	HIS	-	expression tag	UNP P18669


• Molecule 2 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	С	1	Total Cl 1 1	0	0

• Molecule 3 is 2-(N-MORPHOLINO)-ETHANESULFONIC ACID (three-letter code: MES)




(formula: $C_6H_{13}NO_4S$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf		
3	С	1	Total 12	C 6	N 1	O 4	S 1	0	0

• Molecule 4 is 4-(4-chlorophenyl)-2-oxidanyl-6-[(4-phenylphenyl)sulfonylamino]benzoic acid (three-letter code: Q9F) (formula: $C_{25}H_{18}ClNO_5S$) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf				
4	С	1		С	Cl	Н	N	О	S	0	0
1		1	50	25	1	17	1	5	1		V




3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Phosphoglycerate mutase 1



• Molecule 1: Phosphoglycerate mutase 1





4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants	82.19Å 85.59Å 101.30Å	Donogitor
a, b, c, α , β , γ	90.00° 90.00° 90.00°	Depositor
Resolution (Å)	31.91 - 2.80	Depositor
Resolution (A)	31.91 - 2.80	EDS
% Data completeness	98.5 (31.91-2.80)	Depositor
(in resolution range)	98.5 (31.91-2.80)	EDS
R_{merge}	0.24	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	1.63 (at 2.81Å)	Xtriage
Refinement program	PHENIX (1.19.2_4158: ???)	Depositor
D.D.	0.192 , 0.249	Depositor
R, R_{free}	0.192 , 0.248	DCC
R_{free} test set	849 reflections (4.75%)	wwPDB-VP
Wilson B-factor (Å ²)	59.3	Xtriage
Anisotropy	0.503	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.34, 33.4	EDS
L-test for twinning ²	$< L > = 0.49, < L^2> = 0.32$	Xtriage
Estimated twinning fraction	0.019 for k,h,-l	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	3851	wwPDB-VP
Average B, all atoms (\mathring{A}^2)	54.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 4.72% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of <|L|>, $<L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: Q9F, MES, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond angles		
MIOI	Chain	RMSZ	# Z > 5	RMSZ	# Z > 5	
1	В	0.49	0/1934	0.66	0/2624	
1	С	0.43	0/1955	0.62	0/2653	
All	All	0.47	0/3889	0.64	0/5277	

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	В	1882	0	1860	50	0
1	С	1906	0	1878	39	0
2	С	1	0	0	0	0
3	С	12	0	13	0	0
4	С	33	17	0	2	0
All	All	3834	17	3751	86	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

All (86) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.



Atom-1	Atom-2	Interatomic	Clash
1 D 100 DHE 0E1	1 D 170 CI II II C2	distance (Å)	overlap (Å)
1:B:166:PHE:CE1	1:B:170:GLU:HG3	2.12	0.84
1:B:167:TRP:O	1:B:172:VAL:HG23	1.89	0.71
1:C:118:SER:OG	1:C:121:VAL:HG23	1.91	0.71
1:B:172:VAL:HG13	1:B:224:LEU:CD2	2.23	0.68
1:C:208:LEU:HD12	1:C:209:ASN:N	2.09	0.67
1:C:91:HIS:HB3	1:C:154:GLU:HG3	1.77	0.66
1:C:156:LEU:HD13	1:C:188:ASN:HD22	1.61	0.65
1:B:172:VAL:HG13	1:B:224:LEU:HD21	1.79	0.65
1:B:215:PRO:HB2	1:B:233:LEU:HB2	1.79	0.65
1:C:19:GLU:OE1	1:C:21:ARG:HD2	1.97	0.64
1:B:227:ILE:HD12	1:B:227:ILE:O	1.98	0.63
1:B:194:VAL:HG13	1:B:218:TYR:OH	1.98	0.63
1:B:108:GLY:O	1:B:112:VAL:HG23	2.01	0.60
1:B:38:ALA:HB2	1:B:66:THR:HG23	1.83	0.60
1:B:77:MET:HG2	1:C:68:TRP:CH2	2.38	0.58
1:B:38:ALA:CB	1:B:66:THR:HG23	2.32	0.58
1:C:113:LYS:HG2	4:C:303:Q9F:O24	2.05	0.57
1:C:165:PRO:O	1:C:169:GLU:HG3	2.05	0.57
1:C:172:VAL:HB	1:C:173:PRO:HD3	1.85	0.57
1:C:172:VAL:HG13	1:C:224:LEU:CD2	2.35	0.56
1:C:123:PRO:HB2	1:C:124:PRO:HD2	1.87	0.56
1:B:83:ARG:HD3	1:C:78:TRP:O	2.06	0.56
1:C:116:ARG:HD2	4:C:303:Q9F:C02	2.36	0.56
1:C:172:VAL:HG13	1:C:224:LEU:HD22	1.87	0.55
1:B:171:ILE:O	1:B:175:ILE:HG13	2.07	0.54
1:C:215:PRO:HB2	1:C:233:LEU:HB2	1.90	0.54
1:B:168:ASN:HA	1:B:172:VAL:HG21	1.88	0.54
1:B:74:ILE:HD12	1:B:76:GLN:NE2	2.23	0.53
1:C:149:GLN:O	1:C:151:PRO:HD3	2.09	0.52
1:B:166:PHE:CD1	1:B:170:GLU:HG3	2.46	0.51
1:C:52:PHE:CE1	1:C:182:LEU:HB2	2.45	0.51
1:C:142:TYR:O	1:C:145:LEU:HB2	2.12	0.50
1:C:85:TRP:NE1	1:C:145:LEU:HD11	2.26	0.50
1:B:220:LEU:HD23	1:B:226:PRO:HA	1.92	0.50
1:B:119:TYR:CE1	1:B:157:LYS:HB2	2.48	0.49
1:B:172:VAL:HB	1:B:173:PRO:HD3	1.94	0.49
1:C:227:ILE:O	1:C:228:LYS:HB3	2.12	0.49
1:B:53:ASP:OD1	1:B:180:ARG:NE	2.38	0.48
1:B:85:TRP:NE1	1:B:145:LEU:HD21	2.28	0.48
1:B:168:ASN:HA	1:B:172:VAL:CG2	2.44	0.48
1:B:223:ASN:O	1:B:224:LEU:HB2	2.14	0.48
1:B:203:GLU:H	1:B:203:GLU:CD	2.18	0.47
	121201020102		



 $Continued\ from\ previous\ page...$

Continuea from prev		Interatomic	Clash
Atom-1	Atom-2	distance (Å)	overlap (Å)
1:B:95:LEU:HD22	1:B:103:THR:HG23	1.96	0.47
1:C:142:TYR:HB3	1:C:145:LEU:HD22	1.96	0.47
1:C:15:ALA:O	1:C:18:LEU:HB2	2.15	0.47
1:C:89:GLU:HG3	1:C:90:ARG:N	2.29	0.47
1:B:227:ILE:HD12	1:B:227:ILE:C	2.35	0.46
1:B:115:TRP:CZ2	1:B:124:PRO:HD3	2.50	0.46
1:C:177:GLU:O	1:C:177:GLU:HG3	2.15	0.46
1:C:120:ASP:HA	1:C:155:SER:HB2	1.98	0.46
1:B:21:ARG:HA	1:B:98:LEU:O	2.16	0.46
1:B:160:ILE:HD13	1:B:192:GLY:HA2	1.98	0.45
1:C:58:SER:HB2	1:C:88:ASN:OD1	2.16	0.45
1:B:89:GLU:HG3	1:B:90:ARG:N	2.32	0.45
1:B:202:GLU:O	1:B:205:ILE:HG22	2.17	0.45
1:C:237:GLU:HG2	1:C:238:THR:N	2.32	0.44
1:B:230:MET:HE3	1:B:232:PHE:HZ	1.82	0.44
1:C:123:PRO:HB2	1:C:124:PRO:CD	2.48	0.44
1:B:119:TYR:CD1	1:B:157:LYS:HB2	2.52	0.44
1:B:227:ILE:HD11	1:B:228:LYS:HD3	2.00	0.44
1:B:59:VAL:HG22	1:B:85:TRP:HA	2.00	0.43
1:C:93:GLY:O	1:C:96:THR:OG1	2.27	0.43
1:B:190:LEU:O	1:B:194:VAL:HG23	2.18	0.43
1:C:116:ARG:HH11	1:C:117:ARG:CZ	2.32	0.43
1:C:172:VAL:HB	1:C:173:PRO:CD	2.49	0.43
1:C:202:GLU:O	1:C:205:ILE:HG22	2.19	0.43
1:B:161:ALA:O	1:B:165:PRO:CD	2.67	0.42
1:C:167:TRP:CZ2	1:C:197:LEU:HD11	2.54	0.42
1:B:230:MET:CE	1:B:232:PHE:HZ	2.32	0.42
1:B:43:GLN:O	1:B:44:ALA:C	2.58	0.42
1:B:121:VAL:HA	1:B:122:PRO:HD3	1.92	0.42
1:B:230:MET:CE	1:B:232:PHE:CZ	3.02	0.42
1:B:152:SER:O	1:B:153:CYS:HB3	2.20	0.41
1:B:77:MET:O	1:C:64:ILE:HD13	2.19	0.41
1:B:161:ALA:O	1:B:165:PRO:HD3	2.20	0.41
1:B:18:LEU:HA	1:B:21:ARG:HB2	2.02	0.41
1:B:200:LEU:HD12	1:B:205:ILE:HD13	2.03	0.41
1:B:164:LEU:HA	1:B:164:LEU:HD23	1.72	0.41
1:B:233:LEU:HD23	1:B:233:LEU:HA	1.59	0.41
1:C:59:VAL:HG12	1:C:83:ARG:HB3	2.03	0.41
1:C:58:SER:HA	1:C:84:THR:O	2.21	0.41
1:C:119:TYR:HE2	1:C:157:LYS:HG3	1.85	0.41
1:B:56:PHE:CD1	1:B:82:VAL:HB	2.56	0.40



Continued from previous page...

Atom-1	Atom-2	Interatomic	Clash	
Atom-1	Atom-2	$\operatorname{distance} \left(\operatorname{\AA} \right)$	overlap (Å)	
1:C:52:PHE:CD1	1:C:182:LEU:HB2	2.56	0.40	
1:C:205:ILE:HD12	1:C:205:ILE:HA	1.85	0.40	
1:B:113:LYS:O	1:B:117:ARG:HG3	2.22	0.40	

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Favoured Allowed		Percentiles	
1	В	232/262 (88%)	218 (94%)	13 (6%)	1 (0%)	34	66
1	С	235/262 (90%)	224 (95%)	10 (4%)	1 (0%)	34	66
All	All	467/524 (89%)	442 (95%)	23 (5%)	2 (0%)	34	66

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	С	20	ASN
1	В	153	CYS

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	В	199/220 (90%)	193 (97%)	6 (3%)	41 75	



Continued from previous page...

Mol	Chain	Analysed	Analysed Rotameric Outliers		Percentiles		
1	С	201/220 (91%)	199 (99%)	2 (1%)	76 93		
All	All	400/440 (91%)	392 (98%)	8 (2%)	59 84		

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	В	5	LYS
1	В	118	SER
1	В	147[A]	GLU
1	В	147[B]	GLU
1	В	206	MET
1	В	209	ASN
1	С	14	SER
1	С	222	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	С	188	ASN
1	С	209	ASN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

Of 3 ligands modelled in this entry, 1 is monoatomic - leaving 2 for Mogul analysis.



In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol T	Tuno	Type Chain		Link	Bond lengths			Bond angles		
MIOI	Type	Chain	Res	Lilik	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	MES	С	302	-	12,12,12	2.23	1 (8%)	14,16,16	1.88	4 (28%)
4	Q9F	С	303	-	36,36,36	1.79	7 (19%)	52,52,52	2.58	9 (17%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	MES	С	302	-	-	5/6/14/14	0/1/1/1
4	Q9F	С	303	-	-	11/23/23/23	0/4/4/4

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(A)	Ideal(A)
3	С	302	MES	C8-S	-7.28	1.67	1.77
4	С	303	Q9F	S07-N06	5.97	1.73	1.63
4	С	303	Q9F	C09-S07	3.96	1.82	1.76
4	С	303	Q9F	O21-S07	3.76	1.47	1.43
4	С	303	Q9F	O08-S07	3.48	1.47	1.43
4	С	303	Q9F	C05-N06	3.25	1.48	1.42
4	С	303	Q9F	C30-CL31	2.06	1.79	1.74
4	С	303	Q9F	O24-C23	2.03	1.40	1.36

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$\mathbf{Observed}(^o)$	$\operatorname{Ideal}(^{o})$
4	С	303	Q9F	O21-S07-O08	-14.88	101.26	119.55
4	С	303	Q9F	C22-C26-C27	-4.40	113.59	120.86
4	С	303	Q9F	C09-S07-N06	4.32	112.27	106.83
3	С	302	MES	C6-C5-N4	-3.52	104.77	110.10
4	С	303	Q9F	C05-C04-C23	3.27	121.46	117.48
3	С	302	MES	C5-N4-C3	3.25	116.15	108.83



Continued from previous page...

Mol	Chain	Res	Type	Atoms	${f Z}$	$Observed(^o)$	$\operatorname{Ideal}({}^{o})$
4	С	303	Q9F	C25-C26-C27	3.20	126.15	120.86
4	С	303	Q9F	O08-S07-N06	3.13	114.57	106.73
4	С	303	Q9F	C05-C04-C02	-3.04	116.42	121.05
3	С	302	MES	O2S-S-C8	2.78	110.26	106.92
4	С	303	Q9F	C26-C25-C23	-2.37	118.79	120.68
3	С	302	MES	O3S-S-O2S	-2.17	105.98	111.27
4	С	303	Q9F	C14-C13-C12	-2.03	117.84	121.36

There are no chirality outliers.

All (16) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	С	302	MES	C8-C7-N4-C5
3	С	302	MES	C7-C8-S-O1S
3	С	302	MES	C7-C8-S-O3S
4	С	303	Q9F	C25-C26-C27-C28
4	С	303	Q9F	C22-C26-C27-C28
4	С	303	Q9F	C22-C26-C27-C33
4	С	303	Q9F	C25-C26-C27-C33
4	С	303	Q9F	C19-C12-C13-C14
4	С	303	Q9F	C19-C12-C13-C18
4	С	303	Q9F	C11-C12-C13-C14
4	С	303	Q9F	C11-C12-C13-C18
3	С	302	MES	C7-C8-S-O2S
3	С	302	MES	C8-C7-N4-C3
4	С	303	Q9F	C10-C09-S07-O08
4	С	303	Q9F	O01-C02-C04-C05
4	С	303	Q9F	C20-C09-S07-O08

There are no ring outliers.


1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	С	303	Q9F	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring



in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	$\langle { m RSRZ} \rangle$	$\#\mathrm{RSRZ}{>}2$		$OWAB(A^2)$	Q < 0.9
1	В	233/262 (88%)	-0.20	2 (0%)	84 80	39, 51, 71, 82	0
1	С	237/262 (90%)	-0.24	1 (0%)	92 91	38, 52, 69, 91	0
All	All	470/524 (89%)	-0.22	3 (0%)	89 86	38, 52, 70, 91	0

All (3) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	В	147[A]	GLU	2.6
1	С	236	GLU	2.6
1	В	177	GLU	2.4

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-}\mathbf{factors}(\mathbf{\mathring{A}}^2)$	Q < 0.9
4	Q9F	С	303	33/33	0.85	0.33	58,81,100,113	0
3	MES	С	302	12/12	0.97	0.16	53,63,69,77	0



Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-}\mathbf{factors}(\mathbf{\mathring{A}}^2)$	Q < 0.9
2	CL	С	301	1/1	0.99	0.22	50,50,50,50	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers (i)

There are no such residues in this entry.

