

# Full wwPDB X-ray Structure Validation Report (i)

#### Sep 10, 2023 – 07:23 AM EDT

PDB ID	:	4IUQ
Title	:	crystal structure of SHH1 SAWADEE domain
Authors	:	Du, J.; Patel, D.J.
Deposited on		
Resolution	:	2.81  Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org* A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:


MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
$\mathrm{EDS}$	:	2.35.1
buster-report	:	1.1.7(2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber $(2001)$
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.35.1

# 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:  $X\text{-}RAY\;DIFFRACTION$ 

The reported resolution of this entry is 2.81 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$egin{array}{c} { m Whole \ archive} \ (\#{ m Entries}) \end{array}$	${f Similar\ resolution}\ (\#{ m Entries,\ resolution\ range}({ m \AA}))$
$R_{free}$	130704	3140(2.80-2.80)
Clashscore	141614	3569(2.80-2.80)
Ramachandran outliers	138981	3498 (2.80-2.80)
Sidechain outliers	138945	3500 (2.80-2.80)
RSRZ outliers	127900	3078 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain		
1	А	135	76%	21%	
1	В	135	73%	20%	•••



# 2 Entry composition (i)

There are 3 unique types of molecules in this entry. The entry contains 2220 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called SAWADEE HOMEODOMAIN HOMOLOG 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	р	129	Total	С	Ν	0	S	0	0	0
	ГБ	129	1054	655	194	199	6			
1	1 A 1	134	Total	С	Ν	0	S	0	0	0
		104	1098	681	203	208	6	0	0	0

There are 2 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
В	124	SER	-	expression tag	UNP Q9XI47
А	124	SER	-	expression tag	UNP Q9XI47

• Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	В	1	Total Zn 1 1	0	0
2	А	1	Total Zn 1 1	0	0

• Molecule 3 is CYMAL-4 (three-letter code: CVM) (formula:  $C_{22}H_{40}O_{11}$ ).






Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	В	1	Total         C         O           33         22         11	0	0
3	А	1	Total         C         O           33         22         11	0	0



# 3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: SAWADEE HOMEODOMAIN HOMOLOG 1





# 4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants	54.71Å 117.69Å $53.41$ Å	Depositor
a, b, c, $\alpha$ , $\beta$ , $\gamma$	$90.00^{\circ}$ $90.00^{\circ}$ $90.00^{\circ}$	Depositor
Resolution (Å)	40.07 - 2.81	Depositor
Resolution (A)	40.07 - 2.81	EDS
% Data completeness	98.7 (40.07 - 2.81)	Depositor
(in resolution range)	98.9(40.07-2.81)	EDS
R <sub>merge</sub>	0.10	Depositor
$R_{sym}$	0.10	Depositor
$< I/\sigma(I) > 1$	$2.33 (at 2.81 \text{\AA})$	Xtriage
Refinement program	PHENIX (phenix.refine: 1.7_650)	Depositor
$R, R_{free}$	0.201 , $0.261$	Depositor
II, Ilfree	0.200 , $0.257$	DCC
$R_{free}$ test set	420 reflections $(4.78%)$	wwPDB-VP
Wilson B-factor $(Å^2)$	46.0	Xtriage
Anisotropy	0.422	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$ , $B_{sol}(Å^2)$	0.30 , $28.9$	EDS
L-test for $twinning^2$	$<  L  > = 0.50, < L^2 > = 0.33$	Xtriage
Estimated twinning fraction	0.029 for l,-k,h	Xtriage
$F_o, F_c$ correlation	0.92	EDS
Total number of atoms	2220	wwPDB-VP
Average B, all atoms $(Å^2)$	44.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 5.25% of the height of the origin peak. No significant pseudotranslation is detected.

<sup>&</sup>lt;sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



<sup>&</sup>lt;sup>1</sup>Intensities estimated from amplitudes.

# 5 Model quality (i)

## 5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: CVM, ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bo	nd lengths	Bond angles		
	Chain	RMSZ	# Z  > 5	RMSZ	# Z  > 5	
1	А	0.87	0/1119	0.77	1/1511~(0.1%)	
1	В	0.92	1/1073~(0.1%)	0.80	0/1448	
All	All	0.89	1/2192~(0.0%)	0.79	1/2959~(0.0%)	

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	$\operatorname{Ideal}(\operatorname{\AA})$
1	В	239	GLU	CG-CD	-5.59	1.43	1.51

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
1	А	193	ARG	NE-CZ-NH2	-5.08	117.76	120.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	1098	0	1050	17	0
1	В	1054	0	1012	23	0
2	А	1	0	0	0	0
2	В	1	0	0	0	0



Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	А	33	0	40	9	0
3	В	33	0	40	11	0
All	All	2220	0	2142	58	0

Continued from previous page...

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 13.

All (58) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic	Clash
Atom-1	Atom-2	distance (Å)	overlap (Å)
1:A:133:SER:OG	1:A:135:ARG:HG2	1.58	1.03
1:B:208:ASP:HB3	1:B:209:GLN:HE22	1.34	0.93
1:A:133:SER:OG	1:A:135:ARG:CG	2.22	0.87
3:B:302:CVM:O31	3:B:302:CVM:C17	2.30	0.80
1:B:209:GLN:NE2	1:B:209:GLN:N	2.29	0.80
3:B:302:CVM:C30	3:B:302:CVM:O21	2.30	0.80
1:A:250:GLU:N	1:A:250:GLU:OE1	2.17	0.77
1:B:208:ASP:HB3	1:B:209:GLN:NE2	1.99	0.77
3:B:302:CVM:O31	3:B:302:CVM:H17	1.85	0.76
1:A:218:LEU:HD11	1:A:237:ARG:HE	1.52	0.74
3:A:302:CVM:C24	3:A:302:CVM:H192	2.17	0.74
3:A:302:CVM:H15	3:A:302:CVM:O22	1.87	0.73
3:A:302:CVM:H192	3:A:302:CVM:H24	1.71	0.73
1:A:190:GLU:O	1:A:193:ARG:HG2	1.91	0.69
1:B:209:GLN:HE21	1:B:209:GLN:H	1.39	0.67
1:B:256:PRO:O	1:B:257:GLU:CB	2.42	0.66
1:B:209:GLN:NE2	1:B:209:GLN:H	1.93	0.66
3:B:302:CVM:O21	3:B:302:CVM:H302	1.96	0.66
1:B:256:PRO:O	1:B:257:GLU:HB3	1.97	0.65
1:B:206:ARG:O	1:B:207:GLU:C	2.38	0.61
1:A:155:GLU:OE2	1:A:174:ASN:ND2	2.30	0.60
1:A:256:PRO:O	1:A:257:GLU:HB3	2.02	0.59
1:B:209:GLN:N	1:B:209:GLN:HE21	2.00	0.56
3:A:302:CVM:C27	3:A:302:CVM:O23	2.46	0.55
1:B:203:PHE:CE1	1:B:210:ALA:HB1	2.41	0.55
1:A:136:ASP:OD2	1:A:140:TYR:OH	2.29	0.51
1:B:225:HIS:ND1	1:B:226:ASP:O	2.43	0.50
3:A:302:CVM:O12	3:A:302:CVM:O21	2.30	0.50
1:A:222:ARG:HA	1:A:232:CYS:SG	2.52	0.49
1:B:150:VAL:HG22	1:B:156:LEU:CD1	2.43	0.48
3:B:302:CVM:O21	3:B:302:CVM:O25	2.29	0.48



Atom-1	Atom-2	Interatomic $distance (\lambda)$	Clash
		distance (Å)	overlap (Å)
1:B:225:HIS:CE1	1:B:230:CYS:HA	2.48	0.48
3:B:302:CVM:O31	3:B:302:CVM:O21	2.30	0.48
1:A:194:VAL:O	1:A:195:ASN:OD1	2.32	0.48
1:B:200:LEU:HD11	1:B:217:VAL:CG2	2.43	0.47
1:A:203:PHE:HD2	1:A:251:ARG:HG2	1.79	0.47
3:A:302:CVM:O22	3:A:302:CVM:O32	2.30	0.47
1:B:145:PHE:CE1	3:B:302:CVM:H102	2.49	0.47
1:B:186:VAL:HB	1:B:249:LEU:HD11	1.98	0.45
1:B:206:ARG:O	1:B:207:GLU:O	2.34	0.45
3:B:302:CVM:O31	3:B:302:CVM:O22	2.30	0.45
3:B:302:CVM:H31C	3:B:302:CVM:H111	1.58	0.45
1:B:150:VAL:HG22	1:B:156:LEU:HD12	1.99	0.44
1:A:184:ILE:HD11	1:A:256:PRO:HG3	1.99	0.44
3:A:302:CVM:O21	3:A:302:CVM:O14	2.29	0.44
3:B:302:CVM:O21	3:B:302:CVM:C26	2.65	0.44
1:A:206:ARG:HB2	1:A:209:GLN:HG3	2.00	0.43
1:B:209:GLN:NE2	1:B:209:GLN:CA	2.75	0.43
1:A:161:ARG:HD3	1:A:170:ASP:OD1	2.17	0.43
1:A:153:THR:O	1:A:153:THR:CG2	2.65	0.42
1:B:133:SER:OG	1:B:135:ARG:HG3	2.20	0.42
1:B:222:ARG:HA	1:B:232:CYS:SG	2.59	0.42
1:A:133:SER:OG	1:A:135:ARG:HG3	2.13	0.42
3:A:302:CVM:O22	3:A:302:CVM:O23	2.38	0.42
1:B:225:HIS:CG	1:B:230:CYS:HA	2.56	0.41
1:A:129:PHE:CE1	3:A:302:CVM:H32C	2.55	0.41
3:B:302:CVM:H17	3:B:302:CVM:H24	1.15	0.41
1:B:149:ARG:NH1	1:B:157:GLU:OE1	2.44	0.40

Continued from previous page..

There are no symmetry-related clashes.

### 5.3 Torsion angles (i)

#### 5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.



Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	entiles
1	А	132/135~(98%)	127~(96%)	5(4%)	0	100	100
1	В	125/135~(93%)	119 (95%)	5 (4%)	1 (1%)	19	49
All	All	257/270~(95%)	246 (96%)	10 (4%)	1 (0%)	34	66

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	В	207	GLU

#### 5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	А	120/121~(99%)	114 (95%)	6 (5%)	24 56	
1	В	115/121~(95%)	106 (92%)	9~(8%)	12 35	
All	All	235/242~(97%)	220 (94%)	15 (6%)	17 45	

All (15) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	В	126	ASP
1	В	127	LEU
1	В	132	LYS
1	В	135	ARG
1	В	156	LEU
1	В	158	VAL
1	В	163	SER
1	В	209	GLN
1	В	252	ILE
1	А	156	LEU
1	А	161	ARG
1	А	177	THR
1	А	196	VAL
1	А	207	GLU
1	А	246	SER



Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	В	209	GLN

#### 5.3.3 RNA (i)

There are no RNA molecules in this entry.

#### 5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry (i)

Of 4 ligands modelled in this entry, 2 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Mol Type Chain Res L		Type Chain Res Link Bond lengths		ths	E	Bond ang	gles		
IVIOI	туре	Unam	nes		Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z >2
3	CVM	А	302	-	35,35,35	1.16	3 (8%)	48,48,48	2.07	13 (27%)
3	CVM	В	302	-	35,35,35	1.23	5 (14%)	48,48,48	2.33	20 (41%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	CVM	А	302	-	-	9/16/64/64	0/3/3/3



Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	CVM	В	302	-	-	4/16/64/64	0/3/3/3

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Ζ	Observed(Å)	Ideal(Å)
3	А	302	CVM	C16-C15	-3.78	1.42	1.52
3	В	302	CVM	O14-C15	-2.61	1.38	1.44
3	В	302	CVM	O23-C16	-2.38	1.37	1.43
3	В	302	CVM	C16-C15	-2.37	1.46	1.52
3	В	302	CVM	C19-C15	-2.14	1.44	1.51
3	А	302	CVM	O14-C15	-2.07	1.39	1.44
3	В	302	CVM	O14-C13	-2.05	1.36	1.41
3	А	302	CVM	O12-C13	-2.02	1.36	1.40

All (33) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
3	В	302	CVM	C17-C16-C15	-6.20	96.70	110.93
3	В	302	CVM	O14-C15-C19	-6.17	91.09	106.44
3	А	302	CVM	O25-C24-C29	-5.41	98.89	110.35
3	А	302	CVM	O23-C16-C15	-4.90	96.02	109.45
3	В	302	CVM	O25-C26-C30	4.78	118.32	106.44
3	В	302	CVM	C13-O14-C15	-4.65	104.55	113.69
3	А	302	CVM	O14-C13-O12	-4.30	99.79	109.97
3	А	302	CVM	C24-O23-C16	-3.96	108.16	117.96
3	А	302	CVM	O25-C26-C27	3.82	116.63	109.69
3	А	302	CVM	C19-C15-C16	-3.81	102.24	113.33
3	В	302	CVM	O25-C24-C29	-3.68	102.55	110.35
3	А	302	CVM	C17-C16-C15	-3.68	102.50	110.93
3	В	302	CVM	O12-C1-C2	-3.56	97.09	109.56
3	А	302	CVM	O14-C13-C18	-3.34	103.27	110.35
3	В	302	CVM	C30-C26-C27	-3.27	105.36	113.00
3	А	302	CVM	C24-O25-C26	-3.01	107.78	113.69
3	В	302	CVM	O21-C17-C18	-2.77	103.94	110.35
3	В	302	CVM	C4-C6-C11	-2.74	105.57	112.11
3	В	302	CVM	C24-O23-C16	-2.71	111.25	117.96
3	В	302	CVM	O22-C18-C17	-2.63	104.27	110.35
3	В	302	CVM	O14-C13-O12	-2.62	103.78	109.97
3	В	302	CVM	C2-C3-C4	-2.52	104.70	113.62
3	В	302	CVM	O20-C19-C15	-2.51	102.68	111.29
3	В	302	CVM	C8-C7-C6	-2.41	107.59	112.15
3	А	302	CVM	C28-C27-C26	2.40	114.52	110.24



4IUQ
------

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
3	А	302	CVM	C1-O12-C13	-2.28	110.06	113.84
3	В	302	CVM	C19-C15-C16	2.23	119.82	113.33
3	А	302	CVM	C24-C29-C28	-2.18	105.45	110.00
3	В	302	CVM	C4-C6-C7	-2.18	106.90	112.11
3	В	302	CVM	O23-C24-C29	2.18	113.75	108.10
3	В	302	CVM	O25-C26-C27	-2.17	105.75	109.69
3	А	302	CVM	O23-C24-C29	-2.16	102.50	108.10
3	В	302	CVM	C13-C18-C17	2.13	114.44	110.00

Continued from previous page...

There are no chirality outliers.

Mol	Chain	Res	Type	Atoms
3	А	302	CVM	C2-C1-O12-C13
3	В	302	CVM	C17-C16-O23-C24
3	А	302	CVM	O12-C1-C2-C3
3	В	302	CVM	C3-C4-C6-C11
3	А	302	CVM	O14-C13-O12-C1
3	А	302	CVM	C3-C4-C6-C7
3	В	302	CVM	C1-C2-C3-C4
3	А	302	CVM	C1-C2-C3-C4
3	А	302	CVM	C2-C3-C4-C6
3	А	302	CVM	C3-C4-C6-C11
3	В	302	CVM	C3-C4-C6-C7
3	А	302	CVM	O25-C24-O23-C16
3	А	302	CVM	C17-C16-O23-C24

All (13) torsion outliers are listed below:

There are no ring outliers.


2 monomers are involved in 20 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	А	302	CVM	9	0
3	В	302	CVM	11	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the



average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







## 5.7 Other polymers (i)

There are no such residues in this entry.

## 5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



# 6 Fit of model and data (i)

## 6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median,  $95^{th}$  percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	$\langle RSRZ \rangle$	#RSRZ>2	$\mathbf{OWAB}(\mathrm{\AA}^2)$	Q<0.9
1	А	134/135~(99%)	-0.54	0 100 100	22, 41, 78, 103	0
1	В	129/135~(95%)	-0.50	1 (0%) 86 81	23, 38, 74, 130	0
All	All	263/270~(97%)	-0.52	1 (0%) 92 91	22, 39, 76, 130	0

All (1) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	В	208	ASP	2.7

### 6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates (i)

There are no monosaccharides in this entry.


### 6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median,  $95^{th}$  percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	$B-factors(Å^2)$	Q < 0.9
3	CVM	А	302	33/33	0.80	0.23	60, 99, 130, 134	0
3	CVM	В	302	33/33	0.84	0.27	42,111,128,130	0
2	ZN	А	301	1/1	0.97	0.09	70,70,70,70	0
2	ZN	В	301	1/1	0.98	0.14	48,48,48,48	0



The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.







## 6.5 Other polymers (i)

There are no such residues in this entry.

