Full wwPDB X-ray Structure Validation Report (i)

Nov 15, 2023 – 01:27 AM JST

PDB ID	:	6IUI
Title	:	Crystal structure of GIT1 PBD domain in complex with Paxillin LD4 motif
Authors	:	Liang, M.; Wei, Z.
Deposited on	:	2018-11-28
Resolution	:	2.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.36
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.36

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3163 (2.60-2.60)
Clashscore	141614	3518 (2.60-2.60)
Ramachandran outliers	138981	3455 (2.60-2.60)
Sidechain outliers	138945	3455 (2.60-2.60)
RSRZ outliers	127900	3104 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	132	8%, 86%, •, 10%
1	B	132	14%, 82%, •, 14%
2	C	28	7%, 75%, •, 21%
2	D	28	75%, 25%

2 Entry composition (i)

There are 2 unique types of molecules in this entry. The entry contains 2070 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ARF GTPase-activating protein GIT1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	119	916	582	156	174	4	0	0	0
1	B	113	842	531	146	161	4	0	0	0

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	639	GLY	-	expression tag	UNP Q9Z272
A	640	PRO	-	expression tag	UNP Q9Z272
A	641	GLY	-	expression tag	UNP Q9Z272
A	642	SER	-	expression tag	UNP Q9Z272
A	643	GLU	-	expression tag	UNP Q9Z272
A	644	PHE	-	expression tag	UNP Q9Z272
B	639	GLY	-	expression tag	UNP Q9Z272
B	640	PRO	-	expression tag	UNP Q9Z272
B	641	GLY	-	expression tag	UNP Q9Z272
B	642	SER	-	expression tag	UNP Q9Z272
B	643	GLU	-	expression tag	UNP Q9Z272
B	644	PHE	-	expression tag	UNP Q9Z272

- Molecule 2 is a protein called Paxillin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	C	22	159	100	24	33	2	0	0	0
2	D	21	153	94	23	34	2	0	0	0

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	255	GLY	-	expression tag	UNP B4DRY6
C	256	PRO	-	expression tag	UNP B4DRY6
C	257	GLY	-	expression tag	UNP B4DRY6
C	258	SER	-	expression tag	UNP B4DRY6
C	259	GLU	-	expression tag	UNP B4DRY6
C	260	PHE	-	expression tag	UNP B4DRY6
D	255	GLY	-	expression tag	UNP B4DRY6
D	256	PRO	-	expression tag	UNP B4DRY6
D	257	GLY	-	expression tag	UNP B4DRY6
D	258	SER	-	expression tag	UNP B4DRY6
D	259	GLU	-	expression tag	UNP B4DRY6
D	260	PHE	-	expression tag	UNP B4DRY6

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: ARF GTPase-activating protein GIT1

Chain A:



• Molecule 1: ARF GTPase-activating protein GIT1

Chain B:



• Molecule 2: Paxillin

Chain C:



• Molecule 2: Paxillin

Chain D:

4 Data and refinement statistics (i)

Property	Value	Source
Space group	I 4	Depositor
Cell constants a, b, c, α, β, γ	128.80Å 128.80Å 47.64Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	32.20 – 2.60 32.20 – 2.59	Depositor EDS
% Data completeness (in resolution range)	99.8 (32.20-2.60) 95.8 (32.20-2.59)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	1.42 (at 2.61Å)	Xtriage
Refinement program	PHENIX 1.9_1692	Depositor
R, R_{free}	0.194 , 0.226 0.200 , 0.234	Depositor DCC
R_{free} test set	615 reflections (4.99%)	wwPDB-VP
Wilson B-factor (Å^2)	84.2	Xtriage
Anisotropy	0.180	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.28 , 92.3	EDS
L-test for twinning[2]	$< \|L\| > = 0.48$, $< L^2 > = 0.32$	Xtriage
Estimated twinning fraction	0.033 for -k,-h,-l	Xtriage
F_o,F_c correlation	0.95	EDS
Total number of atoms	2070	wwPDB-VP
Average B, all atoms (Å^2)	130.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.59% of the height of the origin peak. No significant pseudotranslation is detected.*

[1]Intensities estimated from amplitudes.

[2]Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\| >5$	RMSZ	$\#\|Z\| >5$
1	A	0.21	0/930	0.36	0/1262
1	B	0.20	0/854	0.36	0/1163
2	C	0.22	0/160	0.29	0/213
2	D	0.21	0/153	0.30	0/204
All	All	0.21	0/2097	0.35	0/2842

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	916	0	934	2	0
1	B	842	0	832	3	0
2	C	159	0	147	1	0
2	D	153	0	141	0	0
All	All	2070	0	2054	5	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (5) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:748:GLN:OE1	2:C:277:LYS:NZ	2.31	0.62
1:B:702:PRO:HB2	1:B:708:ARG:HE	1.71	0.55
1:A:707:VAL:HG22	1:A:763:ILE:HD13	1.89	0.54
1:B:730:PRO:HA	1:B:731:PRO:HD3	1.85	0.46
1:A:724:GLU:O	1:A:742:LEU:HD11	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	115/132 (87%)	113 (98%)	2 (2%)	0	100	100
1	B	109/132 (83%)	108 (99%)	1 (1%)	0	100	100
2	C	20/28 (71%)	20 (100%)	0	0	100	100
2	D	19/28 (68%)	19 (100%)	0	0	100	100
All	All	263/320 (82%)	260 (99%)	3 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	102/116 (88%)	101 (99%)	1 (1%)	76	90
1	B	90/116 (78%)	90 (100%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	C	16/22 (73%)	16 (100%)	0	100	100
2	D	16/22 (73%)	16 (100%)	0	100	100
All	All	224/276 (81%)	223 (100%)	1 (0%)	91	97

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	726	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

There are no ligands in this entry.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å^2)	Q<0.9
1	A	119/132 (90%)	0.10	10 (8%) 11 7	68, 104, 198, 231	0
1	B	113/132 (85%)	0.73	18 (15%) 1 1	72, 137, 231, 271	0
2	C	22/28 (78%)	0.01	2 (9%) 9 6	75, 107, 191, 206	0
2	D	21/28 (75%)	-0.24	0 100 100	98, 152, 182, 197	0
All	All	275/320 (85%)	0.33	30 (10%) 5 3	68, 122, 219, 271	0

All (30) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	702	PRO	8.7
1	B	701	ARG	7.4
1	B	653	ASP	7.1
1	B	703	ALA	6.3
1	A	768	LYS	5.8
1	B	654	VAL	5.4
1	B	731	PRO	5.2
1	A	647	GLY	5.2
1	B	704	LEU	4.7
1	A	736	PRO	4.3
1	A	731	PRO	4.2
1	B	730	PRO	4.2
1	A	767	GLU	4.0
1	A	737	VAL	3.9
1	A	648	LEU	3.8
1	B	765	THR	3.8
1	B	650	SER	3.6
2	C	262	ALA	3.5
1	B	732	GLU	3.2
1	B	707	VAL	3.0
2	C	261	SER	2.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	728	THR	2.9
1	A	729	VAL	2.7
1	A	766	ARG	2.7
1	B	651	THR	2.5
1	B	736	PRO	2.5
1	B	655	ILE	2.4
1	B	694	MET	2.1
1	B	737	VAL	2.1
1	B	652	GLU	2.1

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

There are no ligands in this entry.

6.5 Other polymers (i)

There are no such residues in this entry.