



# Full wwPDB X-ray Structure Validation Report

May 17, 2020 – 08:56 am BST

PDB ID : 5IX8  
Title : Crystal structure of sugar ABC transport system, substrate-binding protein from *Bordetella parapertussis* 12822  
Authors : Chang, C.; Cuff, M.; Joachimiak, G.; Clancy, S.; Joachimiak, A.; Midwest Center for Structural Genomics (MCSG)  
Deposited on : 2016-03-23  
Resolution : 1.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the  symbol.

---

The following versions of software and data (see [references](#) ) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*



The reported resolution of this entry is 1.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	3398 (1.60-1.60)
Clashscore	141614	3665 (1.60-1.60)
Ramachandran outliers	138981	3564 (1.60-1.60)
Sidechain outliers	138945	3563 (1.60-1.60)
RSRZ outliers	127900	3321 (1.60-1.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	321	 2% 94% 5%
1	B	321	 3% 92% 7%

## 2 Entry composition i


There are 4 unique types of molecules in this entry. The entry contains 5558 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Putative sugar ABC transport system, substrate-binding protein.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	S	Se			
1	A	318	Total 2455	C 1566	N 426	O 454	S 2	Se 7	0	5	0
1	B	321	Total 2461	C 1571	N 426	O 455	S 2	Se 7	0	4	0

- Molecule 2 is SULFATE ION (three-letter code: SO<sub>4</sub>) (formula: O<sub>4</sub>S).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	O	S		
2	A	1	Total 5	O 4	S 1	0	0
2	A	1	Total 5	O 4	S 1	0	0
2	A	1	Total 10	O 8	S 2	0	1
2	B	1	Total 5	O 4	S 1	0	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	B	1	Total O S 5 4 1	0	0
2	B	1	Total O S 5 4 1	0	0
2	B	1	Total O S 5 4 1	0	0

- Molecule 3 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C<sub>2</sub>H<sub>6</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O 4 2 2	0	0
3	A	1	Total C O 8 4 4	0	1
3	B	1	Total C O 4 2 2	0	0
3	B	1	Total C O 8 4 4	0	1


- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	281	Total O 281 281	0	0
4	B	297	Total O 297 297	0	0

### 3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Putative sugar ABC transport system, substrate-binding protein

Chain A: 



- Molecule 1: Putative sugar ABC transport system, substrate-binding protein

Chain B: 



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	59.56Å 78.72Å 66.89Å 90.00° 104.83° 90.00°	Depositor
Resolution (Å)	24.98 – 1.60 24.98 – 1.46	Depositor EDS
% Data completeness (in resolution range)	98.9 (24.98-1.60) 85.5 (24.98-1.46)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.17 (at 1.46Å)	Xtrriage
Refinement program	PHENIX	Depositor
R, $R_{free}$	0.140 , 0.181 0.140 , 0.181	Depositor DCC
$R_{free}$ test set	4289 reflections (4.89%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	14.2	Xtrriage
Anisotropy	0.047	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.41 , 51.2	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	5558	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	17.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.10% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.55	0/2499	0.69	1/3391 (0.0%)
1	B	0.61	1/2505 (0.0%)	0.71	4/3400 (0.1%)
All	All	0.58	1/5004 (0.0%)	0.70	5/6791 (0.1%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	99	VAL	CB-CG1	-5.55	1.41	1.52

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	227	MSE	CG-SE-CE	-6.49	84.63	98.90
1	B	227	MSE	CG-SE-CE	-6.25	85.15	98.90
1	B	218	ARG	NE-CZ-NH1	5.59	123.09	120.30
1	B	349	ASP	CB-CG-OD2	-5.16	113.65	118.30
1	B	218	ARG	NE-CZ-NH2	-5.07	117.77	120.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2455	0	2482	9	0
1	B	2461	0	2489	17	0
2	A	20	0	0	0	0
2	B	20	0	0	0	0
3	A	12	0	18	0	0
3	B	12	0	18	2	0
4	A	281	0	0	3	2
4	B	297	0	0	3	2
All	All	5558	0	5007	26	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (26) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:260:GLN:NE2	4:A:602:HOH:O	2.17	0.76
1:A:275:GLU:OE2	4:A:601:HOH:O	2.07	0.72
1:B:275:GLU:OE2	4:B:501:HOH:O	2.15	0.64
1:B:93:ALA:HB2	1:B:102[B]:GLN:HG3	1.81	0.61
1:A:159[A]:GLN:HG3	1:A:296:MSE:HE2	1.90	0.53
1:B:353:GLN:NE2	4:B:502:HOH:O	2.25	0.53
1:B:159:GLN:HG3	1:B:296:MSE:HE2	1.92	0.51
1:B:41:ARG:HH12	3:B:406[A]:EDO:H21	1.75	0.50
1:A:292:PRO:HB2	1:A:295:ILE:HG22	1.92	0.50
1:B:218:ARG:NH2	4:B:508:HOH:O	2.38	0.49
1:A:93:ALA:HB2	1:A:102:GLN:HG3	1.97	0.47
1:B:64:LEU:CD1	1:B:359:THR:HG21	2.45	0.47
1:A:353:GLN:NE2	4:A:603:HOH:O	2.19	0.46
1:B:292:PRO:HB2	1:B:295:ILE:HG22	1.98	0.46
1:B:64:LEU:HD12	1:B:359:THR:HG21	1.98	0.46
1:B:57:LEU:O	1:B:88:VAL:HG22	2.18	0.43
1:B:193:TRP:CD2	1:B:194:PRO:HD3	2.54	0.43
1:B:193:TRP:CG	1:B:194:PRO:HD3	2.54	0.42
1:B:267:VAL:HB	1:B:285:LEU:HD23	2.02	0.42
1:A:140:LYS:NZ	1:A:307:GLU:OE2	2.44	0.41
1:A:267:VAL:HB	1:A:285:LEU:HD23	2.02	0.41
1:A:121:PRO:HG3	1:A:148:VAL:HB	2.02	0.41
1:B:186:PRO:HG3	1:B:215:ALA:HB1	2.02	0.41
1:B:345:TYR:O	1:B:349:ASP:HB2	2.21	0.41
1:B:48:PRO:HG3	3:B:406[B]:EDO:H11	2.02	0.41
1:B:99:VAL:CG1	1:B:103:PHE:CZ	3.04	0.40



All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:A:779:HOH:O	4:B:741:HOH:O[2_556]	1.96	0.24
4:A:812:HOH:O	4:B:579:HOH:O[2_556]	2.18	0.02

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	321/321 (100%)	312 (97%)	9 (3%)	0	100	100
1	B	323/321 (101%)	315 (98%)	8 (2%)	0	100	100
All	All	644/642 (100%)	627 (97%)	17 (3%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	258/249 (104%)	257 (100%)	1 (0%)	91	84
1	B	258/249 (104%)	257 (100%)	1 (0%)	91	84
All	All	516/498 (104%)	514 (100%)	2 (0%)	91	84

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	47	GLU
1	B	357	ILE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

14 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	EDO	A	503	-	3,3,3	0.50	0	2,2,2	0.37	0
2	SO4	B	402	-	4,4,4	0.26	0	6,6,6	0.30	0
2	SO4	B	403	-	4,4,4	0.24	0	6,6,6	0.38	0
2	SO4	A	501	-	4,4,4	0.05	0	6,6,6	0.36	0
2	SO4	B	401	-	4,4,4	0.21	0	6,6,6	0.25	0
3	EDO	A	504[B]	-	3,3,3	0.43	0	2,2,2	0.42	0
2	SO4	A	502	-	4,4,4	0.18	0	6,6,6	0.18	0
3	EDO	A	504[A]	-	3,3,3	0.35	0	2,2,2	0.54	0
2	SO4	B	404	-	4,4,4	0.28	0	6,6,6	0.65	0
3	EDO	B	405	-	3,3,3	0.76	0	2,2,2	0.36	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	SO4	A	505[B]	-	4,4,4	0.16	0	6,6,6	0.27	0
3	EDO	B	406[B]	-	3,3,3	0.44	0	2,2,2	0.24	0
3	EDO	B	406[A]	-	3,3,3	0.45	0	2,2,2	0.64	0
2	SO4	A	505[A]	-	4,4,4	0.22	0	6,6,6	0.26	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	EDO	A	503	-	-	0/1/1/1	-
3	EDO	A	504[B]	-	-	1/1/1/1	-
3	EDO	A	504[A]	-	-	0/1/1/1	-
3	EDO	B	405	-	-	0/1/1/1	-
3	EDO	B	406[B]	-	-	1/1/1/1	-
3	EDO	B	406[A]	-	-	0/1/1/1	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	504[B]	EDO	O1-C1-C2-O2
3	B	406[B]	EDO	O1-C1-C2-O2

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	406[B]	EDO	1	0
3	B	406[A]	EDO	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	311/321 (96%)	-0.20	5 (1%) 72 71	10, 16, 29, 39	0
1	B	314/321 (97%)	-0.27	9 (2%) 51 49	8, 13, 27, 43	0
All	All	625/642 (97%)	-0.23	14 (2%) 62 60	8, 15, 27, 43	0

All (14) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	359	THR	3.8
1	A	345	TYR	3.1
1	A	348	TYR	3.0
1	B	360	ALA	2.9
1	B	345	TYR	2.8
1	B	342	THR	2.5
1	A	346	ALA	2.4
1	A	338	SER	2.1
1	B	358	GLU	2.1
1	B	243	VAL	2.1
1	A	357	ILE	2.1
1	B	343	HIS	2.1
1	B	111	THR	2.1
1	B	348	TYR	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	EDO	B	406[B]	4/4	0.87	0.15	20,20,21,21	4
3	EDO	B	406[A]	4/4	0.87	0.15	19,19,19,20	4
3	EDO	B	405	4/4	0.89	0.14	20,25,29,33	0
3	EDO	A	503	4/4	0.92	0.14	26,29,30,31	0
3	EDO	A	504[A]	4/4	0.96	0.08	17,17,17,17	4
3	EDO	A	504[B]	4/4	0.96	0.08	23,23,24,24	4
2	SO4	A	502	5/5	0.99	0.15	32,33,34,35	0
2	SO4	B	403	5/5	0.99	0.04	16,17,20,23	0
2	SO4	B	404	5/5	0.99	0.04	17,21,24,26	0
2	SO4	A	501	5/5	0.99	0.05	13,13,15,17	0
2	SO4	A	505[B]	5/5	0.99	0.08	12,15,15,16	5
2	SO4	B	401	5/5	0.99	0.14	30,31,32,32	0
2	SO4	B	402	5/5	0.99	0.04	9,11,11,13	0
2	SO4	A	505[A]	5/5	0.99	0.08	15,17,17,17	5

## 6.5 Other polymers [i](#)

There are no such residues in this entry.